

# THE EFFECTS OF LOW POWER LASER ON ELECTROPHYSIOLOGICAL PARAMETERS OF SURAL NERVE IN NORMAL SUBJECTS: A COMPARISON BETWEEN 670 AND 780 nm WAVELENGTHS

M.R. Hadian\* and B. Attarbashi Moghadam

Department of Physical Therapy and Electrophysiology, Faculty of Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran

**Abstract-** There is just one study about the effects of 830 nm low power laser (LPL) on conduction velocity of sural nerve in human. Considering the fact that the sural nerve is a pure sensory nerve, therefore, for determining the effect of LPL on electrophysiological parameters of the nerve in human, this study was carried out as a base for further basic and clinical researches. Thirty eight normal volunteer men participated (20–35 yr.) in this study. LPL (670 and 780 nm) was applied on left and right sural nerves. Electrophysiological parameters such as Onset Latency (OL), Peak Latency (PL), Negative Peak Amplitude (NPA), Peak to Peak Amplitude (PPA) and Duration were measured before and after the application of various doses (0.5, 1.5 and 2.5 J/cm<sup>2</sup>) of LPL. This study showed that both wavelengths of LPL increase the latencies and therefore reduce the nerve conduction velocity (NCV). In addition, LPL application decreased the nerve amplitudes. Among the various intensities, the application of 2.5 J/cm<sup>2</sup> was the most effective ( $P < 0.001$ ). On the other hand, 670 nm wavelength and 2.5 J/cm<sup>2</sup> had the greatest effects on OL in comparison with 780 nm ( $P < 0.04$ ). However, there was no significant difference between the effects of 670 and 780 nm on the other electrophysiological parameters of sural nerve.

Acta Medica Iranica, 41(3): 138-142; 2003

**Key Words:** Low power laser, sural nerve, electrophysiological parameters

## INTRODUCTION

There is only one research about the influence of low power laser (LPL) on conduction velocity of sural nerve in human being (1) and the effects of LPL on the other electrophysiological parameters of the nerve have not been investigated. However, there are some controversial evidences about the effects of different wavelengths and doses of LPL on peripheral nerves of human; these studies were mainly performed on those nerves which contained both sensory and motor fibers. Great House et al., Walsh et al. and Bartlett et. al. reported that application of LPL had no effects on the latency of sensory branch of radial and median nerves (2- 4). However, Synder et. al., Lowe et al, Basford et. al. and Baxter et.al reported that application of LPL increased the latency of sensory branches of median nerves (5-8). Safavi showed that application of LPL increased the latency of sensory branches of radial

nerves in human. Safavi also reported that there was no significant difference between the effects of LPL on right and left radial nerves (9). Accordingly, in determining the effects of LPL on electrophysiological parameters of sural nerve in human, this study was performed as a base for further basic and clinical investigations.

## MATERIALS AND METHODS

Thirty eight normal volunteer men (20-35 yr.) participated in this study. Sensory nerve action potentials (Disa, Dentech Inc. Denmark; Frequency, 20Hz-2kHz; Sensitivity: 20 $\mu$ Volt/Division; Sweep speed: 4ms/Division; Stimulus Duration: 0.2ms) of Sural nerves were antidromically recorded, and electrophysiological parameters such as Onset Latency (OL), Peak Latency (PL), Negative Peak Amplitude (NPA), Peak to Peak Amplitude (PPA) and Duration were measured and, Conduction Velocity (CV) were calculated (10,11). 670 and 780 nm LPL (Chatanoga, USA, Gallium- Aluminum- Arsenide, Continous waves, 3 mW) at various doses (0.5, 1.5 and 2.5 J/cm<sup>2</sup>) were transcutaneously applied on the left and right

Received: 5 January 2002, accepted: 6 January 2003

### \* Corresponding Author:

M.R. Hadian, Department of Physical Therapy and Electrophysiology, Faculty of Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran  
Tel: +98 21 7536134  
Fax: +98 21 7534133  
E-mail: hadian\_ras@yahoo.com

sural nerves throughout the surface anatomical pathway. The irradiation time of laser was calculated based on the following formula

$$\text{Irradiation Time (s)} = \frac{\text{Energy Density (J/cm}^2\text{)} \times \text{Area of Irradiation (cm}^2\text{)}}{\text{Output Power (w)}}$$

(i.e. 167 seconds= 0.5 J/cm<sup>2</sup> for 670 nm and 25 seconds= 0.5 J/cm<sup>2</sup> for 780 nm) (12). Thereafter, sensory nerve action potentials (SNAPs) of the nerves were separately recorded after the application of LPL at different wavelengths and doses (e.g. 670 nm, 0.5, 1.5 and 2.5 J/cm<sup>2</sup>). Similar method was exactly repeated for 780 nm at the same doses. Throughout the experiment, room temperature was kept at 30°C and skin temperature was also monitored through the digital thermometer. For statistical analysis, SPSS for windows, version 7 (SPSS Inc., 1997); Paired t-test and Friedman Two Way ANOVA were used (13).

## RESULTS

The results of application of 670 and 780 nm LPL on electrophysiological parameters of sural nerve are compared before and after the laser irradiation at different doses (Fig. 1,2,3 and 4). For convenience the effect of 670 and 780 nm are therefore described separately.

### a) 670 nm (0.5,1.5 and 2.5 J/cm<sup>2</sup>)

Irradiation of 0.5 J/cm<sup>2</sup> laser had increased significantly OL (Paired t-test, P<0.04), (Fig. 1) and decreased significantly PPA (Paired t-test, P<0.01), (Fig. 4).

Irradiation of 1.5 J/cm<sup>2</sup> laser had increased significantly OL (Paired t-test, P<0.01), (Fig. 1) and

decreased significantly NPA & PPA (Paired t-test, P<0.01), (Fig. 3 and 4).

Irradiation of 2.5 J/cm<sup>2</sup> laser had increased significantly OL and PL (Paired t-test, P<0.001), (Fig. 1 and 2) and decreased significantly NPA and PPA (Paired t-test, P<0.001), (Fig. 3 and 4).

### b) 780 nm (0.5,1.5 and 2.5 J/cm<sup>2</sup>)

Irradiation of 0.5 J/cm<sup>2</sup> laser had increased significantly OL and PL (Paired t-test, P<0.03), (Fig. 1 and 2) and decreased significantly PPA (Paired t-test, P<0.01), (Fig. 4).

Irradiation of 1.5 J/cm<sup>2</sup> laser had increased significantly OL (Paired t-test, P<0.04), (Fig. 1) and decreased significantly NPA and PPA (Paired t-test, P<0.04), (Fig. 3 and 4).

Irradiation of 2.5 J/cm<sup>2</sup> laser had increased significantly OL and PL (Paired t-test, P<0.001), (Fig. 1 and 2) and decreased significantly NPA and PPA (Paired t-test, P<0.001), (Fig. 3 and 4).

### c) Comparison among the different doses

Nonparametric Friedman two way ANOVA (13) was used to compare the effects of different doses of 670 nm on electrophysiological parameters of the nerve. The same procedure was exactly repeated for 780 nm.

Among the different doses of 670 nanometer laser, significant difference was only seen among the mids of OL (before irradiation, after irradiation of 0.5, 1.5 and 2.5 J/cm<sup>2</sup> respectively, P<0.04); furthermore, the greatest difference was seen after 2.5 J/cm<sup>2</sup> (Fig. 1).

With regard to 780 nm, significant difference was only seen among the mids of OL (before irradiation, after irradiation of 0.5,1.5 and 2.5 J/cm<sup>2</sup> respectively, P<0.04); furthermore, the greatest difference was seen after 2.5 J/cm<sup>2</sup> (Fig. 1).




Fig. 1. Comparison of onset latencies before and after the irradiation of LPL at different doses and wavelengths

The effects of LPL on electrophysiological parameters of sural nerve




Fig. 2. Comparison of peak latencies before and after the irradiation of LPL at different doses and wavelengths




Fig. 3. Comparison of negative peak amplitudes before and after the irradiation of LPL at different doses and wavelengths




Fig. 4. Comparison of peak to peak amplitudes before and after the irradiation of LPL at different doses and wavelengths

Archive of SID

**d) Comparison among the different wavelengths**

Paired t-test was used to compare the effects of 670 and 780 nm. Application of 2.5 J/cm<sup>2</sup> had significantly increased OL (P<0.04), (Fig. 1); while, no significant differences were seen among the other parameters and doses.

**DISCUSSION**

There is no research about the influence of 670 and 780 nm laser at various doses on sensory nerve of human in the literature. The only citation is about the effects of 830 nm laser on NPL parameter of SNAPs at 0.5, 1 and 1.5 J/cm<sup>2</sup> (1). As mentioned earlier, the effects of LPL on other electrophysiological parameters of sensory nerve in human have not been investigated. The results of this study showed that the application of continuous beam of Gallium-Aluminum-Arsenide laser with 670 nm (3 mW) and 780 nm (20 mW) at maximum intensity (2.5 J/cm<sup>2</sup>, in this study) had significant effects on all electrophysiological parameters of sural nerve with the exception of the duration. These findings are in accordance with Synder et al., Lowe et al., Basford et al., Baxter et al. and Safavi who also reported that application of LPL increased the latency of sensory branches of median and radial nerves (5-9). However, Great House et al., Walsh et al. and Bartlett et al. reported that application of pulsed beam (904 and 820 nm) had no effect on latency, amplitude and NPL of sensory branch of radial nerve (2-4). This controversy might be due to the fact that these authors used pulsed beam, 904 and 820 nm wavelengths. Application of both wavelengths of the LPL had increased the OL and PL of SNAPs. This finding shows that nerve conduction is affected with laser beam. This might be due to indirect effects of laser on increment of ATP production in cells and correspondingly activation of Na<sup>+</sup> ion channels. Alternatively, bioelectric and bioenergetic effects of laser on ions transition through the cell membrane can affect cell membrane potentials. Indeed, there are some reports regarding hyperpolarization of cell membrane and increment of threshold of afferent fibers (14-17). 670 nm LPL (2.5 J/cm<sup>2</sup>) was more effective than the same dose of 780 nm laser on electrophysiological parameters of sural nerve. Lesser depth of penetration of 670 nm laser might have a greater effect on the sural nerve because it lies superficially on the lateral aspect of the ankle. The above findings might be useful in pain management (18-20). Application of LPL might cause presynaptic inhibition of those afferent fibers that signal pain to the central nervous system. However,

further studies should be performed in normal and patient subjects.

**REFERENCES**

1. Cambier D, Blom K, Witvrouw E, Ollevier G, De Muynck M, Vanderstraeten G. The influence of low intensity laser irradiation on conduction characteristics of peripheral nerve: A randomized, controlled, double blind study on sural nerve. *Lasers Med Sci* 2000; 15(3): 195–200.
2. Great House DG, Currier DP, Gilmore RL. Effects of clinical laser on superficial nerve conduction. *Physiotherapy* 1985; 65(8): 1184–1187.
3. Walsh DM, Baxter GD, Allen JM. Lack of effect of pulsed low-intensity (820 nm) laser irradiation on nerve conduction in the human superficial radial nerve. *Lasers Surg Med* 2000; 26(5): 485–490.
4. Bartlett WP, Quillen WS, Gonzalez JL. Effect of gallium aluminum arsenide triple-diode laser on median nerve latency in human subjects. *J Sport Rehabil* 1999; 8(2): 99–108.
5. Synder L, Mackler, Christopher EB. Effects of helium–neon laser irradiation on peripheral sensory nerve latency. *Physical Therapy* 1988; 68(2): 223–225.
6. Lowe A, Baxter GD, Walsh DM, Allen JM. Effects of low intensity laser (830 nm) irradiation on skin temperature and antidromic conduction latencies in the human median nerve: Relevance of radiant exposure. *Laser Surg Med* 1994; 14(1): 40–46.
7. Basford JR, Hallman HO, Matsumoto JY, Moyer SK, Buss JM, Baxter GD. Effects of 830 nm continuous wave laserdiode irradiation on median nerve function in normal subjects. *Laser Surg Med* 1993; 13(6): 597–604.
8. Baxter GD, Walsh DM, Allen JM, Lowe AS, Bell AG. Effects of low intensity laser-irradiation upon conduction in the human median nerve in vivo. *Exp Physiol* 1994; 79(2): 227–234.
9. Safavi Z. Effects of low intensity laser-irradiation upon electrophysiological parameters of the human sensory branch of radial nerve. *MSC thesis* 1997; (68).

### The effects of LPL on electrophysiological parameters of sural nerve

10. Kimura J. Electrodiagnosis in disease of muscle and nerve, principle and practice. Second ed. F A Davis 1989.
11. Oh SJ. Clinical EMG and nerve conduction studies. Second ed. Williams and Wilkins 1993.
12. Kitchen SS, Partridge CJ. A review of low level laser therapy; Part I: Background, Physiological effects and hazards. *Physiotherapy* 1991; 77(3): 161–163.
13. Siegel S. Nonparametric statistics for the behavioral sciences. Second ed. Mc Graw-hill 1988.
14. Matsushia H, Ito A, Kakami K, Fukaya M. Effect on the action potential of the low power Nd–Yag laser as irradiated directly to the nerve. *Aichi Gakkuin Sci* 1989; 2: 19–28.
15. Rochkind S, Rousso M, Nissan M, Villarreal M, Barr. Nea L, Rees DG. Systemic effects of low power laser irradiation on the peripheral and central nervous system, cutaneous wounds and burns. *Laser Surg Med* 1989; 9(2): 174–82.
16. Rochkind S, Vogler I, Barr. Nea L. Spinal cord response to laser treatment of injured peripheral nerve. *Spine* 1990; 15(1): 6–10.
17. Rochkind S, Quaknine GE. New trend in neuroscience: low power laser effect on peripheral and central nervous system (basic science, pre-clinical and clinical studies). *Neurol Res* 1992; 14(1): 2–11.
18. Shuster MA, Isaev VM, Rechitskii VI, Agafonov BV. Treatment of trigeminal neuralgia, ganglioneuritis pterigopalatine ganglion and other types of prosopalgias by helium–neon laser irradiation of the pterigopalatine ganglion. *Zh Nevropatol Psikhiatr* 1988; 88(7): 96–8.
19. Wong E, Lee G, Zucherman J, Mason DT. Successful management of female office workers with “repetitive stress injury“ or “carpal tunnel syndrome“ by a new treatment modality- application of low power laser. *Int J Clin Pharmacol Ther* 1995; 33(4): 208–211.
20. Hadian MR, Fakhari Z. The effect of low power laser on carpal tunnel syndrome. Fourth congress international federation of societies for Hand Therapy; Canada 1998; 24-26.

Archive of SID