

# ICTAL AND INTERICTAL EEG ABNORMALITIES IN 100 MIGRAINEURS WITH AND WITHOUT AURA

H. Pourmahmoodian, M. Kahani, M. Ghaffarpour, M. H. Harrirchian, M. Ghabaee\* and A. Fallah

Department of Iranian Center of Neurological Research, Imam Khomeini Hospital, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

**Abstract-** There are several conflicting reports about the EEG of the migraineurs. In this study we report the ictal and interictal EEGs of 100 migraineurs, in comparison with control group. The range age for patient and control groups were 9-48 (mean:  $26 \pm 1.8$ ) and 10-46 (mean:  $23 \pm 2.1$ ) years respectively. 32% of the patients were less than 14 years old and the remaining 68% were more than 14 years. In the patient group, 68% of cases had migraine without aura and 32% suffered from migraine with aura. Hemiplegic and basilar migraines were observed in one and two of our patients respectively. Gender and age had no effect on the type of migraine. Family history for first degree relatives was found in 64% of patients, without being influenced by gender or type of migraine. Male to female ratio was 1/1.6 (38/62). Abnormal EEG was found to be much more frequent in migraineurs than the control group (47% vs. 7%). Children had an overall somewhat more abnormal EEGs, compared with adult group (53% vs. 44% or 17/32 vs. 30/68), though slow discharges were detected more in adult group. The most common abnormality was slow high voltage waves, which was observed in 33/47 (70%) of abnormal recordings. The less common findings, in decreasing order of frequency were: focal (slow, sharps or mixed) discharges in 14/47 (29%), epileptiform (alone or associated with slow waves) in 4/47 (8.5%), diffuse beta and frontal intermittent delta, activity each being in 1/47 (2.1%) of abnormal recordings.

© 2006 Tehran University of Medical Sciences. All rights reserved.

*Acta Medica Iranica*, 45(4): 309-315; 2007

**Key words:** Migraine, Aura, Electroencephalography, Basilar migraine, Hemiplegic migraine.

## INTRODUCTION

Headache is one of the humanity's most common afflictions. The utility of EEG in the diagnosis of headache has been controversial. During the past 50 years, a variety of electroencephalographic abnormalities have been reported in patients with migraine, with an incidence ranging from 11% to 74% (1). Although EEG is not useful in the routine evaluation of headaches including migraine, but may be of benefit in those headaches which have unusual

symptoms, suggesting possible seizure disorder (2,3). It is also claimed that EEG is clearly indicated in cases with acute headache attacks, when either epilepsy, basilar migraine, migraine with prolonged aura or alternating hemiplegic migraine is suspected (4), keeping the fact in the mind that the final diagnosis must mainly depend on clinical judgment however. A few controlled and blinded studies have shown focal slow activity in 0-15% and spikes in 0.2-9% of patients with migraine, generally not different from control group (5-7, 8). However spikes are reported more common in migraineurs than in headache-free control subjects by other authors (9). Lauritzen *et al.* and Westmoreland (10, 11) reported slowing, depression of background amplitude and also normal EEG during visual aura of migraineurs, others (12-14) showed definite

Received: 9 May 2006, Revised: 31 Jul. 2006, Accepted: 6 Aug. 2006

### \* Corresponding Author:

Majid Ghaffarpour, Department of Neurology, Imam Khomeini Hospital, School of Medicine, Medical Sciences/Tehran University, Tehran, Iran

Tel: +98 21 66940033, Fax: +98 21 66940033

E-mail: icnr@sina.tums.ac.ir

unilateral intermittent or bilateral delta activity during attacks of basilar and hemiplegic migraines. De Carlo et al (15) reported usefulness of EEG, particularly in children suffering from migraine with aura during ictal phase, because they found abnormalities in 80% of cases. Gallo and Winter (16) described persistence of photic driving to 20 Hz flashes or above (H-response) in nearly 90% of patients with idiopathic and post-traumatic migraine-like headache and epilepsy, while 80% of headache-free subjects lacked a response above 14 Hz. Lack of alpha blocking during intermittent photic stimulation, a trend towards a greater driving response, and an increased alpha power asymmetry are also reported (17-21). Appearance of slow waves and disturbance of consciousness have been reported in classic migraine with different clinical presentations (global amnesia, stupor, clouding of consciousness) (22-24). Lauritzen demonstrated that reduction of cerebral blood flow is not of sufficient magnitude to explain the focal symptoms and also abnormal discharges in classic migraine, rather it may be caused by neuronal dysfunction (25).

## MATERIAL AND METHODS

This prospective cross-sectional case control study was conducted in our center within January 2004 to November 2005. 100 migraineurs, diagnosed according to international headache society (HIS) criteria and randomized from patients referred to clinic of neurology. Patients with history of epilepsy, brain lesion and recurrent headaches not compatible with HIS criteria were excluded. We excluded also drug abusers and patients with stupor during EEG recording. Equal numbers of healthy volunteers were chosen as control group. Patients less than 14 years old were considered children subgroup, and those above 14 years old as adult subgroup.

Ages, gender, family history, types of migraine as well as EEG abnormalities were evaluated. EEG

was performed, when ever possible, during headache (ictal phase) and between attacks of headaches (interictal phase). Recording were performed by a 21 channel Nehocoden machine, using the international 10-20 system. Each recording session lasted for a minimum 30 minutes, with 3 min of hyperventilation (H.V) and intermittent photic stimulation (IPS) with a flash frequency ranging from 1 to 30 Hz as described by Gallo and Winter. The EEGs were evaluated and then evaluated by another experienced physician, blinded to the patient's identities. EEGs were recorded ictally (within 48 h of onset of the headache) in 14 cases, and interictally (5-10 days after attack of migraine) in the remainder 86 patients. Although a pronounced slowing during H.V has often been considered in migraine, but this response develops also in healthy individuals, so we decided not to use the slowing during H.V in this study. Data were gathered and analyzed by SPSS software using Fisher's exact  $X^2$  test.

## RESULTS

The age range for patient and control groups were 9-48 (mean:  $26 \pm 1.8$ ) and 10-46 (mean:  $23 \pm 2.1$ ) years respectively. The ratio of children to adults and also migraine with aura (MWA) to migraine without aura (MWOA) were in both 1:2.1 (32/68).

Male to female ratio was 1:1.6 (38/62). Family history for first degree relatives was positive in 64% of patients, 21 of whom had MWA and the remaining suffered from MWOA, thus the type of migraine had no statistically significant effect on the positive family history ( $P = 0.81$ ). With regard to age-subgroups, we detected that 10 cases among the children and group 22 patients among the adults had MWA, whereas 16 cases in the children group and 52 cases of adults suffered from MWOA, indicating that age was also ineffective on the type of migraine (Table 1.  $P 0.41$ ).

**Table 1.** Frequency of MWA and MWOA in the children and adults ( $P = 0.41$ )

	Children		Adults		Total	
	No	%	No	%	No	%
MWA	10	31.2	22	68.8	32	100
MWOA	16	23.5	52	76.5	68	100

**Table 2.** Frequency of MWA and MWOA in male and females ( $P = 0.94$ )

	Male (38)		Female (62)		Total	
	No	%	No	%	No	%
MWA	12	37.5	20	62.5	32	100
MWOA	26	38.2	42	61.8	68	100

Male to female ratio in patients with MWA and MWOA were 12/20 and 26/42 respectively, thus gender also had no effect on the type of migraine (Table 2,  $P = 0.94$ ). Abnormal EEG was found much more frequently in migraineurs than the control group (47% vs. 7%) (Table 3,  $P = 0.00$ ), 18 cases with abnormal EEG had MWA and remaining 29 suffered from MWOA, suggesting that there was no relation between type of migraine and rate of abnormal EEG (Table 4,  $P = 0.20$ ). Male to female ratio in patient and control groups with abnormal EEG were 18/29 and 3/4 respectively confirming that gender also had no effect on the rate of electroencephalographic abnormalities, but age of patients was somewhat an effective factor, because children to adult ratio in patient group was 32/68

whereas it was 20/80 in control group, on the other hand 17 cases in patient group and 3 cases in control group were less than 14 years old, so with a  $P$  value of 0.029 there was a meaningful relation between the age and abnormal EEG (Table 5). Interictal EEGs were abnormal in 41% (36/86).

Slow discharges (focal, hemispheric, bilateral) were detected in 33% of patient group and 4% of control groups, thus with a  $P$  value of 0.000 it was statistically meaningful. From 33 cases with slowing, 21 patients were adults, in other words 60% of slow discharges had occurred in the adult group, thus it was also age dependent ( $P, 0.03$ ), but neither gender nor type of migraine were effective on the rate of slowing. Many of the patients had more than one type of abnormality in their EEGs.

**Table 3.** Frequency of abnormal EEG in patient and control groups ( $P = 0.000$ )


	Abnormal		Normal		Total	
	No	%	No	%	No	%
Migraineurs	47	47	53	53	100	100
Control	7	7	93	93	100	100

**Table 4.** Frequency of abnormal EEG in MWA and MWOA ( $P = 0.20$ )

	Abnormal		Normal		Total	
	No	%	No	%	No	%
MWA	18	56.2	14	43.8	32	100
MWOA	29	42.6	39	57.4	68	100

**Table 5.** Frequency of abnormal EEG in the age subgroups ( $P = 0.029$ )


	Patient group		Control group	
	No	%	No	%
Children	17(from32)	53	3(from7)	43
Adult	30(from68)	44	4(from7)	57




**Fig. 1.** Frequency of the range of age in patients with migraine and in control group.

The most common abnormality was slow high voltage waves, which was observed in 33/47 (70%) of abnormal recordings. Two-third of them was located over occipital region, ipsilateral to the side of headache. The less common findings, in decreasing order of frequency were: focal (slow, sharps or mixed) discharges in 14/47 (29%), epileptic form (alone or associated with slow waves) in 4/47 (8.5%), diffuse beta and frontal intermittent delta, each in 1/47 (2.1%) of abnormal recordings. Five of the EEGs with lateralized slowing were associated with interhemispheric asymmetry of alpha amplitude (Fig. 1-4).

Focal discharges (slow or sharps) were observed in 14% of migraineurs and in 2% of control group, which was meaningful ( $P = 0.001$ ), and located mostly over occipitotemporal region ipsilateral to the side of headache in patients with unilateral headache. We noticed that 9/14 patients with focal discharge were male, whereas this feature was 1/1 in the control group, thus masculinity had a relation with focality of the discharges ( $P = 0.029$ ), but five out of nine patients with focal discharges were less than 14 years old, so age was ineffective on the rate of focality ( $P = 0.37$ ).



**Fig. 2.** Frequency of the different types of EEG abnormalities in migraineurs



**Fig. 3.** Frequency of slow wave in patients with migraine and in control group

Epileptiform discharges were detected in only 4% of patients (8.5% of abnormal EEGs), without any relation to gender, age or type of migraine. H-response was detected in 21% of patients, and we noticed no obvious alpha blocking during IPS.

## DISCUSSION

From Welter and confusing reports, we understand that, protean and non-specific abnormalities may build up in the EEG of migraineurs. On the other hand diversity of findings may be a mine-field for uncritical physician, particularly in differentiating migraine from epilepsy, especially in children. The problem lies mainly in the high incidence of abnormalities including epileptiform discharges, found in 22-47% of children and adolescents with migraine (26, 27). Although it is doubtful that the EEG exceeds clinical judgment in separating migraine from epilepsy, in some situations however, the conditions are entwined, as in the syndromes of basilar migraine, visual phenomena, occipital spikes and seizures. Previous studies (28-30) have reported slowing with an interhemispheric asymmetry of the alpha rhythm, in the interictal (between headache attacks) EEGs of both MWO and MWOA. In our study slowing was found in 33% of patients which is nearly twice that of previous reports and only five of them (all with aura) showed obvious alpha asymmetry. Slowing was more frequent in adults and occurred both in ictal and interictal EEGs. Schoermen and Francesco (31, 32) did not report slowing during interictal phase of the migraineurs. Women had more slowing in some reports (32) but our study did not confirm this issue. Epileptic activity is the most intriguing abnormality in

migraineurs, which has been reported in 0.4% to 20% of cases (29). These abnormalities include focal and generalized high voltage spikes and spike-wave complexes suppressed by eye opening and have been observed in migraine with visual aura, basilar migraine, childhood epilepsy with occipital paroxysms and benign rolandic epilepsy (26, 34-36). Four of our patients had epileptiform discharges (8.5% of abnormalities) which correlate with the least rate of the previous studies.

To differentiate attack of migraine from epilepsy, Panayiotopoulos (37, 38) concluded that if each characteristic of visual aura is identified, the diagnosis of migraine is easy. Brincicoth (39) showed that the presence of specific clinical features (amaurosis, scotoma and positive family history) together with bilateral EEG abnormality and no changes during IPS is related to migraine. In a recent study (40) periodic lateralized epileptiform discharges (PLEDs) were reported. Our patients with epileptiform discharges did not show any PLEDs pattern. Basilar migraine is usually first encountered during childhood or teenagers. Several types of abnormal EEG were reported in basilar migraine including 1) an excess of beta activity in the ictal phase in children (41), 2) predominant delta activity during attack of headache and normal EEG between the attacks (42), 3) slowing in posterior region or slowing with spikes and sharp-wave complexes (43-45) and 4) unusual association of acute confusional state with FIRDA (frontal intermittent rhythmic delta activity) during attack of migraine (45). We had two patients suffering from basilar migraine; one of them showed diffuse beta activity with the absence of alpha rhythm during headache and the second one had frontal intermittent slow activity. We have no explanation for the latter finding.

In conclusion, our study showed that there was no significant variation between electroencephalograms of MWA and MWOA, and that interictal EEGs did not add further information to the clinical and neurological examination. We also noticed, in contrast to Francesco *et al.* (31) that, most of patients had abnormal interictal EEG (41%), and despite the fact that, abnormal EEGs were found more frequently in children, in regard with slowing, in adults it exceeded than that among children.

## Acknowledgement

The authors acknowledge the help of Dr. Hatmi. N, Consultant epidemiologist from the faculty of medicine, department of epidemiology, for providing statistical help, and Dr. Zahid Hussain Khan, Professor of Anesthesiology for editing the manuscript.

## Conflict of interests

The authors declare that they have no competing interests.

## REFERENCES

1. Sand T. EEG in migraine: a review of the literature. *Funct Neurol* 1991; 6(1): 7-22.
2. Rosenberg J, Alter M, Byrne TD, Daube JR, Franklin G, Goldstein ML et al. Practice parameter: The electroencephalogram in the evaluation of headache. Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 1995; 45:1411-1413.
3. EFNS Task Force. Neurophysiological tests and imaging procedures in headache patients. Overview and Recommendations. *Eur J Neurol* 2001; in press.
4. Gordon N. Alternating hemiplegia of childhood. *Dev Med Child Neurol* 1995;37:464-468.
5. Giel R, de Vlieger M, van Vliet AGM. Headache and the EEG. *Electroencephalogr Clin Neurophysiol* 1999, 21:492-495.
6. Whitehouse D, Pappas JA, Escala PH, Livingstone S. Electroencephalographic changes in children with migraine. *N Engl J Med* 1967;276:23-27.
7. Rowan AJ. The electroencephalographic characteristics of migraine. *Arch Neurol Biol (Madr) Suppl* 1974;37:95-113.
8. Ninck B. Migraine and epilepsy. *Eur Neurol* 1970;3:168-178.
9. Schachter SC, Ito M, Wannamaker BB, Rak I, Ruggles K, Matsuo F et al. Incidence of spikes and paroxysmal rhythmic events in overnight ambulatory computer-assisted EEGs of normal subject: a multicenter study. *J Clin Neurophysiol* 1998;15:251-255.
10. Lauritzen M, Trojaborg W, Olesen J. EEG during attacks of common and classical migraine. *Cephalalgia* 1981; 1:63-66.

11. Westmoreland BF. EEG in the evaluation of headaches. In: Klass DW, Daly DD, eds. *Current Practice of Clinical Electroencephalography*. New York: Raven Press, 1979; p. 381-394.
12. Bickerstaff ER. Basilar artery migraine. *Lancet* 1961; 1:15-17.
13. Walser H, Isler H. Frontal intermittent rhythmic delta activity, impairment of consciousness and migraine. *Headache* 1982; 22:74-80.
14. Jacome DE. EEG features in basilar artery migraine. *Am J Dis Child* 1978; 132:278-281.
15. De Carlo L, Cavaliere B, Arnaldi C, Faggioli R, Soriani S, Scarpa P. EEG evaluation in children and adolescents with chronic headaches. *Eur J Paediatr* 1999; 158: 247-248.
16. Golla FL, Winter AL. Analysis of cerebral responses to flicker in patients complaining of episodic headache. *Electroencephalogr Clin Neurophysiol* 1959; 11: 539-549.
17. Tsounis S, Varfis G. Alpha rhythm power and the effect of photic stimulation in migraine with brain mapping. *Clin Electroencephalogr* 1992; 23: 1-6.
18. Genco S, de Tomasso M, Prudenzeno AMP, Savarese M, Puca FM. EEG features in juvenile migraine: topographic analysis of spontaneous and visual evoked brain electrical activity. *Cephalalgia* 1994; 14:41-46.
19. Puca FM, de Tomasso M, Tota P, Sciruicchio V. Photic driving in migraine: correlations with clinical features. *Cephalalgia* 1996; 16:246-250.
20. Fachetti D, Marsile C, Faggi L, Donati E, Kokodoko A, Poloni M. Cerebral mapping in subjects suffering from migraine with aura. *Cephalalgia* 1990; 10:279-284.
21. Lia C, Carenini L, Degioz C, Bottachi E. Computerized EEG analysis in migraine patients. *Ital J Neurol Sci* 1995; 16:249-254.
22. Caplan L, Chedru F, Lhermitte F, Mayman C. Transient global amnesia and migraine. *Neurology* 1981; 31:1167-70.
23. Plum F, Posner JB. *Diagnosis of stupor and coma*. 3<sup>rd</sup> ed. Philadelphia: FA Davis Company, 1982:5.
24. Tinaper P, Cortelli P, Sacquegna T, Lugaesi E. Classic migraine attack complicated by confusional. State: EEG and CT study. *Cephalgia* 1985; 5:63-68.
25. Lauritzen M, Olsen J. Regional cerebral blood flow during migraine attacks by xenon-133 inhalation and emission tomography. *Brain* 1984; 107:447-461.
26. Froelich WA, Carter CC, O'Leary JL. Headache in childhood, Electroencephalographic evaluation of 500 cases. *Neurology* 1960; 639-642.
27. Pinsky AL, Sommer D. The diagnosis and treatment of migraine in children. *Neurology*, 1979; 29:306-510.
28. Puca F, de Tommaso M. Clinical neurophysiology in childhood headache. *Cephalalgia* 1999; 19:137-146.
29. De Tommaso M, Sciruicchio V, Guido M, Sasanelli G, Puca FM. EEG spectral analysis in migraine without aura attacks. *Cephalalgia* 1998; 18:324-328.
30. Kramer U, Nevo Y, Neufeld MY, Harel S. The value of EEG in children with chronic headaches. *Brain Dev* 1994; 16:304-308.
31. Schoenen J, Jamart B, Delwaide PJ. Electroencephalographic mapping in migraine during the critical and intercritical periods. *Rev Electroencephalogr Neurophysiol Clin* 1987; 17(3):289-299.
32. Francesco P, Carlo F. Ictal and interictal EEG findings in children with migraine. *J Headache and pain* 2004; 5:23-29.
33. Smyth V.O.G, winter A.L. *The EEG in migraine, Electroencephalography and clinical Neurophysiology*, Elsevier Science Ireland Ltd. 1964; Vol 16. P:194-202.
34. Froelich WA, Carter CC, O'Leary JL, Rosenbaum HE. Headache in childhood: electroencephalographic: evaluation of 500 cases. *Neurology* 1960; 10:639-642.
35. Hockady JM, Witty CWM. Factors determining the electroencephalogram in migraine: a study of 560 patients according to clinical type of migraine. *Brain* 1969; 92:769-788.
36. Panayiotopoulos CP. Benign nocturnal childhood occipital epilepsy: a new syndrome with nocturnal seizures, tonic deviation of the eyes and vomiting. *J Child Neurol* 1989; (1): 43-49.
37. Panayiotopoulos CP. Visual phenomena and headache in occipital epilepsy: a review, a systematic study and differentiation from migraine. *Epileptic Disord* 1999; (4): 205-216.
38. Panayiotopoulos CP. Elementary visual hallucinations, blindness, and headache in idiopathic occipital epilepsy: differentiation from migraine. *J Neurosurg Psychiatry* 1999; 66(4): 536-540.
39. Brinciotti M, Di Sabato ML, Matricardi M, Guidetti V. Electroclinical features in children and adolescents with epilepsy and/or migraine, and occipital epileptiform EEG abnormalities. *Clin Electroencephalogr* 2000; 31(2): 76-82.
40. Garcia-Morales I, Garcia MT, Galan Davila L, Gomez-Escalonilla C, Saiz Diaz R, Martinez-Salio A, de la Pena P, Tejerina A. Periodic lateralized epileptiform discharges: etiology, clinical aspects, seizures, and evolution in 130 patients. *J Clin Neurophysiol* 2002; 19:172-177.

41. Parain D, Samson-Dollfus C Electroencephalograms in basilar artery migraine. *Electroencephalogr Clin Neurophysiol* 1984; 58:392-399.
42. Passier PE, Vredevelde JW, de Lrom MC Basilar migraine with severe EEG abnormalities. *Headache* 1994; 34(1):56-58.
43. Ramelli GP, Sturzenegger M, Donati F, Karbowski K EEG findings during basilar migraine attacks in children. *Electroencephalogr Clin Neurophysiol* 1984; 107:374:378.
44. De Romanis F, Buzzi G, Assenza S, Brusa L, Cerbo R Basilar migraine with electroencephalographic findings of occipital apike-wave complexes: a long-term study in seven children. *Cephalalgia* 1993; 13:192-196.
45. Panayiotopoulos CP Basilar migraine? Seizures, and severe epileptic EEG abnormalities. *Neurology* 1980; 30(10): 1122-1125.
46. Pietrini V, Terzano MG, D Andrea G, Parrino L, Cananzi AR, Ferro-Milone F Acute confusional migraine: clinical and electroencephalographic aspects. *Cephalalgia* 1987; 7(1):29-37.

Archive of SID