

ASSOCIATION BETWEEN ENZYMATIC AND NON-ENZYMATIC ANTIOXIDANT DEFENSE WITH ALZHEIMER DISEASE

A. Vaisi-Raygani^{1,2,4*}, Z. Rahimi^{1,3}, M. Zahraie⁴, M. Noroozian⁵, and A. Pourmotabbed⁶

1) Department of Clinical Biochemistry, Kermanshah University of Medical Sciences, Kermanshah, Iran

2) Reproductive Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran

3) Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran

4) Department of Clinical Biochemistry, Tehran University of Medical Sciences, Tehran, Iran

5) Department of Neurology, Tehran University of Medical Sciences, Tehran, Iran

6) Department of Physiology and Pharmacology, Kermanshah University of Medical Sciences, Kermanshah, Iran

Abstract- The etiopathogenesis of dementia in Alzheimer's disease (AD) is still unclear. However, long-term oxidative stress is believed to be one of the major contributing factors in progression of neuronal degeneration and decline of cognitive function in AD. In order to assess the presence of oxidative stress in AD, we examined the enzymatic activities of the erythrocyte Cu-Zn superoxide dismutase (Cu-Zn SOD), glutathione peroxidase (GSH-Px), catalase (CAT), and plasma level of total antioxidant status (TAS) in AD and control groups (age and sex-matched). The results showed that the Cu-Zn SOD activity was significantly higher and the level of GSH-Px and TAS activities were significantly lower in AD subjects than that in the control group (2111 ± 324 U/grHb, 43.7 ± 11.6 U/grHb, and 1.17 ± 0.23 mmol/L compared with 1371 ± 211 U/gHb; $t = -2.17$, $p = 0.036$, 56.3 ± 9.5 U/gHb; $t = 3.8$, $p = 0.014$, and 1.54 ± 0.2 mmol/L; $t = 11.18$, $P < 0.001$, respectively). While, the erythrocyte CAT activity was lower in AD subjects compared to the control group, the difference was not statistically significant ($t = 1.3$, $P = 0.15$). These findings support the idea that the oxidative stress plays an important role in the pathogenesis underlying AD neurodegeneration. In addition, the enzymatic activity of the erythrocyte Cu-Zn SOD and GSH-Px and the plasma level of TAS can be used as a measure of the oxidative stress and a marker for pathological changes in the brain of patients with AD.

© 2008 Tehran University of Medical Sciences. All rights reserved.

Acta Medica Iranica, 46(1): 11-16, 2008

Key words: Alzheimer disease, oxidative stress, superoxide dismutase, glutathione peroxidase, catalase, total antioxidant status

INTRODUCTION

The etiopathogenesis of dementia in Alzheimer is still unclear (1-3).

Received: 24 Oct. 2005, Revised: 14 Mar. 2006, Accepted: 19 Apr. 2006

*** Corresponding Author:**

Asad Vaisi-Raygani, Department of Clinical Biochemistry, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Tel: +98 831 4274622, 9183331207, Fax: +98 831 4274623

E-mail: vaisira@yahoo.com

Recent findings indicate that cellular events involving oxidative stress may be a basic mechanism of neuro-degenerative disease. Severe oxidative stress progressively leads to cell dysfunction and ultimately cell death. Oxidative stress is an imbalance between pro-oxidants and/or free radicals on one hand, and anti-oxidizing systems on the other (2,3-7). Oxidative stress results from generation of oxygen free radicals, hydrogen peroxide, hydroxyl

radical, hydroperoxide, dioxygen and nitric oxide, collectively termed as reactive oxygen species (ROS). ROS is hypothesized to be main etiologic factor for progressive and specific neuronal degeneration which is observed in the Alzheimer disease (AD) (1,8,9). These ROS are highly reactive toward proteins, lipids and DNA molecules causing damage to these macromolecules and possibly leading to dysfunction or death of the cell (8,10).

There are many intrinsic free radical scavenger systems, which involve enzymatic and non-enzymatic reactions. One of the enzymatic antioxidant defense systems is copper- zinc super oxide dismutase (Cu-Zn SOD) that converts super oxide radicals to hydrogen peroxide (H_2O_2). Glutathione peroxidase (GSH-Px) and catalase (CAT) will then convert H_2O_2 to a water molecule. Cu-Zn SOD, GSH-Px, and CAT together provide the primary antioxidant defense mechanism (4,8,11,12).

The non-enzymatic antioxidant defense system includes ascorbic acid (vitamin C), α -tocopherol (vitamin E), glutathione (GSH), β -Carotene, and vitamin A. There is a balance between both the activities and intracellular levels of these antioxidants that are essential for the survival of organisms and their health (4,10,13-15). It is known that the brain bears relatively low antioxidant protection, and also contains high levels of polyunsaturated fatty acids that makes it prone to increased lipid peroxidation (3,9,16).

Recent studies indicate that amyloid beta- peptide ($A\beta$) can be neurotoxin by mechanisms involving the generation of H_2O_2 , ROS and lipid peroxidation (5,17-20). It has been suggested that the brain of the AD patients are affected by inordinate oxidative stress (21). This may be due to an increased SOD and/ or decreased GSH-Px and CAT activities, leading to elevation of H_2O_2 concentration in AD. The generated H_2O_2 is used for hydroxyl radical production via Fenton and Haber-Weiss reactions (4,6,8,12,22). Thus, identifying marker(s) that can be used as a measure of the oxidative stress associated with the pathologic changes in the brain of patients with AD is highly beneficial.

We investigated the level of the enzymatic activities of erythrocyte Cu-Zn SOD, GSH-Px, and CAT, factors involved in an antioxidant defense

mechanism, and the total antioxidant status (TAS) of plasma in AD patients.

MATERIALS AND METHODS

Our case-control study consisted of 91 AD cases (40 males and 51 females with mean age of 75 ± 10 years) with no reported family history of AD or dementia and 91 unrelated controls (40 males and 51 females with a mean age of 73.5 ± 11.5 years). AD cases were recruited from the Shariati hospital and Roozbeh neurology and psychiatry center at University of Tehran, in Tehran, Iran. The diagnosis of probable AD was based upon the NINCDS/ADRDA clinical diagnostic criteria (McKhann et al. and DSM IV 1994) (23), brain CT-scan and/or MRI. Subjects were excluded if they had a family history of dementia (more than one first-degree relative with any type of dementia). Volunteer control subjects were consisted of age/sex and ethnic background matched cases, who were spouses of clinically ascertained AD or dementia patients. These subjects were chosen based on their medical history and physical examination. Their cognitive function was assessed using Mini Mental State Examination (MMSE) (24) and they did not show signs of dementia.

A written and informed consent was obtained from the subject or his/her legal guardian, and twelve hours after fasting blood samples were collected from each subject. The samples were centrifuged at 500g for 15 min and serums were collected. To obtain packed erythrocytes, the remaining erythrocytes were washed repeatedly with an isotonic solution of NaCl (0.9%) until a colorless supernatant was observed. To obtain erythrocyte hemolysate, 500 μ l packed erythrocyte was hemolyzed by addition of four volumes of cold double distilled water. The resulting suspension was centrifuged twice to eliminate all of the cell membranes as described (25). The hemolysates were used to determine Cu-Zn SOD, GSH-Px, and CAT activity.

Erythrocyte Cu-Zn SOD and GSH-Px activities were measured using the commercially available kits (Randox. Lab. Ltd. Ireland cat No.SD125 and RS

505 respectively). Erythrocyte CAT activity was measured by the method of Aebi et al (22). Plasma TAS level was measured using commercially available kit (Randox, Lab. Ltd. Ireland cat No. NX 2332). Data were expressed as mean \pm standard deviation (SD). Two-tailed student's t-test were used to compare the data between AD and control groups. Statistical significance was assumed at the $p < 0.05$ levels. The SPSS statistical software package version 11.5 was used for all of the statistical analysis.

RESULTS

The age, gender, erythrocytes Cu-Zn SOD, GSH-Px, and CAT activities and plasma TAS concentration in AD and control groups are given in Table 1. The average age of AD subjects (75 ± 9.3 years) was slightly higher than that of the control (73.2 ± 11.8), but gender distribution was the same in both groups. The mean plasma concentration of TAS and the activity of erythrocyte GSH-Px were lower in those with Alzheimer disease than in control subjects (1.17 ± 3.5 mmol/L vs 1.54 mmol/L; $t=11.18$, $P < 0.001$ and 43.7 ± 11.6 U/gHb vs 56.3 ± 9.5 U/gHb; $t=3.8$, $P=0.014$, respectively). As shown in Table 1, the erythrocyte CAT activity was also lower in AD subjects compared to control, the difference, however, was not statistically significant ($t=1.3$, $P=0.15$).

Table 1. Oxidative stress parameters in AD patients and normal controls.

Parameters	AD	Control	t*	P*
Age	75 ± 9.3	73.2 ± 11.8	-1.15	0.26
Sex(M/F)	40/51	40/51	--	--
SOD(U/gHb)	2111 ± 324	1371 ± 211	-2.17	0.03
GSH-Px(U/gHb)	43.7 ± 11.6	56.3 ± 9.5	3.8	<0.01
CAT(U/gHb)	117 ± 15	127 ± 12	1.3	0.15
TAS(mmol/L)	1.17 ± 0.23	1.54 ± 0.2	11.18	<0.001

Values are mean \pm SD

AD=Alzheimer's disease, SOD= superoxide dismutase, GSH-Px=glutathione peroxidase, CAT=catalase, TAS=total antioxidant statuses,

*By student's - t-test.

The activity of erythrocyte SOD, on the other hand, was significantly higher in those with Alzheimer's disease than the control subjects (2111 ± 324 U/gHb vs 1371 ± 211 U/gHb; $t= -2.17$, $P=0.036$).

DISCUSSION

Oxidative stress is believed to play an important role in neuronal dysfunction and ultimately cell death (12). AD has been hypothesized to be associated with oxidative stress (4,7). In this study we have found that the activity of erythrocyte Cu-Zn SOD is substantially increased in AD patients compared with age-matched control subjects, but erythrocyte GSH-Px activity and plasma TAS concentrations are reduced. These observations suggest a role for oxidative stress in the pathogenesis of AD neuro-degeneration (1,4,16). The high level of Cu-Zn SOD and the low level of GSH-Px and CAT enzymes in AD patients may result in accumulation of H_2O_2 and excessive free radicals in their brain, leading to neuro-degeneration. This is consistent with previous studies demonstrating a decreased GSH-Px (4,12,20,26) and an increased SOD (2,4-6,12,27-29) activity in Alzheimer disease and advanced aging compared to healthy controls. A significant increase in Cu-Zn SOD activity has also been found in the brain of AD patients (11) along with a defect in the metabolism of ROS that causes neuronal apoptosis (30). It has been suggested that the systems that detoxify reactive oxygen species, including enzymatic antioxidants such as glutathione peroxidase (GSH-Px) and catalase are decreased in the hippocampus of AD, while superoxide dismutase (SOD) is increased (31). Ceballos et al (32) however, reported that activity of SOD and GSH-Px are similar in AD and the same age control groups. While Licastro et al (27) reported that the activity of GSH-Px was increased in AD patients and Marcus et al (33) reported activity of SOD was decreased in AD patients.

As shown in Table 1, we have also found a significant decrease in the TAS concentration in the AD patients compared to the control.

Antioxidant activity decreased in the presence of low level of TAS leading to a compensatory elevation of SOD activity.

The decrease of antioxidant activity in the presence of low level of TAS leading to a compensatory elevation of SOD activity. Hyper production of H₂O₂ and inadequate antioxidant enzyme activities resulted in cell damage. Also, high free radical production in AD patients leads to a rapid consumption of plasma antioxidants that is not concomitant with activation of antioxidant enzymes (4).

Our finding is in accordance with previous studies and the epidemiological data, demonstrating that antioxidants may have a beneficial effect on many age-related diseases, such as AD (16,34). Furthermore, even though studies support the hypothesis that subjects with AD are malnourished-particularly in the last phase of the disease-more recent by several researches have demonstrated lower plasma antioxidant levels in the early AD stages in well-nourished subjects (12). The plasma non-enzymatic antioxidant profile of AD patients shows that the major components of the antioxidant defense system are affected in this condition. In fact, recently an epidemiological prospective cohort study shows that high dietary intake of Vitamin C and Vitamin E may lower the risk of Alzheimer's disease (11).

Finally, the measurement of peripheral markers of oxidative stress (3,4,33,34) could be used in order to biologically assess the efficacy of antioxidant supplementation in AD and mild cognitive impairment (MCI). Recent studies indicate that amyloid beta- peptide (A β) can be neurotoxin by mechanisms involving the generation of H₂O₂, ROS and lipid peroxidation (5,17-20). β amyloid toxicity is eliminated by free radical scavengers. The ability of SOD to protect against A β cytotoxicity suggests that the activation of the pathway for generation of ROS occurs upstream of superoxide (SOD activity increased). The antioxidant enzymes CAT and GSH-Px are also associated with senile plaques (30). The ability of CAT and GSH-Px to breakdown H₂O₂ generated in response to A β is a suggestive mechanism for CAT and GSH-Px protection (35).

Some in vitro studies have observed that β -amyloid aggregated peptide, a characteristic feature of AD, is toxic to neurons in culture likely through generation of free radicals and by induction of lipid oxidation (36). Marcus et al (33) have reported that the temporal lobes of AD patients consistently showed significant differences in activity for the above studied three enzymes, suggesting that abnormalities in the antioxidant system may lead to neuronal cell death in AD. These results correlate well with the regional pathology of AD and provide additional evidence for a relationship between the development of AD and a breakdown of the antioxidant system. A causal relationship between these results and the development of AD remains to be determined. If causal, then an alteration in the gene for SOD, CAT, GSH-Px, or three may have occurred. If not causal, then the abnormalities in the antioxidant system, observed in these studies, may be due to the effects of AD. Our studies indicate that learning about the antioxidant defense mechanism in AD may lead to unraveling the role of free radicals in this neuro-degenerative disease. In addition, our data suggests that the erythrocyte and the plasma level of Cu-Zn SOD, GSH-Px, and TAS can be used as a marker for pathological changes in the brain of patients with AD.

REFERENCES

1. Apelt J, Bigl M, Wunderlich P, Schliebs R. Aging-related increase in oxidative stress correlates with developmental pattern of beta-secretase activity and beta-amyloid plaque formation in transgenic Tg2576 mice with Alzheimer-like pathology. *Int J Dev Neurosci* 2004; 22(7): 475-484.
2. Delibas N, Ozcankaya R, Altuntas I. Clinical importance of erythrocyte malondialdehyde levels as a marker for cognitive deterioration in patients with dementia of Alzheimer type: a repeated study in 5-year interval. *Clinical Biochemistry* 2002; 32: 137-141.
3. Saito A, Maier,CM, Narasimhan P, Nishi T, Song YS, Yu F, Liu J, Lee YS, Nito C, Kamada H, Dodd RL, Hsieh LB, Hassid B, Kim EE, Gonzalez M, Chan PH. Oxidative stress and neuronal death/survival signaling in cerebral ischemia. *Mol Neurobiol* 2005; 31(1-3):105-116.

4. Berr C, Richard MJ, Gourlet V, Garrel C, Favier A, Enzymatic antioxidant balance and cognitive decline in aging the EVA study. *Eur J Epidemiol.* 2004; 19(2): 133-138.
5. De Leo ME Borrello S, Passantino M, Palazzotti B, Mordente A, Daniele A, Filippini V, Galeotti T, Masullo C. Oxidative stress and over expression of manganese superoxide dismutase in patients with Alzheimer's disease, *Neuroscience Letters.* 1998; 250:173-176.
6. Maier CM, Chan PH. Role of superoxide dismutases in oxidative damage and neurodegenerative disorders. *Neuroscientist.* 2002; 8(4):323-334.
7. Veurink G, Fuller SJ, Atwood CS, Martins RN. Genetics, lifestyle and the roles of amyloid beta and oxidative stress in Alzheimer's disease. *Ann Hum Biol* 2003;30(6): 639-67.
8. Bourdel-Marchasson I, Delmas-Beauvieux MC, Peuchant E, Richard-Harston S, Decamps A, Reignier B, Emeriau JP, Rainfray M. Antioxidant defenses and oxidative stress markers in erythrocytes and plasma from normally nourished elderly Alzheimer patients. *Age Ageing.* 2001; 30(3):235-241.
9. Ozcankaya R, Delibas N. Malondialdehyde, superoxide dismutase, melatonin, iron, copper, and zinc blood concentrations in patients with Alzheimer disease: cross-sectional study. *Croat Med J.* 2002; 43(1):28-32.
10. Meydani M. Antioxidant and cognitive function, *Nutrition Reviews* 2001; 59(8): S75-S82.
11. Panter SS, Scott MD. Elevated temporal cortex superoxide dismutase in Alzheimer's Disease. *Soc Neurosci Abstr* 1991; 17: 1072.
12. Rinaldi P, Polidori MC, Metastasio A, Mariani E, Mattioli P, Cherubini A, Catani M, Cecchetti R, Senin U, Mecocci P. Plasma antioxidants are similarly depleted in mild cognitive impairment and in Alzheimer's disease. *Neurobiol Aging.* 2003;24(7): 915-919.
13. J-Mate's M, Cristina PE, Rez-Go M, Nu-Ezdecastero. Antioxidant Enzymes and Human Diseases. *Clini Bioch* 1999; 32(8): 595-603.
14. Palmer AM, Burhs MA. Selective increase in lipid peroxidation in the inferior temporal cortex in Alzheimer's disease. *Brain Res* 1994; 645:338-42.
15. Sigalov AB, Stern LJ. Enzymatic repair of oxidative damage to human apolipoprotein A-I. *FEBS Lett.* 1998; 433:196-200.
16. Aksenov MY, Aksenova MV, Butterfield DA., Geddes JW, Markesbery WR., Protein oxidation in the brain in Alzheimer's disease. *Neuroscience.* 2001; 03(2):373-383.
17. Aksenov MY, Aksenova MV, Markesbery WR, Butterfield DA. Amyloid beta-peptide (1-40)-mediated oxidative stress in cultured hippocampal neurons. Protein carbonyl formation, CK BB expression, and the level of Cu, Zn, and Mn SOD mRNA. *J Mol Neurosci.* 1998; 10(3): 181-192.
18. Aksenov MY, Aksenova MV, Carney JM, Butterfield DA. Alpha 1-antichymotrypsin interaction with A beta (1-42) does not inhibit fibril formation but attenuates the peptide toxicity. *Neurosci Lett.* 1996; 18; 217(2-3):117-120.
19. Halliwell B. Reactive oxygen species and the central nervous system, *J. Neurochem* 1992; 59: 1609-1623.
20. Sagara Y, Tan S, Maher P, Schubert D. Mechanisms of resistance to oxidative stress in Alzheimer's disease. *Biofactors.* 1998; 8(1-2): 45-50.
21. Apelt J, Lessig J, Schliebs R. Beta-amyloid-associated expression of intercellular adhesion molecule-1 in brain cortical tissue of transgenic Tg2576 mice. *Neurosci Lett.* 2002; 23;329(1): 111-115.
22. Aebi h. Catalase .in Bergmeyer H.U, editor., *Methods of enzymatic analysis Vol 11.* Verlag: Chemie Weinheim. 1974; P.673-684
23. Mckhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM, Clinical diagnosis of Alzheimer's disease: report of NINCDS-ADRDA work group under the auspices of the Department of Health and Human Services task force of Alzheimer's disease. *Neurology.* 1984; 34: 939-944.
24. Folstein MF, Folstein SE, McHugh PR. Mini-Mental state, A practical method for grading the cognitive state of patients for the clinician. *J Psychiatry Res.* 1975; 12:189-198.
25. Martin Mateo MC, Martin B, SantosBeneti M, Rabadan I. Catalase activity in erythrocytes from colon and gastric cancer patients Influence on nickel, lead, mercury, and cadmium. *Biol Trace Elem Res* 1997; 7:79-90.
26. Famulari AL, Marschoff ER, Llesuy SF, Kohan S, Serra JA, Dominguez RO, Repetto M, Reides C, Sacerdote de Lustig E. The antioxidant enzymatic blood profile in Alzheimer's and vascular diseases. Their sociation and a possible assay to differentiate demented subjects and controls. *J Neurol Sci.* 1996;15;141(1- 2): 69-78.

27. Licastro F, Pedrini S, Davis LJ, Caputo L, Tagliabue J, Savorani G, Cucinotta D, Annoni G. Alpha-1-antichymotrypsin and oxidative stress in the peripheral blood from patients with probable Alzheimer disease: a short-term longitudinal study. *Alzheimer Dis Assoc Disord.* 2001;15(1): 51-55.
28. Rossi L, Squitti R, Pasqualetti P, Marchese E, Cassetta E, Forastiere E, Rotilio G, Rossini PM, Finazzi-Agro A. Red blood cell copper, zinc superoxide dismutase activity is higher in Alzheimer's disease and is decreased by D-penicillamine. *Neurosci Lett.* 2002; 30(329):137-140.
29. Serra JA, Marschoff ER, Dominguez RO, Guareschi EM, Famulari AL, Pagano MA, de Lustig ES. Oxidative stress in Alzheimer's and vascular dementias: masking of the antioxidant profiles by a concomitant Type II diabetes mellitus condition. *J Neurol Sci.* 2004; 15;218(1-2):17-24.
30. Milton NGM. Inhibition of catalase activity with 3-amino-trizole enhances the cytotoxicity of the Alzheimer's amyloid- beta peptide. *Neuro Toxicology.* 2001; 22: 767-774.
31. Balazs L, Leon M. Evidence of antioxidative challenge in the Alzheimer's brain. *Neurochem Res.* 1994; 19:1131-1137.
32. Ceballos I, Merod M, Nicole A, Theverim M, Hellier G, Legrain S, Berr C. Peripheral antioxidant enzyme activities and selenium in elderly subjects and dementia of Alzheimer's type-place of the extracellular glutathione peroxidase. *Free Rad Biol Med.* 1996; 20(15):79-87.
33. Marcus DL, Thomas C, Rodriguez C, Simberkoff K, Tsai JS, Strafaci JA, Freedman ML. Increased peroxidation and reduced antioxidant enzyme activity in Alzheimer's disease. *Exp Neurol.* 1998; 150(1): 40-44.
34. Mecocci P, Polidori MC, Cherubini A, Ingegneri T, Mattioli P, Catani M, Rinaldi P, Cecchetti R, Stahl W, Senin U, Beal MF. Lymphocyte oxidative DNA damage and plasma antioxidants in Alzheimer disease. *Arch Neurol.* 2002; 59(5):794-798.
35. Ramassamy C, Averill D, Beffert U, Theroux L, Lussier-Cacan S, Cohn JS, Christen Y, Schoofs A, Davignon J, Poirier J. Oxidative Insults Are Associated with Apolipoprotein E Genotype in Alzheimer's Disease Brain. *Neurobiology of Disease.* 2000; 7: 23-37
36. Behl, C., Davis, J. B., Lesley, R., & Schubert, D. Hydrogen peroxide mediates amyloid b protein toxicity. *Cell.* 1994; 77: 817-827.