Dendritic Cell Stimulation by IFN-β Alters T Cell Function via Modulation

of Cytokine Secretion in Diabetes Type 1

Saeid Abediankenari¹, Maryam Ghasemi^{2*}, and Mohammad Bagher Eslami³

¹ Department of Microbiology and Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

² Department of Pathology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran ³ Department of Pathobiology, School of Health, Tehran University of Medical Sciences, Tehran, Iran

Department of Famorology, School of Health, Fental Oniversity of Mealear Sciences, Fental, Ha

Received: 26 May 2008; Received in revised form: 17 Jun. 2008; Accepted: 1 Jul. 2008

Abstract- During antigen capture and processing, mature dendritic cells (DC) express large amounts of peptide-MHC complexes and accessory molecules on their surface. We investigated the role of IFN- β in induction HLA-G expression on the monocyte derived DC and cytokine profile in diabetes type 1. We accomplished secretary pattern and total cytokine production of the Th1 cytokine (IL-2, γ IFN) and Th2 cytokines (IL-4, IL-10) before and after mixed leukocyte reaction (MLR) of 30 diabetic patients and 30 normal subjects. In this study a significant increase of IL-10 and γ IFN reduction after IFN- β Therapy in culture in presence of HLA-G bearing DC as compared to control were seen. It is seen that dendritic cell causes IL-10 production of T cell in vitro that reduce T cell activation from diabetes patients and normal subjects resulted to the production and expression of HLA-G on these cells from both groups. Using mixed leukocyte reaction, it was found that IFN- β -treated dendritic cell mediated the inhibition of autologous T cell activation via IL-10 production and level of HLA-G on dendritic cell may be correlated to disease activity in diabetes patients and it could also serve as a useful marker for disease progress and treatment.

© 2009 Tehran University of Medical Sciences. All rights reserved.

Acta Medica Iranica 2009; 47(5): 341-347.

Key words Cytokine, HLA-G, IFN- β (Interferon β), diabetes type 1, autoimmunity, dendritic cell

Introduction

Diabetes type 1 is associated with an imbalance between T helper (Th) 1 and Th2 cytokines (1,2). The pathophysiological mechanism initiating this autoimmune response remains to be determined. Human studies suggested that cytokines play important roles in initiating the pathogenesis of type 1 diabetes. Cytokines are associated with destruction of pancreatic cells and development of diabetes type 1 (3). In diabetes type 1, the Th1/Th2 ratio varied and characterized by a lower production of the Th2 cytokines IL-4 and IL-10 and a higher secretion of Th1 cytokines IFN-y and IL-2 (4-6). It demonstrated that IFN-y stimulate antigen presenting cell (APC) to produce inflammatory monokines to destroy beta cells (7,8). The important mechanisms of how T lymphocytes are activated and the tolerance to the beta cells antigen is diminished remain to be elucidated. However, an exact event in the antigen recognition and activation is presentation of antigen by APC to T cells.

Variety of positive as well as negative regulatory signals is provided by APC in particular by dendritic cells (9-11). Interferon-beta (IFN- β), a type I Interferon, is normally induced by viruses or ds RNA (12) which is considered an immunomodulatory molecule for dendritic cells (13). The immune response is naturally regulated by various mechanisms aimed at controlling hyperactivity and preventing self destruction. An association between HLA and type 1 diabetes is generally accepted (9,10,14). HLA-G a non-classical MHC class I antigen is characterized by a limited polymorphism and is encoded by an altered native transcription of spliced mRNAs resulting in at least seven different isoforms, including membrane bound HLA-G1, G2, G3, G4 and HLAG5 (formerly HLAG15), G6 (formerly HLA-G25), and G7 proteins (15). These were first detected on extravillus cytotrophoblast cells (16). These antigens play an important role in cancer, pregnancy and transplantation .Moreover functional studies have identified HLA-G as a key mediator in immune tolerance (17-20).

^{*}Corresponding Author: Maryam Ghasemi

Department of Pathology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

Tel: +98 911 3553194, Fax: +98 151 3543248, E-mail: a.zghasemi@yahoo.com

Dendritic cell stimulation by IFN-β alters T cell function

Whereas, dendritic cell might have tolerogenic function even in their mature form, we studied effect of IFNbeta with GM-CSF (Granulocyte- Monocyte colony stimulating factor) and IL-4 (Interleukin 4) on DCs in regulation of T cells.

In this study we showed that IFN-beta-GM-CSF-IL-4-LPS activated dendritic cells mediate inhibition of T cell activation by producing IL-10 and IL-4 in MLR. Thus this cell may be involved in the immune regulatory pathway in the pathogenesis of type I diabetes.

Patients and Methods

The patients were those admitted in the endocrinology department of Shariati Hospital, a teaching hospital affiliated to Tehran University of Medical Sciences. The diagnosis of illness was made, according to clinical and laboratory findings, by consultant endocrinologist. The patients who took part in this study were informed clearly of the nature of this investigation and they gave the consultant endocrinologist a written consent for taking blood sample for measuring certain variables as well as using the blood cells for the present study .The blood samples which served as normal control were obtained from blood transfusion services.

To study the immune phenotype and cell surface expression of monocyte derived DC bearing HLA-G, 50ml blood samples were collected from 30patients with diabetes type 1, with median age 29.3 and 30 normal controls with median age 30.76. All diabetic patients were under treatment with insulin and showed a mean fasting blood sugar of 162 mg/dl (Table 1).

Blood mononuclear cells (MNC) were isolated from heparinzed blood by centrifugation on a Ficoll Histipaque 1.077 (sigma, U.S.A). The cells from the interphase were collected and washed three times with RPMI 1640 medium (sigma). Cell viability was determined by trypan blue exclusion. Monocytes 90% isolated from MNC by plastic flasks adherence. $5-8 \times 10^6$ /ml of MNC were then cultured in 5ml RPMI 1640 medium supplemented with 10% fetal calf serum, 50units/ml penicillin and 50µg/ml streptomycin. After 2h of incubation at 37°C in an atmosphere of 5% CO2, the adherent cells were cultured in complete medium containing 1000 units /ml rhGM-CSF (Serotec, UK), 500unit/ml rhIL-4 (serotec, UK) and IFN- β (1000 units/ml, Avonex; Biogen, Cambridge) to generate DC.

Non adherent cells were Harvested for T cell isolation. The T cells (> 95%) were isolated. CD^{3+} T cells were purified respectively using MACS CD^{3+} magnetic beads, CD^{3+} T cell isolation kit (Miltenyi Biotec). These cells were counted and then frozen in 10% dimetylsulfoxide containing a mixture of 60% RPMI and 30% FCS.

We stimulated DCs with 50μ g/ml insulin (Actrapid Novo Nordisk, Bagsvared, Denmark) on day 4 for antigen processing by these cells for 72h at 37°C, in 5% CO₂.

The immature monocyte-derived DCs were then activated by adding TNF- α (10ng/ml) into the culture to promote maturation of these cells and HLA-G expression.

Expressions of molecular markers on the immune cells were determined by flowcytometry on a fluorescent activated cell sorter (B.D).

Harvested cells were washed twice with PBS supplemented with 1% BSA. Fc receptors on cells were preblocked with excess human IgG (Sigma-Aldrich) on ice for 15 min. Cells were stained for 30 min at 4°C with the following FITC-conjugated Abs: anti-HLR-DR and anti-CD83 and PE-conjugated Abs: CD1a (eBioscience) or isotype controls (eBioscience)

Finally in order to identity and enumerate the cells, the cell suspensions were analyzed by flowcytometry.

Total RNA was extracted from treated DCs with IFN β and from jeg-3 coriocarcinoma cell line which was used as a positive control for HLA-G expression (National cell bank of country), using RNAzol Bee reagent (Biosite, TABY, Sweden) based on the manufacture instructions. RNA was checked for Quality by electrophoresis in 1.5% agarose gel. The transcriptional levels of HLA-G were evaluated by reverse transcriptase PCR in comparison of those of jeg-3 cells.

First strand cDNA was synthesized using $3\mu g$ of total RNA in 20 μ l reaction mixture consisting of 5X reaction buffer 4μ l; 10mM dNTPs 2μ l; 100 μ M dithiotheritol (DTT) 1.5 μ l; 10pMol/ml random hexamer(N6)1 μ l, and M-MLV reverse transcriptase 200units (metabion). Thereafter, the mixture was incubated at 42°C for 45 min.

Table 1. Features of diabetic patients and normal subjects

 who participated in the study

who participated in the study					
Features	Diabetes	Normal			
Total Numbers	30	30			
Gender	M(19),F(11)	M(25),F(5)			
Age in years(Range)	28.5(13-49)	28.86(22-44)			
Insulin dependent	+	-			
Familial History of	+	-			
Diabetes					
Fasting Blood Sugar	160.6 ± 57.4	88.6 ± 9.3			

Primers used in this study are 5-TCATGCTGAGA TGGAAGCAG-3; 5-TCTCCACAGCACAGCCAGC-3 for HLA-G and 5-TGGCCACGGCTGCTTCCAGC-3; CAGGAGGAGCAATGATCTTGAT-3 for beta actin.

Briefly, 25µl reaction mixture of PCR was prepared using 2.5 μ l of 10 × buffer , 3 μ l of 25mM mgcl2 , 1.5 μ l dNTPs (10Mm), 10 pmol of each primer and 1 unit of Amplify Tag Gold DNA polymerase(Perkin Elmer Biosystems, USA)) . PCR was performed in 35 cycles, initiated by 1 cycle at 95°C for activating the tag Gold's DNA polymerase followed by 94°C; 30sec,60°C ; 30sec, 72°C; 30sec and final extension of 10minutes at 72°C. Amplification of β actin was performed using 26 cycles.PCR amplification products and 100bp molecular weight markersVIII (Roche) was separated on 1% agarose gels containing etidium bromide. The primers for HLA-G produced PCR amplification products of 118bp, whereas the β actin primers gave a product of 321bp.Control PCR amplifications without cDNA were always carried out in parallel were consistently negative. All PCR amplification was performed in duplicate.

Then, isolated T cells and autolagous DC were co cultured in 96 U-shaped–bottomed plates that 10×10^4 T cell were cultured in each well with 1×10^4 DC in a final volume of 200µl per well. All samples were run in triplicates .Cultures were incubated at 37°C in a humidified 5% CO2 atmosphere for 5 days and then pulsed with 200 µl 3-[4,5-dimethylthiazolyl]-2,5-diphenyl-tetrazolium bromide(MTT; Sigma) as a color indicator of metabolic activity , denoting cell proliferation (21). Supernatant harvested at 4h later, and then dimethylsulfoxide (DMSO) was added (200 µl). The color change was read in an Elisa reader at 550nm wave length.

Cytokines concentrations were determined in duplicate in serum and culture supernatants using commercially available human IL-2, IL-10, IL-4 and γIFN Bender MedSystems Immunoassay Kit (Bender Med-Systems GmbH Campus Vienna Biocenter 2 A-1030 Viena,Austria Europe). Serum- free supernatant of Dc-Tcell cultures with and without HLA-G were harvest 5 days of incubation and frozen down for subsequent cytokine analysis.

Statistical analysis

Mature Dc

For statistical analysis, we used paired *t*-tests. Each paired *t*-test was calculated for diabetic patients as well as the normal donors .The *P* values were determined in all cases and they were considered statistically significant at P<0.05.

Results

Phenotype analysis of monocyte-derived DCs was carried out using flowcytometry.

Analysis of phenotype showed the expression of HLA-DR and minimal level of CD-83 on their surface before stimulation with Ag.Expression of the CD83 marker and HLA-DR after antigenic stimulation indicating the maturation of dendritic cells (Figure 1).

The functional potential of mature monocyte-derived DC to induce proliferation of T lymphocytes was investigated by MLR in presence or absence of HLA-G. Proliferation of T lymphocytes co-cultured with mature DC was found to decrease by about 40% in presence of DC bearing HLA-G (Figure 2, Table 2).



Immature D C

Figure 1. Flowcytometry results of immature and mature dendritic cells treated with IFN-beta in comparison with controls

Acta Medica Iranica, Vol. 47, No. 5 (2009) 343

Dendritic cell stimulation by IFN-ß alters T cell function

DC bearing HLA-G in MLR inhibited Tcell proliferation in the normal subjects as well as diabetic patients. Since we compared the T cell proliferation responses in DC expressing HLA-G and in DC lacking HLA-G, upon addition of MTT, T- Cell proliferation was inhibited in co-culture system containing DC bearing HLA-G in both diabetic as well as normal subjects as indicated by metabolic activity assay (MTT) (Table 3, Figure 3).

Interferon- γ , IL-2, IL-10 and IL-4 cytokines release assays were carried out in MLR supernatant and serum using commercially available Elisa kits. The results are illustrated in table 4.



Figure 2. RT-PCR analyses of HLA-G and β actin gene expression in diabetes type 1 and controls after treatment with IFN- β .The cDNA samples were from,: jeg-3 cell line as positive control (lane1); dendritic cell from patients with diabetes type 1 (Lane 2, 3) DCs from a normal (Lane 4, 5); negative control with out cDNA (lane 6); DNA weight marker VIII (lane 7)



Figure 3. Mean of Mixed leukocyte reaction in diabetes patients (left) and normal subjects (right) with control evaluated by MTT assay. Dendritic cells were cultured with T cells (1 to 10).

Table 2. Effect of IFN- β on expression of HLA-G on DCs in vitro as evaluated by flowcytometry. The results are percent of DC bearing HLA-G

Cells	Total Sam-	Dc without IFN-β		Dc + 1000units IFN-β		_
	ples	Mean	SD	Mean	SD	_
Diabetic DC	30	0.2	0.06	33%	0.11	< 0.05
Normal DC	30	0.5	0.15	35%	0.09	< 0.03

Table 3. Effect of DC-bearing HLA-G on CD3⁺T-Cell proliferation in MLR, evaluated by MTT assay. The data are OD of MTT at 550 nm read in Elisa reader

MLR	Total Samples	DC Without HLA-G Dc with HLA-G		P value	
	F	Mean \pm SD	Mean \pm SD	1 (1111)	
Diabetic	30	427.26±56.58	213.06±62.17	< 0.05	
Normal	30	382.36±81.86	211.3±58.9	< 0.05	

Table 4. Cytokine production profile (γ IFN, IL-2, IL-4, IL-10 [pg/ml]) in supernatant of culture media after 6 days and serum. Values are mean ±SD.

Cytokines	γIFN	IL-2	IL-4	IL-10
Supernatant of MLR(Dc without HLA-G +TCell) in Diabetes	2.39±0.53	10.4 ± 4.26	2.76±1.25	1.47±1.13
patients				
Supernatant of MLR(Dc without HLA-G +TCell) in normal	2.28 ± 0.39	9±3.8	$3.87{\pm}1.09$	1.83 ± 0.62
subjects				
Supernatant of MLR(Dc with HLA-G +TCell) in Diabetes	1.25±0.49	3.51±2.03	4.01±1.39	3.26 ± 1.51
patients				
Supernatant of MLR(Dc with HLA-G +TCell) in Normal	1.26 ± 0.34	5.2 ± 1.80	$3.89{\pm}1.24$	3.69 ± 1.30
subjects				
Serum cytokines concentration in Diabetes patients	12.56±0.44	5.01 ± 0.63	4.52 ± 0.37	1.57 ± 0.66
Serum cytokines concentration in Normals subjects	5.88 ± 1.24	6.71±0.91	5.61±1.35	4.32±1.67



Figure 4. γ IFN concentration in the supernatant of culture media in diabetes patients and normal. Dendritic cells were cultured with T cells. Values are mean \pm SD.Statistically significant at p<0.05.



Figure 5. IL-4 concentration in the supernatant of culture media in diabetes patients and normal. Dendritic cells were cultured with T cells. Values are mean \pm SD. Statistically significant at p<0.05.



Figure 6. IL-10 concentration in the supernatant of culture media in diabetes patients and normal. Dendritic cells were cultured with T cells .Statistically significant at p<0.05.

Discussion

We studied a comprehensive and follow up analysis of Th1/Th2 secretion in MLR supernatant in presence IFNbeta-treareted dendritic cell and without IFN- β in diabetes patients and controls. An important step in the present study was to induce HLA-G on dendritic cell derived from monocytes. This was achieved by treatment of dendritic cells in vitro with IFN- β . This cytokine has been shown to promote the differentiation of blood monocytes to DC and to contribute to DC maturation (22). It was also demonstrated in this study that IFN- β up regulated considerably HLA-G expression on dendritic cells (Figure 2 and Table 2). In a variety of research it has been demonstrated that DC bearing HLA-G e the potential to down regulate and to inhibit T cell activities (23-27). The cells which are inhibited by HLA-G bearing DC carry specific receptors for HLA-G. The presence of HLA-G receptors on CD4 Tcells has also been shown for immunoglobulin like transcript 2 (ILT-2) indicating this molecule a possible candidate mediating inhibitory effects (28-30). In the present study, the inhibition of Tcells proliferation was observed only when dendritic cells were treated with IFN- β (Table 3, Figure 3). Thus IFN- β and HLA-G molecules on the dendritic cells are two components contributing to this phenomenon, limiting cells proliferation; thereby reducing the potential of T-cells .We demonstrated that Dc bearing HLA-G in MLR causes higher production of IL-4, IL-10 and lower production of IL-2, IFNy in diabetes patient and controls (Figure 4, 5 and 6). The serum increased secretion of Th1 cytokines and higher Th1/Th2 ratio in our diabetes group is consistent with others, before treatment with IFN β (Figure 7). It is also interest that several evidences suggest that autoimmune diabetes is associated with decreased IL-4 secretion rather than enhanced secretion of Th1 cytokines (31). In this view IL-10 can directly inhibit Tcell activation, through inhibition of IL-2 production. Functionally; IL-10 inhibits the capacity of peripheral blood Dc to stimulate alloreactive T cells (32). Wilson and et al indicated that increased yIFN and decreased IL-4 secretion resulted a progression of type 1 diabetes (33). Our results suggest that some desirable immune modulator effects of IFN- β may be accompanished via the up regulation of the immune tolerogenic molecule of HLA-G and IL-10.

It is quite evident that the cells bearing HLA-G are an important negative immune-regulatory constituent for this suppressing effect. It is speculated that IFN- β by increasing HLA-G on dendritic cells can also down regulate inflammatory response in vivo mediated by Tcells in islet tissues of pancreatic gland. Levels of HLA-G on dendritic cells may also be correlated to disease activity in diabetes patients and it could also serve as a useful marker for disease progress and treatment. We would like to propose that HLA-G may be a factor limiting autoimmune response, thus potentially constituting a new component for controlling and suppressing the detrimental role of T-cells in the pathogenesis of diabetes type 1.

Acknowledgments

We appreciate the assistance of Dr. Tavakkoli PhD of Genetic. This study was carried out in the immunology division of the School of medicine in Mazandaran University of Medical Sciences in collaboration with Dept.of Endocrinology and Metabolic Research, Tehran University of Medical Sciences.

References

- Berman MA, Sandborg CI, Wang Z, Imfeld KL, Zaldivar F Jr, Dadufalza V, et al. Decreased IL-4 production in new onset type I insulin-dependent diabetes mellitus. J Immunol 1996;157(10):4690-6.
- 2. Rabinovitch A, Suarez-Pinzon WL. Roles of cytokines in the pathogenesis and therapy of type 1 diabetes. Cell Biochem Biophys 2007;48(2-3):159-63.
- 3. Suarez-Pinzon WL, Rabinovitch A. Combination therapy with epidermal growth factor and gastrin delays autoimmune diabetes recurrence in nonobese diabetic mice transplanted with syngeneic islets. Transplant Proc 2008;40(2):529-32.
- 4. Kallmann BA, Lampeter EF, Hanifi-Moghaddam P, Hawa M, Leslie RD, Kolb H. Cytokine secretion patterns in twins discordant for Type I diabetes. Diabetologia 1999;42(9): 1080-5.
- 5. Liu D, Cardozo AK, Darville MI, Eizirik DL. Doublestranded RNA cooperates with interferon-gamma and IL-1 beta to induce both chemokine expression and nuclear factor-kappa B-dependent apoptosis in pancreatic beta-cells: potential mechanisms for viral-induced insulitis and betacell death in type 1 diabetes mellitus. Endocrinology 2002;143(4):1225-34.
- Radu DL. Cellular mechanisms involved in experimental insulin-dependent diabetes mellitus. Roum Arch Microbiol Immunol 2001;60(3):203-26.
- Rabinovitch A. Immunoregulatory and cytokine imbalances in the pathogenesis of IDDM. Therapeutic intervention by immunostimulation? Diabetes 1994;43(5):613-21.
- Castano L, Eisenbarth GS. Type-I diabetes: a chronic autoimmune disease of human, mouse, and rat. Annu Rev Immunol 1990;8:647-79.
- Azam A, Eisenbarth GS. Immunopathogenesis and immunotherapeutic approaches to type 1A diabetes. Expert Opin Biol Ther 2004;4(10):1569-75.
- 10. Roep BO. The role of T-cells in the pathogenesis of Type 1 diabetes: from cause to cure. Diabetologia 2003;46(3):305-21.
- 11Wilson DB. Immunology: Insulin auto-antigenicity in type 1 diabetes. Nature 2005;438(7067):E5.
- 12. Biron CA. Interferons alpha and beta as immune regulators: a new look. Immunity 2001;14(6):661-4.
- 13. Karp CL, Biron CA, Irani DN. Interferon beta in multiple sclerosis: is IL-12 suppression the key? Immunol Today 2000;21(1):24-8.

- 14. Jun HS, Yoon JW. The role of viruses in type I diabetes: two distinct cellular and molecular pathogenic mechanisms of virus-induced diabetes in animals. Diabetologia 2001;44(3):271-85.
- 15. Le Rond S, Le Maoult J, Créput C, Menier C, Deschamps M, Le Friec G, et al. Alloreactive CD4+ and CD8+ T cells express the immunotolerant HLA-G molecule in mixed lymphocyte reactions: in vivo implications in transplanted patients. Eur J Immunol 2004;34(3):649-60.
- 16. Solier C, Aguerre-Girr M, Lenfant F, Campan A, Berrebi A, Rebmann V, et al. Secretion of pro-apoptotic intron 4retaining soluble HLA-G1 by human villous trophoblast. Eur J Immunol 2002;32(12):3576-86.
- 17. Selmani Z, Naji A, Zidi I, Favier B, Gaiffe E, Obert L, et al. Human leukocyte antigen-G5 secretion by human mesenchymal stem cells is required to suppress T lymphocyte and natural killer function and to induce CD4+CD25highFOXP3+ regulatory T cells. Stem Cells 2008;26(1):212-22.
- Bukur J, Rebmann V, Grosse-Wilde H, Luboldt H, Ruebben H, Drexler I, et al. Functional role of human leukocyte antigen-G up-regulation in renal cell carcinoma. Cancer Res 2003;63(14):4107-11.
- Créput C, Durrbach A, Menier C, Guettier C, Samuel D, Dausset J, et al. Human leukocyte antigen-G (HLA-G) expression in biliary epithelial cells is associated with allograft acceptance in liver-kidney transplantation. J Hepatol 2003;39(4):587-94.
- 20. Fainardi E, Rizzo R, Melchiorri L, Vaghi L, Castellazzi M, Marzola A, et al. Presence of detectable levels of soluble HLA-G molecules in CSF of relapsing-remitting multiple sclerosis: relationship with CSF soluble HLA-I and IL-10 concentrations and MRI findings. J Neuroimmunol 2003;142(1-2):149-58.
- 21. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. J Immunol Methods 1983;65(1-2):55-63.
- 22. Severa M, Remoli ME, Giacomini E, Ragimbeau J, Lande R, Uzé G, et al. Differential responsiveness to IFN-alpha and IFN-beta of human mature DC through modulation of IFNAR expression. J Leukoc Biol 2006;79(6):1286-94.

- 23. Baştürk B, Karakayali F, Emiroğlu R, Sözer O, Haberal A, Bal D, et al. Human leukocyte antigen-G, a new parameter in the follow-up of liver transplantation. Transplant Proc 2006;38(2):571-4.
- 24. Takeuchi S, Furue M. Dendritic cells: ontogeny. Allergol Int 2007;56(3):215-23.
- 25. Reder AT, Genç K, Byskosh PV, Porrini AM. Monocyte activation in multiple sclerosis. Mult Scler 1998;4(3):162-8.
- 26. Wiendl H, Mitsdoerffer M, Hofmeister V, Wischhusen J, Bornemann A, Meyermann R, et al. A functional role of HLA-G expression in human gliomas: an alternative strategy of immune escape. J Immunol 2002;168(9):4772-80.
- 27. Lila N, Rouas-Freiss N, Dausset J, Carpentier A, Carosella ED. Soluble HLA-G protein secreted by allo-specific CD4+ T cells suppresses the allo-proliferative response: a CD4+ T cell regulatory mechanism. Proc Natl Acad Sci U S A 2001;98(21):12150-5.
- 28. Chang CC, Ciubotariu R, Manavalan JS, Yuan J, Colovai AI, Piazza F, et al. Tolerization of dendritic cells by T(S) cells: the crucial role of inhibitory receptors ILT3 and ILT4. Nat Immunol 2002;3(3):237-43.
- 29. Colonna M, Samaridis J, Cella M, Angman L, Allen RL, O'Callaghan CA, et al. Human myelomonocytic cells express an inhibitory receptor for classical and nonclassical MHC class I molecules. J Immunol 1998;160(7):3096-100.
- Hunt JS, Petroff MG, McIntire RH, Ober C. HLA-G and immune tolerance in pregnancy. FASEB J 2005;19(7):681-93.
- Falcone M, Yeung B, Tucker L, Rodriguez E, Krahl T, Sarvetnick N. IL-4 triggers autoimmune diabetes by increasing self-antigen presentation within the pancreatic Islets. Clin Immunol 2001;98(2):190-9.
- 32. Buelens C, Willems F, Delvaux A, Piérard G, Delville JP, Velu T, et al. Interleukin-10 differentially regulates B7-1 (CD80) and B7-2 (CD86) expression on human peripheral blood dendritic cells. Eur J Immunol 1995;25(9):2668-72.
- 33. Wilson SB, Kent SC, Patton KT, Orban T, Jackson RA, Exley M, et al. Extreme Th1 bias of invariant Valpha24JalphaQ T cells in type 1 diabetes. Nature 1998;391(6663):177-81.