# Failure to Thrive Severity Determination by New Design

## **Curves in Standard Growth Charts**

Firouzeh Hosseini<sup>\*1</sup>, Babak Borzouei<sup>\*2</sup>, and Mehrangiz Vahabian<sup>2</sup>

<sup>1</sup> Department of Pediatrics, Besat Hospital, Hamedan University of Medical Sciences, Hamadan, Iran <sup>2</sup> Faculty of Medicine, Hamedan University of Medical Sciences, Hamadan, Iran

Received: 5 Aug. 2010; Received in revised form: 18 Dec. 2010; Accepted: 10 Jan. 2011

**Abstract-** Failure to thrive (FTT) is a perplexing disorder commonly seen by primary care physicians. Identification of FTT and an assessment of severity of the undernutrition are important to identify children at risk, and to provide appropriate intervention for preventing malnutrition and developing sequela. One of different anthropometric methods for categorizing undernutrition in childhood is Waterlow and Gomez classifications. Unfortunately, these classifications are not an easy task for the practitioners due to its difficult calculations. In this paper, Waterlow and Gomez classifications have been shown on the Standard Growth Charts (WHO 2006). Using these curves, the practitioners can determine severity of FTT by plotting weight, height and age of the patients without any calculations.

© 2011 Tehran University of Medical Sciences. All rights reserved. *Acta Medica Iranica*, 2011; 49(12): 795-800.

Keywords: Failure to thrive (FTT); Anthropometric methods; Standard growth charts

## Introduction

Failure to thrive is a perplexing disorder. The diagnosis is often in the eve of the beholder, laboratory tests are usually unrevealing and the effectiveness of treatment is uncertain (1). While definition of FTT varies widely across studies with criteria ranging from weight < 5th percentile to < 3rd percentile, a fall down 2 percentile lines, less than 80% of normal weight and a low thrive index (2). Until now, five different anthropometric methods for categorizing undernutrition in childhood were applied to children with failure to thrive (3). Between these methods, Waterlow and Gomez classifications have been used to more than others by pediatricians all over the world (4,5). FTT severity determination is very important for primary care physicians because they have different protocols for management and treatment of these children. For example, management of less severe FTT takes place at home but historically, the patients with severe FTT should be hospitalized, until the cause of the FTT is determined (6). In addition, many of primary care physicians have difficulty in using standard growth charts. Key to diagnosing FTT is finding the time in

busy clinical practice to accurately measure and plot a child's weight, height and then assess the trend (7).

In a report from England, 54% of general practitioner didn't diagnose FTT, while a child's weight for age fell below two major percentile lines (8). In addition, a pilot study performed at a family practice residency clinic found that from 29 children diagnosed with FTT, all of them had a growth charts that were incorrectly plotted and diagnosis was delayed in 41% of the patients (9). FTT severity determination may be more difficult because of its hard calculation. In this paper, Waterlow and Gomez classifications have been shown on the WHO standard growth charts 2006.

#### **Patients and Methods**

Comparing Standard Growth Charts (NCHS Revision 2000) to New Growth Charts (WHO 2006); the latter was selected because NCHS growth reference was based on data from several samples of children from a single country and suffers from a number of technical and biological drawbacks that make it inadequate to monitor the rapid and changing rate of early childhood growth (10).

**Corresponding Authors:** 

Tel: +98 811 8380462, 918 8153600, Fax: +98 811 8380208, E-mail: babakborzouei@yahoo.com

\* These authors equally contributed as corresponding authors

Firouzeh Hosseini: Department of Pediatrics, Besat Hospital, Hamedan University of Medical Sciences, Hamadan, Iran Tel: +98 811 8242427, Fax: +98 811 2640064, E-mail: hosseini@umsha.ac.ir

Babak Borzouei: Faculty of Medicine, Hamedan University of Medical Sciences, Hamadan, Iran

Original information regarding weight, height and weight for height/length under 60 months of New Growth Charts (WHO 2006) were selected .Age was monthly considered for this article. 1 year meaning 12 completed months, 2 years meaning 24 completed months, 3 years meaning 36 completed months, 4 years meaning 48 completed months, 5 years meaning 60 completed months, 18 months meaning 1 year and 6 completed months, and 30 months meaning 2 years and 6 completed months.

In Gomez classification, mild, moderate, and severe FTT is equivalent to 75-90%, 60-74% and less than 60% of standard weight, respectively, and in Waterlow classification, for height, the corresponding valves are 90-95%, 85-89% and less than 85%, and for the weight for height, the valves are 81-90%, 70-80% and less than 70% (5,11,12). Getting the information regarding 50th percentile of weight, height and weight for height of New Growth Charts (WHO 2006) and based on the definition of FTT severity (Gomez and Waterlow classifications) the following calculations were performed: 90%, 75%, and 60% of ideal weight for age (50th percentile or the median weight for age) for both gender, 95%, 90%, and 85% of ideal height for age (50th percentile or the median height for age) for both genders, and 90%, 81%, and 70% of ideal weight for height (50th percentile or the median weight for height) for both genders. Then the curves were drawn.

## Results

The curves for boys and girls age birth to 60 months were generated for length/height for age, weight for age, weight for length (45 to 110 cm) and weight for height (65 to 120 cm). After dividing 50th percentile of weight for age under 60 months for both genders to %90, %75 and %60 results in weight for age (wasting) FTT severity curves (Figures 1 and 2).

After dividing 50th percentile of length for age under 24 months for both genders to 95%, 90%, and 85% results in length for age (stunting) FTT severity curves. After dividing 50th percentile of height for age 24 to 60 months for both genders to 95%, 90% and 85% results in height for age (stunting) FTT severity curves. It was possible to construct both length for age (birth to 2 years) and height for age (2 to 5 years) standards fitting a unique model, yet still reflect the average difference between recumbent length and standing height (Figures 3 and 4).

After dividing 50th percentile of weight for length from 45 to 110 cm for both genders to 90%, 81% and 70% results in weight for length FTT severity curves (Figures 5 and 6).

After dividing 50th percentile of weight for height from 65 to 120 cm for both genders to 90%, 81% and 70% results in weight for height FTT severity curves (Figures 7 and 8).



Figure 1. Weight for age (wasting) FTT severity curves for boys age birth to 5 years



Figure 2. Weight for age (wasting) FTT severity curves for girls age birth to 5 years



Figure 3. Length/ height for age (stunting) FTT severity curves for boys age birth to 5 years



Figure 4. Length/ height for age (stunting) FTT severity curves for girls age birth to 5 years

Acta Medica Iranica, Vol. 49, No. 12 (2011) 797



Figure 5. Weight for length FTT severity curves for boys age birth to 2 years



Figure 6. Weight for length FTT severity curves for girls age birth to 2 years



Figure 7. Weight for height FTT severity curves for boys age 2 to 5 years



Figure 8. Weight for height FTT severity curves for girls age 2 to 5 years

#### Discussion

FTT in the young infant and toddler must be considered a medical emergency if the growth curve documents weight < 70% of the predicted weight for length/height (6). Rudolf and Logan concluded that FTT is associated with slight decrement in IQ-3 points and their weight and height is more significantly affected at three years follow up (2). In general, weight gain is used as the measure for identifying babies and young children who are failing to thrive, and it is generally assumed that the more severely underweight the worse the prognosis (3). Growth data were presented in variety of forms including standard deviation scores, mean centile position, percent of normal weight/height, and weight for height (2). Recently it has been argued that clinicians should use the new charts which identify the slowest growing children while correcting for regression to the mean (13). Although the different methods of FTT assessment have been compared in several articles and accordingly each has its own advantages and disadvantages and none is better than the others (3,14,15). But in a single citation by Olsen, they recommended that no single measurement seems adequate on its own for identifying nutritional growth delay (16). Waterlow and Gomez classifications seem to classifications. be more practical than other Unfortunately, FTT severity determination with each other different methods is not an easy task for the practitioners due to its difficult calculations. For using WHO Standard Growth Charts to determine the severity of FTT, a practitioner has to do the following processes:

1- Recording information regarding gender, age, weight and height of a patient.

2- Determine ideal weight (median) and ideal height (median) for age of the patient.

3- Dividing ideal weight, ideal height by percents described by Gomez and Waterlow classifications.

4- Comparing this information with each other and determining the severity of FTT. However, using the curves recommended by the authors of the present article, the practitioners need to perform only 2 processes:

1- Recording patients' information (gender, age, weight and height).

2- Comparing this information with the new curves.

Using these curves, practitioners can determine the severity of FTT by simply plotting weight, height, and age of the patients under 60 months in less time and with no calculating error. In conclusion, FTT severity determination is very essential for the primary general practitioners and the pediatricians. Using these curves is highly recommended specially in developing countries.

#### Acknowledgements

We are grateful to Dr Cynthia Ogden for her guideline to take original information.

#### References

- 1. Bauchner H. Atoms. Arch Dis Child 2005;90:881a.
- Rudolf MC, Logan S. What is the long term outcome for children who fail to thrive? A systematic review. Arch Dis Child 2005;90(9):925-31.

- Raynor P, Rudolf MCJ. Anthropometric indices of failure to thrive. Arch Dis Child 2000;82:364-5
- Waterlow JC, Buzina R, Keller W, Lane JM, Nichaman MZ, Tanner JM. The presentation and use of height and weight data for comparing the nutritional status of groups of children under the age of 10 years. Bull World Health Organ 1977;55(4):489-98.
- Bauchner H. Failure to thrive. In: Behrman RE, Kligman RM, editors. Nelson Textbook of Pediatrics. Philadelphia: WB Saunders; 2004. p. 133-4.
- Block RW, Krebs NF; American Academy of Pediatrics Committee on Child Abuse and Neglect; American Academy of Pediatrics Committee on Nutrition. Failure to thrive as a manifestation of child neglect. Pediatrics 2005;116(5):1234-7.
- Krugman SD, Dubowitz H. Failure to thrive. Am Fam Physician 2003;68(5):879-84.
- 8. Batchelor JA. Has recognition of failure to thrive changed? Child Care Health Dev 1996;22(4):235-40.
- 9. Krugman SD, Jablonski KA, Dubowitz H. Missed opportunities to diagnose failure to thrive in a family medicine resident practice. Pediatr Res 2000;47:204A.

- 10. WHO Multicentre Growth Reference Study Group. WHO Child Growth Standards based on length/height, weight and age. Acta Paediatr Suppl 2006;450:76-85.
- Gómez F, Ramos-Galvan R, Frenk S, Cravioto JM, Chávez R, Vazquez J, et al. Mortality in second and third degree malnutrition. J Trop Pediatr 1956;2:77-83.
- 12. Waterlow JC. Classification and definition of proteincalorie malnutrition. Br Med J 1972;3(5826):566-9.
- 13. Cole TJ. Conditional reference charts to assess weight gain in British infants. Arch Dis Child 1995;73(1):8-16.
- Wright JA, Ashenburg CA, Whitaker RC. Comparison of methods to categorize undernutrition in children. J Pediatr 1994;124(6):944-6.
- 15. Wright C, Avery A, Epstein M, Birks E, Croft D. New chart to evaluate weight faltering. Arch Dis Child 1998;78(1):40-3.
- 16. Olsen EM, Petersen J, Skovgaard AM, Weile B, Jørgensen T, Wright CM. Failure to thrive: the prevalence and concurrence of anthropometric criteria in a general infant population. Arch Dis Child 2007;92(2):109-14.