

Assessing the Fauna of Aquatic Insects for Possible Use for Malaria Vector Control in Large River, Central Iran

Mansoureh Shayeghi¹, Jalil Nejati¹, Leila Shirani-Bidabadi¹, Mona Koosha¹, Mehdi Badakhshan¹,
Mulood Mohammadi Bavani¹, Kourosh Arzamani¹, Nayyereh Choubdar¹, Fatemeh Bagheri¹, Abedin Saghafipour²,
Arshad Veysi¹, Fateh Karimian¹, Amir Ahamd Akhavan¹, and Hassan Vatandoost¹

¹ Department of Medical Entomology and Vector Control, School of Public Health,
Tehran University of Medical Sciences, Tehran, Iran

² Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of
Medical Sciences, International Campus (TUMS-IC), Tehran, Iran

Received: 27 Jul. 2014; Accepted: 24 Dec. 2014

Abstract- Insects with over 30,000 aquatic species are known as very successful arthropods in freshwater habitats. Some of them are applied as biological indicators for water quality control, as well as the main food supply for fishes and amphibians. The faunistic studies are the basic step in entomological researches; the current study was carried out emphasizing on the fauna of aquatic insects in Karaj River, northern Iran. A field study was carried out in six various sampling site of Karaj River during spring 2013. The aquatic insects were collected using several methods such as D-frame nets, dipping and direct search on river floor stones. Specimens were collected and preserved in Ethanol and identified by standard identification keys. Totally, 211 samples were collected belonging to three orders; Plecoptera, Trichoptera and Ephemeroptera. Seven genera (Perla, Isoperla, Hydropsyche, Cheumatopsyche, Baetis, Heptagenia and Maccaffertium) from five families (Perlidae, Perlodidae, Hydropsychidae, Batidae, Heptageniidae) were identified. The most predominant order was Plecoptera followed by Trichoptera. Karaj River is a main and important river, which provides almost all of water of Karaj dam. So, identification of aquatic species which exist in this river is vital and further studies about systematic and ecological investigations should be performed. Also, monitoring of aquatic biota by trained health personnel can be a critical step to describe water quality in this river. Understanding the fauna of aquatic insects will provide a clue for possible biological control of medically important aquatic insects such as Anopheles as the malaria vectors.

© 2015 Tehran University of Medical Sciences. All rights reserved.

Acta Med Iran 2015;53(9):523-532.

Keywords: Aquatic insects; Faunistic; Karaj River; Iran

Introduction

Insects are known as very successful arthropods in freshwater and brackish water habitats. The success of aquatic insects to exploit of these habitats can be confirmed by their diversity and abundance in most types of freshwater resources. Over 30,000 aquatic insect species have been identified that exist in freshwater; in contrast, only several hundred are marine (1).

The larval stage of mayflies (Ephemeroptera), dragonflies and damselflies (Odonata), stoneflies (Plecoptera), alderflies (Megaloptera), lacewings (Neuroptera), flies (Diptera), caddiesflies (Trichoptera),

moths (Lepidoptera) and wasps (Hymenoptera) are known as residents of freshwater, although the adults are terrestrial. Also, some species of beetles (Coleoptera) and bugs (Hemiptera) are aquatic but which their larval, nymphal and adult stages exist in the water which called fully aquatic (2).

Some orders such as Ephemeroptera, Plecoptera and Trichoptera (EPT), are known as an environmental condition indicator in water resources. In the United States, the number of EPT species is often applied as a “biological indicator” of water quality (3).

One of the environmental topics that the world faces today is the water crisis (4). Industrial activities and urbanization have raised pollution in rivers, streams, and

Corresponding Author: H. Vatandoost

Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran
Tel: +98 21 88951393, Fax: +98 21 88951393, E-mail address: vatando@tums.ac.ir

lakes in developing countries. Most sewage enters waterways without adequate treatment, and the water quality is being degraded (5).

The use of aquatic insects is a standard method and an efficient, fast, and inexpensive technique of management of freshwater resources, monitoring of aquatic biota in water recourses by trained health personnel can be a critical step to describe water quality in developing countries (6). For instance, Trichoptera species are suitable and important for monitoring physicochemical effects, and are suggested for widespread use in biomonitoring programs in developed countries (7,8).

In addition to “biological indicator” role, in freshwater ecosystems, aquatic insects have another important function in nutrient recycling and decomposition (9). These abilities are due to various functions of their feeding such as shredders, scrapers, filter feeders and predators. Also the role of some aquatic insects such as Trichopteran larvae has been proven in trophic dynamics and energy flow (6). Moreover, they are the main food supply for fishes and amphibians (2).

Although, in Iran, published faunistic studies about aquatic insects are rare, however the results indicate the richness and biodiversity of fauna. Some researchers have identified and reported the various families of orders Odonata, Ephemeroptera, Hemiptera, Trichoptera, Coleoptera and Diptera from Iran (6,10-17). Also, the aquatic beetle fauna of Iran was nearly known. Fifty one species belonging to 40 genera and 14 families from the Esat Azerbaijan province have been reported. Similar studies have been conducted in Fars, Guilan, Mazandaran, Ardabil and Khuzestan Provinces. Dytiscidae, Gyrinidae, Haliplidae, Hydrophilidae and Noteridae had the most abundance (18-23). Furthermore, in the new study which was done on Iranian Simulidae, 23 nominal species were identified (24).

Due to the importance of freshwater insects as

mentioned previously, and few faunistic studies in Iran, this study was carried out with a focus on the fauna of aquatic insects in Karaj River.

Materials and Methods

Study area

This study was conducted on Karaj River located in Karaj province in the north of the central plateau of Iran. This river with 67 km in length is situated in 36°11'N 51°35'E, comes from the Alborz Mountains and leads to Amir-Kabir (Karaj) Dam (25,26). Some studies have mentioned 450 × 106 m³ for the annual mean flow rate at its entrance to the Dam (27). Karaj River has permanent flow and its higher branches due to snow contain more water. Thus, the hydrological regime of the Karaj River can be classified as Nivopluvial (snowy-rainy). This area has been formed from various stones including; sandstone, dolomite, shale, limestone, quartzite, and tuff (25) (Figure 1).

Collection methods and identification

Aquatic specimens were collected using several methods including D-frame nets, dipping and direct search on river floor stones in 6 various sampling sites during spring 2013. The various habitat types were examined; full and partial sunlight, shaded, with and without vegetation, as well as in fast and slow flow. Specimens were picked up using forceps and placed in glass jars with labels and preserved in Ethanol (Figure 2). Labels indicate the sampling site and date of collection. All collected samples were identified using morphological characteristics of standard keys (2,28,29).

Results

Totally, 211 samples were collected belonging to three orders; Plecoptera, Trichoptera, and Ephemeroptera (Table1).

Table 1. Aquatic insects collected from the various sampling sites, Karaj River, 2013

No	Order	Family	Genus	Total No.
1	Plecoptera	Perlidae	Perla	105
2		Perlodidae	Isoperla	1
3	Trichoptera	Hydropsychidae	Hydropsyche	30
4		Hydropsychidae	Cheumatopsyche	8
5		Batidae	Baetis	25
6	Ephemeroptera	Heptagenidae	Heptagenia	24
7		Heptagenidae	Maccafferrum	18

Seven genuses (Perla, Isoperla, Hydropsyche, Cheumatopsyche, Baetis, Heptagenia and Maccafferrum) from five families (Perlidae, Perlodidae, Hydropsychidae, Batidae, Heptagenidae) were identified.

The most predominant family and genus were Perlidae (49.7%) and Perla (49.7%) belonging to Plecoptera order. It was followed by Hydropsyche

(14.2%) from Trichoptera order.

Isoperla genus (Plecoptera: Perlodidae) has the lower population size (0.5%).

The identified species are listed in Table 1. Also the captured nymphs of Plecoptera and Ephemeroptera are shown in figures 3 and 4, respectively. Figures 2 and 5 show Trichopteran eggs, larvae and case.



Figure 1. (a, b) Two sampling sites, Karaj River



Figure 2. (a) Trichopteran larva, case and eggs. (b, c, d) Bring out and pick up the larvae with forceps



Figure 3. Plecoptera: (a) Nymph. (b) Lateral gills on thoracic segments




Figure 4. Ephemeroptera: (a) Mature nymph. (b, c) Abdominal gills. (d) Caudal filaments



Figure 5. Trichoptera: (a) larvae. (b) Anal proleg and claw. (c, d) Pupal cases

Discussion

This investigation provides the first formal data with regard to the aquatic insects' fauna in a part of Karaj River as a main and important river in Iran.

In the present study, were collected various species belonging to Plecoptera. This order which commonly named the stoneflies, comprise about 2000 species (30). There are many investigations on stoneflies biology in rivers. These papers describe that the adults are most abundant near streams, particularly those with a rapid flow. They lay several thousand eggs in the water and the nymphs locate on the underside of stones. According to the literatures, many species of this order breed only in cool running waters or cold lakes and the upper limit of temperature had been reported around 250 C. The nymphs' tracheal gills are not leaf-like as the May flies, but they are thread-like structures containing tracheas.

The nymphs feed on small insects, such as May fly nymphs, and possibly on the vegetable matter (diatoms) but they are desirable food for fish. After fully grown the nymphs leave the water and molt for the last time on land (30,31).

In our study, Plecoptera species were collected in spring. Also the literature review showed their nymph stage is not tolerant to the warm conditions. In warm lands they are usually limited to forested rivers (30,32).

In Iran, in a similar study, this order has been reported from the north of the country (32) but in contrast to our result, it wasn't collected in Kashan rivers in the center (31).

In a recent investigation, two families of Plecoptera were identified: Perlidae and Perlodidae. In similar research Perlodidae was reported from the north of Iran but Perlidae not collected (32).

The Perlidae was the most predominant family in our study. It has a total of 84 species in 15 genera. This

family is widespread throughout the world except Antarctica and parts of Africa. Their life cycle varies between one and three years. The adults appear in the summer, are dynamic and usually be attracted to the light traps. Perlidae is collected in cool and clear rivers and sometimes can be found in warm rivers. Their nymph can move quickly. In stagnant water, due to no water movement over their gills, they shift the body up and down to get oxygen. Also they hunt all types of invertebrates as a predator. The larvae of this family are commonly found under stones where an abundance of prey can be found. The Perlodidae also can be found in running waters under stones. They are sometimes found along the edges of cold lakes. These stoneflies have one generation per year. Most of them emerge in the spring or fall and pass the hot summer months as aestivation in eggs stage (33).

Trichoptera was another aquatic insects' order which collected from Karaj River. This order has reported from northeast of Tehran Province, Lavasan River (6) and center of Iran (31), previously. The successive project for preparation of the Trichoptera World Checklist (TWC) was done in 2009. The TWC recorded valid extant taxa including; 13,574 species, 308 subspecies and 609 genera of 47 families. According to the World Checklist database, there are 62 Trichopteran species in Iran comprising 14 families (34).

The family Hydropsychidae which was identified is a cosmopolitan group of Trichoptera. In similar results this family was collected from various parts of Iran; center: including Kashan and Isfahan (31,35), North: Mazandaran (36), Northeast: Bojnord, South and West: Bushehr and Hamedan (37). Larvae of Hydropsychidae family are easily identified by the hard sclerotized plates on each thoracic segment and vastly branched gills on ventral of abdominal segments (6). They make silken seine-like nets for collection of food such as floating tiny organic matter or drifting invertebrates (38). Like most Trichoptera larvae, hydropsychids are sensitive to changes in dissolved oxygen levels and organic pollution (39). The Biotic Index Values (RBP) of Hydropsychidae as one of the Iranian caddisflies is reported 4.0 (6).

The genus *Hydropsyche* (Trichoptera: Hydropsychidae) was identified in this study with the second most abundance. These larvae are filter-feeding and consume everything that trapped in their nets, such as algae and tiny animals (40). As well as *Cheumatopsyche* larvae are significant ecological components in freshwater lotic ecosystems all over the world that necessitate these species for accurate water-

quality assessment (41).

The order Ephemeroptera (common name may fly) which was identified in the present study is one of the most archaic of winged insect groups. In the nymphal stages, they can be found in lakes or streams under rocks. The mature nymphs come to the water surface and turn into flying insects. Unlike most insects they typically have two winged stages. It is the only insects' order that molts after getting functional wings. The first winged stage is called the subimago, which is a sub-adult stage. Then they molt to form the true adult with reproductive ability, called the imago (42).

Baetidae is a family of mayflies which was identified in this study. It has about 900 described species that are classified as one of the smallest groups of mayflies. Their breeding sites include a wide range of habitats from lakes and streams to ditches. Some species can be found in polluted streams (43,44). This family has been reported from saline habitats such as the wetlands of the Zarrineh estuary at the south of Urmia Lake in northwestern Iran (10). They feed mainly on algae in the nymphal stage as strong swimmers. Due to their good swimming, they name minnow mayfly. (43,44).

Heptageniidae (Synonym: Ecdyonuridae) is another family of Ephemeroptera collected in this study. It has over 500 species distributed in most regions of the world. Heptageniids breed generally in streams with the fast flow but some species have been observed in stagnant waters. The nymphs have a flattened shape and usually dark color (45). In consistence to our study, both of families Heptageniidae and Baetidae were reported from Kashan Rivers, the center of Iran (31).

Although the mentioned aquatic insects normally have no medically important, some of them such as Trichoptera can cause allergic responses in sensitive persons (6). Researchers have showed that some of the orders such as caddis flies are known as biological indicators for water quality assessment (6). They explained that the groups which are sensitive to pollution (Ephemeroptera, Plecoptera, Trichoptera) in contaminated areas are reduced and vice versa, the resistant groups Diptera (Chironomidae and Simuliidae) are increased (45).

Karaj River is a main and important river which provides almost all of water of Karaj (Amir-Kabir) dam. Thus, identification of aquatic species which exist in this river is vital. The lack of indigenous taxonomic knowledge and local identification keys were the limitations of the present study, although the international identification keys were available.

It is recommended that further studies about systematic and ecological investigations should be supported and performed on aquatic insects which exist in this important habitat. As well as, monitoring of aquatic biota by trained health personnel can be a critical step to describe water quality in this river.

In Iran, several studies are conducted regarding malaria in which different aspects are published. These include insecticide resistance monitoring (14,46-56) sibling species, molecular study, new record (57-63) novel methods for vector control (64-69), faunistic study (70,71), use of plants for larval control (53,72-85) using bed nets and long lasting impregnated nets (86-92), morphological studies (93-95), malaria epidemiology (96-101), ecology of malaria vectors (13,16,17,102-106), community participation (91,107), vector control (12), repellent evaluation (108), anthropophic index of malaria vectors (109-111), and training (112,113).

The presence of predators in the breeding places may reduce the abundance of *Anopheles* and result in a reduction of malaria cases.

Acknowledgement

We would like to thank Ms. Rafiei and Ms. Abolhasani for providing laboratory equipment. This research was financially supported by School of Public Health, Tehran University of Medical Sciences (TUMS). The authors declare that there is no conflict of interest.

References

- Merritt RW, Cummins KW, editors. An introduction to the aquatic insects of North America. 1st ed. Dubuque: Kendall Hunt; 1996: p.1158
- Subramanian KA, Sivaramkrishnan KG. Aquatic Insects of India-A Field Guide. India Biodiversity. (Accessed in May 2015, 4, at <http://indiabiodiversity.org/biodiv/content/documents/document-3345fd78-8c3b-4e1c-8978-8a3e29efa493/830.pdf>).
- Compin A, Cereghino R. Sensitivity of aquatic insect species richness to disturbance in the Adour-Garonne stream system (France). *Ecological Indicators* 2003;3(2):135-42.
- Aldhous P. The world's forgotten crisis. *Nature* 2003;422(6929):251.
- Langergraber G, Muellegger E. Ecological Sanitation—a way to solve global sanitation problems? *Environ Int* 2005;31(3):433-44.
- Malekei-Ravasan N, Bahrami A, Shayeghi M, et al. Notes on the Iran Caddisflies and Role of Annulipalpians Hydropsychid Caddisflies as a Bio-monitoring Agent. *J Arthrop Borne Dis* 2013;7(1):71-82.
- Dohet A. Are caddisflies an ideal group for the biological assessment of water quality in streams? *Nova Suppl Ento* 2002;15:507-20.
- Lenat DR. A biotic index for the southeastern United States: derivation and list of tolerance values, with criteria for assigning water-quality ratings. *J North Am Benthol Soc* 1993; 12(3):279-90.
- Balachandran C, Dinakaran S, Chandran MS, et al. Diversity and Distribution of Aquatic Insects in Aghanashini River of Central Western Ghats, India. (Accessed in May 2015, 2, at http://wgbis.ces.iisc.ernet.in/energy/water/paper/lake2012_aquatic_insects/results.htm).
- Ahmadi R, Mohebbi F, Hagigi P, et al. Macro-invertebrates in the Wetlands of the Zarrineh estuary at the south of Urmia Lake (Iran). *Int J Environ Res* 2011;5(4):1047-52.
- Shayeghi M, Vatandoost H, Gorouhi A, et al. Biodiversity of Aquatic Insects of Zayandeh Roud River and Its Branches, Isfahan Province, Iran. *J Arthrop Borne Dis* 2014;8(2):197-203.
- Vatandoost H, Abai M, Abbasi M, et al. Designing of a laboratory model for evaluation of the residual effects of deltamethrin (K-othrine WP 5%) on different surfaces against malaria vector, *Anopheles stephensi* (Diptera: Culicidae). *J Vector Borne Dis* 2009; 46(4):261-7.
- Nejati J, Vatandoost H, Oshghi MA, et al. Some ecological attributes of malarial vector *Anopheles superpictus* Grassi in endemic foci in southeastern Iran. *Asian Pac J Trop Biomed* 2013; 3(12):1003-8.
- Vatandoost H, Shahi H, Abai M, et al. Larval habitats of main malaria vectors in Hormozgan province and their susceptibility to different larvicides. *Southeast Asian J Trop Med Public Health* 2004; 35(Suppl 2):22-5.
- Vatandoost H, Oshaghi M, Abaie M, et al. Bionomics of *Anopheles stephensi* Liston in the malarious area of Hormozgan province, southern Iran, 2002. *Acta Trop* 2006;97(2):196-203.
- Vatandoost H, Shahi M, Hanafi-Bojd A, et al. Ecology of *Anopheles dthali* Patton in Bandar Abbas district, Hormozgan Province, southern Iran. *J Arthrop Borne Dis* 2007;1(1):21-7.
- Vatandoost H, Emami S, Oshaghi M, et al. Ecology of malaria vector *Anopheles culicifacies* in a malarious area of Sistan va Baluchestan province, south-east Islamic Republic of Iran. *East Mediterr Health J* 2011;17(5):439-45.

18. Darilmaz MC, Incekara Ü, Vafaei R. Contributions to the knowledge of Iranian Aquatic Adephaga. SPIXIANA 2013;36(1):149-52.
19. Hosseinie S. A report on the aquatic Coleoptera of Bushehr, Iran. Newsletter of the Balfour-Browne Club 1992;50(1):10.
20. Hosseinie S. Report on the aquatic Coleoptera of Hormozgan, Iran. Latissimus (Newsletter of the Balfour-Browne club) 1992;1(1):22.
21. Hosseinie S. A survey of the water Beetles of Khuzestan, Iran. Latissimus (Newsletter of the Balfour-Browne club) 1994;4(1):23.
22. Hosseinie S. Report on the aquatic Coleoptera of Guilan, a province in the North of Iran. Latissimus. Newsletter of the Balfour-Browne club 1995:1:11.
23. Hosseinie S. Water beetles in the Mazandaran Province, north of Iran. Latissimus (Newsletter of the Balfour-Browne club) 1995;5:20.
24. Khazeni A, Adler PH, Telmadareiiy Z, et al. The Black Flies (Diptera: Simuliidae) of Iran. Zootaxa 2013;3718(6):600.
25. Khatami S, Riyazi B, Modiri Asari S. The quality assessment of Karaj River based on diversity of benthic macroinvertebrates families. Environ Sci Technol 2006;1(9):71-8.
26. Khavari R, Ghorashi M, Arian M, et al. Geomorphic Signatures of Active Tectonics in the Karaj Drainage Basin in south central Alborz, North of Iran. Geoscience 2010; 19(1):67-74.
27. Gholikandi GB, Orumieh H, Haddadi S, et al. Application of multivariate statistical techniques for surface water quality assessment: case study of Karaj River, Iran. Water Resources Management VI 2011;145:361.
28. Elliott JM, Humpesch UH, Macan TT, editors. Larvae of the British Ephemeroptera: a key with ecological notes. Freshwater Biological Assn 1988; 49:1-145.
29. Bouchard R. Guide to Aquatic Invertebrate Families of Mongolia. Saint Paul, Minnesota, USA 2012. <http://midge.cfans.umn.edu/research/mongoliaguide/>
30. Fernald H, Sheparad Harold H. Applied Entomology, an introductory textbook of insects in their relations to Man. 4th ed. New York: Mcgraw-Hill Book Company, Inc; 1942: p. 400
31. Dehghani R, Almasi H, Asadi M. The Fauna of Kashan Aquatic Insects. Feyz J 2005;8(4):24-9.
32. Abbaspour R, Hedayatifard M, Alizadeh Sabet H, et al. Bioassessment of Macrobenthic Fauna of the Cheshmeh Kileh River, Northern Iran. Am Eur J Agric Environ Sci 2013; 13(6):747-53.
33. Gleick P, Schneider S, editors. Encyclopedia of climate and weather. 2nd ed. New York: Oxford University Press; 1996; p. 817-23.
34. Morse JC. Trichoptera world checklist. Trichoptera World Checklist. (Accessed in May 2015, 2, <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.352.5310&rep=rep1&type=pdf>).
35. Mey W. Beitrag zur Trichoptera-Fauna Armeniens und des Iran (Trichoptera). Entomol Nachr Ber 2004;48(1):81-7.
36. Mirmoayedi A, Malicky H. An updated check-list of caddisflies (Insecta, Trichoptera) from Iran, with new records. Zool Middle East 2002;26(1):163-8.
37. Chvojka P. Contribution to the knowledge of the caddisfly fauna (Trichoptera) of Iran: description of new species and new distributional data. Acta Entomol Mus Nat Pragae 2006;46:245-55.
38. Wiggins GB, editor. Caddisflies: the underwater architects. Toronto: University of Toronto Press; 2004.
39. Barbour MT, Gerritsen J, Snyder B, et al. Rapid bioassessment protocols for use in streams and wadeable rivers. 2nd ed. Washington: USEPA; 1999.
40. Thorp JH, Covich AP, editors. Ecology and classification of North American freshwater invertebrates. 2nd ed. Academic Press; 2009.
41. Burington ZL. Larval taxonomy, phylogeny and historical biogeography of the genus Cheumatopsyche (Trichoptera: Hydropsychidae) in North America. Clemson Univ., 2011.
42. Burks BD, editor. The mayflies or Ephemeroptera of Illinois. 1st ed. State of Illinois: Champaign: Illinois Natural History Survey; 1953.
43. Chinery M, editor. Guide to the Insects of Britain and Western Europe. London: Collins; 1986.
44. Wang TQ, McCafferty WP. Heptageniidae (Ephemeroptera) of the World. I. Phylogenetic higher classification. Trans Am Entomol Soc 2004;130(1):11-45
45. Ghane A, Ahmadi M, Esmaelie A, et al. The Biological assessment of Chafroud River (Guilan Province) using Macro benthos demographic structure. JWSS, Isfahan Univ Technol 2006;1:247-57.
46. Salari Lak S, Vatandoost H, Entezarmahdi M, et al. Monitoring of insecticide resistance in *Anopheles sacharovi* (Favre 1903) in borderline of Iran, Armenia, Naxcivan and Turkey, 2001. Iran J Public Health 2002;31(3-4):96-9.
47. Enayati A, Vatandoost H, Ladonni H, et al. Molecular evidence for a kdr like pyrethroid resistance mechanism in the malaria vector mosquito *Anopheles stephensi*. Med vet Entomol 2003;17(2):138-44.
48. Vatandoost H, Mashayekhi M, Abaie M, et al. Monitoring of insecticides resistance in main malaria vectors in a malarious area of Kahnooj district, Kerman province,

- southeastern Iran. J Vector Borne Dis 2005;42(3):100-8.
49. Davarai B, Vatandoost H, Ladonni H, et al. Comparative efficacy of different imagicides against different strains of *Anopheles stephensi* in the malarious areas of Iran, 2004-2005. Pak J Biol Sci 2006;9(5):885-92.
 50. Hanafi-Bojd A, Vatandoost H, Jafari R. Susceptibility status of *Anopheles dthali* and *An. fluviatilis* to commonly used larvicides in an endemic focus of malaria, southern Iran. J Vector Borne Dis 2006;43(1):34-8.
 51. Davari B, Vatandoost H, Oshaghi M, et al. Selection of *Anopheles stephensi* with DDT and dieldrin and cross-resistance spectrum to pyrethroids and fipronil. Pestic Biochem Physiol 2007;89(2):97-103.
 52. Abai M, Mehravaran A, Vatandoost H, et al. Comparative performance of imagicides on *Anopheles stephensi*, main malaria vector in a malarious area, southern Iran. J Vector Borne Dis 2008;45(4):307-12.
 53. Vatandoost H, Khazani A, Rafinejad J, et al. Comparative efficacy of neem and dimethyl phthalate (DMP) against malaria vector, *Anopheles stephensi* (Diptera: Culicidae). Asian Pac J Trop Med 2008;1(3):1-6.
 54. Vatandoost H, Zahirnia A. Responsiveness of *Anopheles maculipennis* to different imagicides during resurgent malaria. Asian Pac J Trop Med 2010;3(5):360-3.
 55. Vatandoost H, Hanafi-Bojd AA. Indication of pyrethroid resistance in the main malaria vector, *Anopheles stephensi* from Iran. Asian Pac J Trop Med 2012;5(9):722-6.
 56. Soltani A, Vatandoost H, Oshaghi MA, et al. Baseline Susceptibility of Different Geographical Strains of *Anopheles stephensi* (Diptera: Culicidae) to Temephos in Malarious Areas of Iran. J Arthropod-Borne Dis 2013;7(1):55-6.
 57. Dezfouli SN, Oshaghi M, Vatandoost H, et al. rDNA-ITS2 based species-diagnostic polymerase chain reaction assay for identification of sibling species of *Anopheles fluviatilis* in Iran. Southeast Asian J Trop Med Public Health 2003; 34(1):56-60.
 58. Naddaf SR, Oshaghi MA, Vatandoost H. Confirmation of two sibling species among *Anopheles fluviatilis* Mosquitoes in south and southeastern Iran by analysis of cytochrome oxidase I gene. J Arthropod-Borne Dis 2012;6(2):144-8.
 59. Naddaf S, Oshaghi M, Vatandoost H, et al. Molecular characterization of *Anopheles fluviatilis* species complex in the Islamic Republic of Iran. East Mediterr Health J 2003;9(3):257-65.
 60. Oshaghi M, Sedaghat M, Vatandoost H. Molecular characterization of the *Anopheles maculipennis* complex in the Islamic Republic of Iran. East Mediterr Health J 2003;9(4):659-66.
 61. Sedaghat M, Linton YM, Oshaghi M, et al. The *Anopheles maculipennis* complex (Diptera: Culicidae) in Iran: molecular characterization and recognition of a new species. Bull Entomol Res 2003;93(06):527-35.
 62. Azari-Hamidian S, Abai M, Ladonni H, et al. *Anopheles pedtaeniatus* (Leicester) new to the Iranian mosquito fauna with notes on *Anopheles hyrcanus* group in Iran. J Am Mosq Control Assoc 2006;22(1):144-6.
 63. Oshaghi M, Shemshad K, Yaghobi-Ershadi M, et al. Genetic structure of the malaria vector *Anopheles superpictus* in Iran using mitochondrial cytochrome oxidase (COI and COII) and morphologic markers: A new species complex? Acta Trop 2007;101(3):241-8.
 64. Soltani A, Vatandoost H, Jabbari H, et al. Use of expanded polystyrene (EPS) and shredded waste polystyrene (SWAP) beads for control of mosquitoes. J Arthropod-Borne Dis 2008;2(2):12-20.
 65. Soltani A, Vatandoost H, Jabbari H, et al. Field efficacy of expanded polystyrene and shredded waste polystyrene beads for mosquito control in artificial pools and field trials, Islamic Republic of Iran. East Mediterr Health J 2012;18(10):1042-8.
 66. Omrani SM, Vatandoost H, Oshaghi MA, et al. Fabrication of an olfactometer for mosquito behavioural studies. J Vector Borne Dis 2010;47(1):17-25.
 67. Omrani S, Vatandoost H, Oshaghi M, et al. Differential responses of *Anopheles stephensi* (Diptera: Culicidae) to skin emanations of a man, a cow, and a Guinea pig in the olfactometer. J Arthropod-Borne Dis 2010;4(1):1-16.
 68. Chavshin AR, Oshaghi MA, Vatandoost H, et al. Identification of bacterial microflora in the midgut of the larvae and adult of wild caught *Anopheles stephensi*: A step toward finding suitable paratransgenesis candidates. Acta Trop 2012;121(2):129-34.
 69. Omrani S, Vatandoost H, Oshaghi M, et al. Upwind responses of *Anopheles stephensi* to carbon dioxide and L-lactic acid: an olfactometer study. East Mediterr Health J 2012;18(11):11344-42.
 70. Oshaghi M, Vatandoost H, Gorouhi A, et al. Anopheline species composition in borderline of Iran-Azerbaijan. Acta Trop 2011; 119(1):44-9.
 71. Moosa-Kazemi S, Vatandoost H, Nikookar H. Culicinae (Diptera: culicidae) mosquitoes in chabahar County, sistian and baluchistan Province, southeastern Iran. J Arthropod-Borne Dis 2009;3(1):29-35.
 72. Hadjiakhoondi A, Aghel N, Zamanizadeh-Nadgar N, et al. Chemical and biological study of mentha spicata essential oil from Iran. DARU 2000;8(1-2):19-21.
 73. Hadjiakhoondi A, Vatandoost H, Jamshidi A, et al. Chemical Constituents of Efficacy of Cymbopogon Olivieri (Boiss) Bar Essential Oil Against Malaria Vector, *Anopheles Stepensi*. DARU 2003; 11(3):125-8.

74. Hadjiakhoondi A, Vatandoost H, Abousaber M, et al. Chemical composition of the essential oil of *Tagetes minuta* L. and its effects on *Anopheles stephensi* larvae in Iran. *J Med Plants* 2000;7(1):33-9.
75. Oshaghi M, Ghalandari R, Vatandoost H, et al. Repellent effect of extracts and essential oil of *Citrus limon* (Rutaceae) and *Melissa officinalis* (Labiatae) against main malaria vector, *Anopheles stephensi* (Diptera: Culicidae) in Iran. *Iran J Public Health* 2003;32(4):47-52.
76. Vatandoost H, Vaziri V. Larvicidal activity of a neem tree extract (Neemarin) against mosquito larvae in the Islamic Republic of Iran. *East Mediterr Health J* 2004;10(4-5):573-81.
77. Sadat Ebrahimi S, Hadji Akhoundi A, Fereiduonian N, et al. The components of *Tagetes minuta* L. and its biological activities against malaria vector, *Anopheles stephensi* in Iran. *J Med Plants* 2005;4(1):43-7.
78. Hadjiakhoondi A, Vatandoost H, Khanavi M, et al. Biochemical investigation of different extracts and larvicidal activity of *Tagetes minuta* L on *Anopheles stephensi* larvae. *Iran J Pharm Sci* 2005;1(2):81-4.
79. Hadjiakhoondi A, Sadeghipour-Roodsari H, Vatandoost H, et al. Fatty acid composition and toxicity of *Melia azedarach* L. fruits against malaria vector *Anopheles stephensi*. *Iran J Pharm Sci* 2006;2(2):97-102.
80. Shahi M, Hanafi-Bojd A, Iranshahi M, et al. Larvicidal efficacy of latex and extract of *Calotropis procera* (Gentianales: Asclepiadaceae) against *Culex quinquefasciatus* and *Anopheles stephensi* (Diptera: Culicidae). *J Vector Borne Dis* 2010;47(3):185-8.
81. Khanavi M, Toulabi PB, Abai MR, et al. Larvicidal activity of marine algae, *Sargassum swartzii* and *Chondria dasyphylla* against malaria vector *Anopheles stephensi*. *J Vector Borne Dis* 2011;48(4):241-4.
82. Sedaghat M, Dehkordi AS, Abai M, et al. Larvicidal Activity of Essential Oils of Apiaceae Plants against Malaria Vector, *Anopheles stephensi*. *J Arthropod-Borne Dis* 2011;5(2):51-9.
83. Sedaghat MM, Dehkordi AS, Khanavi M, et al. Chemical composition and larvicidal activity of essential oil of *Cupressus arizonica* EL Greene against malaria vector *Anopheles stephensi* Liston (Diptera: Culicidae). *Pharm Res* 2011;3(2):135-9.
84. Vatandoost H, Sanei Dehkordi A, Sadeghi S, et al. Identification of chemical constituents and larvicidal activity of *Kelussia odoratissima* Mozaffarian essential oil against two mosquito vectors *Anopheles stephensi* and *Culex pipiens* (Diptera: Culicidae). *Exp Parasitol* 2012;132(4):470-4.
85. Khanavi M, Vatandoost H, Dehaghi NK, et al. Larvicidal Activities of Some Iranian Native Plants against the Main Malaria Vector, *Anopheles stephensi*. *Acta Med Iran* 2013;51(3):141-7.
86. Vatandoost H, Dehakia M, Djavadia E, et al. Comparative study on the efficacy of lambda-cyhalothrin and bifenthrin on torn nets against the malaria vector, *Anopheles stephensi* as assessed by tunnel test method. *J Vector Borne Dis* 2006;43(3):133-5.
87. Moosa-Kazemi S, Vatandoost H, Raeisi A, et al. Deltamethrin impregnated bed nets in a malaria control program in Chabahar, Southeast Baluchistan, IR Iran. *J Arthropod-Borne Dis* 2007;1(1):43-51.
88. Rafinejad J, Vatandoost H, Nikpoor F, et al. Effect of washing on the bio-efficacy of insecticide-treated nets (ITNs) and long-lasting insecticidal nets (LLINs) against main malaria vector *Anopheles stephensi* by three bioassay methods. *J Vector Borne Dis* 2008;45(2):143-50.
89. Vatandoost H, Ramin E, Rassi Y, et al. Stability and wash resistance of local made mosquito bednets and detergents treated with pyrethroids against *Anopheles stephensi*. *J Arthropod-Borne Dis* 2009;3(1):18-29.
90. Soleimani-Ahmadi M, Vatandoost H, Shaeghi M, et al. Field evaluation of permethrin long-lasting insecticide treated nets (Olyset sup® sup) for malaria control in an endemic area, southeast of Iran. *Acta Trop* 2012; 123(3):146-53.
91. Soleimani-Ahmadi M, Eshraghian MR, Madani AH, et al. Effects of educational intervention on long-lasting insecticidal nets use in a malarious area, southeast Iran. *Acta Med Iran* 2012;50(4):279-87.
92. Vatandoost H, Mamivandpoor H, Abai MR, et al. Wash Resistance and Bioefficacy of Alpha-cypermethrin Long Lasting Impregnated Nets (LLIN-Interceptor®) against *Anopheles stephensi* using Tunnel Test. *J Arthropod-Borne Dis* 2013;7(1):31-45.
93. Doosti S, Azari-Hamidian S, Vatoost H, et al. Taxonomic differentiation of *Anopheles sacharovi* and *An. maculipennis* sl (Diptera: Culicidae) larvae by seta 2 (antepalmar hair). *Acta Med Iran* 2006;44(1):21-7.
94. Doosti S, Vatandoost H, Oshaghi M, et al. Applying morphometric variation of seta 2 (Antepalmar Hair) among the larvae of the members of the Maculipennis Subgroup (Diptera: Culicidae) in Iran. *J Arthropod-Borne Dis* 2007;1(1):28-37.
95. Emami S, Vatandoost H, Oshaghi M, et al. Morphological method for sexing anopheline larvae. *J Vector Borne Dis* 2007;44(4):245-9.
96. Vatandoost H, Ashraf H, Lak SS, et al. Factors involved in the re-emergence of malaria in borderline of Iran, Armenia, Azerbaijan and Turkey. *Southeast Asian J Trop Med Public Health* 2003;34(2):6-14.
97. Hanafi-Bojd A, Vatandoost H, Philip E, et al. Malaria

- situation analysis and stratification in BandarAbbas County, southern Iran, 2004-2008. Iran J Arthropod-Borne Dis 2010;4(1):31-41.
98. Hanafi-Bojd A, Vatandoost H, Oshaghi M, et al. Entomological and epidemiological attributes for malaria transmission and implementation of vector control in southern Iran. Acta Tropica 2012;121(2):85-92.
 99. Hemami MR, Sari AA, Raeisi A, et al. Malaria elimination in Iran, importance and challenges. Int J Prev Med 2013;4(1):88-94.
 100. Vatandoost H, Rashidian A, Jafari M, et al. Demonstration of malaria situation analysis, stratification and planning in Minab District, southern Iran. Asian Pac J Trop Med 2011;4(1):67-71.
 101. Hanafi-Bojd A, Vatandoost H, Oshaghi M, et al. Spatial analysis and mapping of malaria risk in an endemic area, south of Iran: A GIS based decision making for planning of control. Acta Trop 2012;122(1):132-7.
 102. Vatandoost H, Oshaghi M, Abaie M, et al. Bionomics of *Anopheles stephensi* Liston in the malarious area of Hormozgan province, southern Iran, 2002. Acta Trop 2006;97(2):196-203.
 103. Hanafi-Bojd AA, Azari-Hamidian S, Hassan V, et al. Spatio-temporal distribution of malaria vectors (Diptera: Culicidae) across different climatic zones of Iran. Asian Pac J Trop Med 2011;4(6):498-504.
 104. Soleimani-Ahmadi M, Vatandoost H, Shaeghi M, et al. Vector ecology and susceptibility in a malaria-endemic focus in southern Islamic Republic of Iran. East Med Health J 2012;18(10):1034-41.
 105. Soleimani-Ahmadi M, Vatandoost H, Hanafi-Bojd A-A, et al. Environmental characteristics of anopheline mosquito larval habitats in a malaria endemic area in Iran. Asian Pac J Trop Med 2013;6(7):510-5.
 106. Mehravaran A, Vatandoost H, Oshaghi MA, et al. Ecology of *Anopheles stephensi* in a malarious area, southeast of Iran. Acta Med Iran 2012;50(1):61-5.
 107. Hanafi-Bojd A, Vatandoost H, Oshaghi M, et al. Knowledge, attitudes and practices regarding malaria control in an endemic area of southern Iran. Southeast Asian J Trop Med Public Health 2011;42(3):491-501.
 108. Vatandoost H, Hanafi-Bojd A. Laboratory evaluation of 3 repellents against *Anopheles stephensi* in the Islamic Republic of Iran. East Mediterr Health J 2008;14(2):260-7.
 109. Oshaghi MA, Chavshin AR, Vatandoost H, et al. Effects of post-ingestion and physical conditions on PCR amplification of host blood meal DNA in mosquitoes. Exp Parasitol 2006;112(4):232-6.
 110. Oshaghi MA, Chavshin AR, Vatandoost H. Analysis of mosquito bloodmeals using RFLP markers. Exp Parasitol 2006;114(4):259-64.
 111. Oshaghi M, Yaghoobi F, Vatandoost H, et al. *Anopheles stephensi* biological forms, geographical distribution, and malaria transmission in malarious regions in Iran. Pak J Biol Sci 2006;9(2):294-8.
 112. Vatandoost H, Mesdaghinia A, Zamani G, et al. Development of the Regional Malaria Training Centre in Bandar-e Abbas, Islamic Republic of Iran. East Mediterr Health J 2004;10(1-2):215-24.
 113. Mesdaghinia AR, Vatandoost H, Hanafi-Bojd AA, et al. Conducting international diploma course on malaria program planning and management (1996-2012). J Arthropod-Borne Dis 2013;7(2):100-12.