

# Nasal Carriage of Uncommon Coagulase-Negative Staphylococci in Nurses and Physicians of Tehran University Hospitals

Elaheh Salimi<sup>1</sup>, Zahra Pakbaz<sup>2</sup>, Mohammad Reza Pourmand<sup>1,2</sup>,  
Payman Avakh Majelan<sup>2</sup>, and Sanaz Dehbashi<sup>2</sup>

<sup>1</sup>Urology Research Center, Tehran University of Medical Sciences, Tehran, Iran

<sup>2</sup>Department of Pathobiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Received: 17 Jan. 2015; Accepted: 04 Jul. 2015

**Abstract-** Coagulase-negative staphylococci (CoNS) have been identified as a major cause of nosocomial infections. Nasal carriage of CoNS in nurses and physicians is known to be an important risk factor for potential hospital infections. This study was carried out to investigate the prevalence of nasal carriage of uncommon coagulase-negative staphylococci among nurse and physician staffs of Tehran University Hospitals. A total of 116 CoNS were isolated from anterior nares of the study participants working in different wards of the hospitals. Thirteen uncommon CoNS were identified using phenotypic and biochemical methods, were subsequently confirmed by API kits. *Staphylococcus xylosum*, *Staphylococcus haemolyticus*, and *Staphylococcus capitis* species accounted for 53.85%, 30.77%, and 15.38% from the isolates, respectively. Six isolates (46.15%) were found to be resistant to methicillin. In conclusion, screening of healthcare workers for uncommon CoNS colonization along with identification and testing for susceptibility of cultured isolates is of paramount importance in strengthening effective nosocomial infection control and prevention measures.

© 2016 Tehran University of Medical Sciences. All rights reserved.

*Acta Med Iran*, 2016;54(5):330-333.

**Keywords:** Healthcare workers; Nasal carriage; Coagulase negative staphylococci

## Introduction

Healthcare workers may carry pathogenic and opportunist bacteria in their nasal cavity, and subsequently transmit these organisms to patients and other healthcare workers (1,2). Staphylococci are frequent opportunist pathogens and important commensal bacteria residing in human nasal mucosa (3). Nasal carriage of these organisms in physicians and nurses has been identified as a risk factor for nosocomial infections (4).

To date, coagulase-negative staphylococci (CoNS) are generally recognized as an important cause of severe infections among community and healthcare workers (5). Nonetheless, there are clinical microbiology laboratories still not practicing the detection of CoNS based on the species of bacteria. The asymptomatic carriage of CoNS species and the increasing incidence of complications from infections other than the *S. epidermidis* species, such as *S. haemolyticus*, *S. capitis*, *S. warneri* and *S. lugdunensis*, laid the ground for

learning more about the epidemiology and pathogenic potential of the different Staphylococcal species (6,7).

Furthermore, the ability of CoNS species to develop resistance against a wide spectrum of antibiotics needed consideration in the control and prevention of infections. These microorganisms may also have the potential to transfer their resistance to methicillin resistant *S. aureus* (MRSA). Thus, understanding antibiotic resistance of bacterial species is not only limited to isolates from patients but also is imperative to the carriers as well (8,9). Decolonization is considered to be one of the methods for controlling transmission of methicillin or vancomycin resistant Staphylococcus strains from healthcare workers to patients. However, complete eradication of nasal carriage is not possible but with their reduction, the risk of transfer within the healthcare system can be decreased (10,11).

This study was aimed at detecting the uncommon species of staphylococci residing in the anterior nares of physicians and nurses working in Tehran University hospitals, and determining the susceptibility of isolates

**Corresponding Author:** M.R. Pourmand

Department of Pathobiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran  
Tel: +98 21 8895491, Fax: +98 21 88954910, E-mail address: mpourmand@tums.ac.ir

against antibiotics.

## Materials and Methods

### Collection of samples and identification of bacteria

The samples were collected from nurses and physicians working in different wards of the hospitals. The swabs taken from the anterior nares were subsequently cultured on blood agar. Plates were incubated at 35°C for 24 hours. The elementary identification of staphylococcal isolates was performed using phenotypic and biochemical methods (catalase, oxidase, mannitol fermentation, DNase, coagulase and

different disk tests). Suspicious types of any CoNS isolates were further identified at species level using API STAPH kit test according to the manufacturer's instructions.

### Susceptibility to antimicrobial agents

According to the guidelines of the Clinical and Laboratory Standards Institute (CLSI), disc diffusion susceptibility testing was performed on uncommon CoNS isolates using Mueller Hinton agar. Depending on the zone size, each antimicrobial agent was recorded and interpreted as resistant, susceptible or intermediate (Table 1).

**Table 1. The antibacterial susceptibility in uncommon CoNS isolates among nurses and physicians of Tehran University Hospitals**

Antibiotics	<i>S. xylosus</i>		<i>S. haemolyticus</i>		<i>S. capitis</i>		Total	
	Susceptible No %	Resistant No %	Susceptible No %	Resistant No %	Susceptible No %	Resistant No %	Susceptible No %	Resistant No %
Penicillin	5(71.4)	2(28.6)	2(50.0)	2(50.0)	0(0.0)	2(100)	7(53.85)	6(46.16)
Oxacillin	5(71.4)	2(26.4)	2(50.0)	2(50.0)	0(0.0)	2(100)	7(53.85)	6(46.16)
Mupirocin	6(85.7)	1(14.3)	4(100.0)	0(0.0)	1(50.0)	1(50.0)	11(84.61)	2(15.39)
Tetracycline	4(57.1)	3(42.9)	2(50.0)	2(50.0)	0(0.0)	2(100)	6(46.15)	7(53.85)
Erythromycin	5(71.4)	2(26.4)	2(50.0)	2(50.0)	0(0.0)	2(100)	7(53.85)	6(46.16)
Clindamycin	4(57.1)	3(42.9)	3(75.0)	1(25.0)	1(50.0)	1(50.0)	8(61.53)	5(38.47)
Rifampin	4(57.1)	3(42.9)	4(100.0)	0(0.0)	1(50.0)	1(50.0)	9(69.23)	4(31.77)
Tobramycin	4(57.1)	3(42.9)	4(100.0)	0(0.0)	1(50.0)	1(50.0)	9(69.23)	4(31.77)
Ciprofloxacin	7(100)	3(42.9)	4(100.0)	0(0.0)	0(0.0)	2(100)	11(84.61)	2(15.39)
Co-trimoxazole	7(100)	0(0.0)	4(100.0)	0(0.0)	1(50.0)	1(50.0)	12(92.30)	1(7.70)

## Results

From total of 116 identified CoNS from the anterior nares of the nurses and physicians, 13 of them were identified to be uncommon CoNS. Further analysis revealed that *S. xylosus*, *S. haemolyticus* and *S. capitis* accounted for 53.85% (n=7), 30.77% (n=4), 15.38% (n=2) of 13 uncommon CoNS isolates respectively.

A high rate of resistance was observed against tetracycline (53.85%), penicillin (46.15%) and erythromycin (46.15%). Whereas the lowest resistance identified was for co-trimoxazole (7.7%) and ciprofloxacin (15.38%). The pattern of antimicrobial susceptibility of uncommon CoNS isolates to other antimicrobial agents is summarized in the Table 1.

The oxacillin resistance in uncommon CoNS accounted for 46.15%. Six isolates (46.15%) were found to be resistant to methicillin. That is, two isolates from each the three species were resistant to oxacillin.

## Discussion

This study explored the nasal carriage of uncommon CoNS in nurses and physicians of Tehran University hospitals. Thirteen isolates of uncommon CoNS were detected consisting of *S. xylosus* (53.85%), *S. haemolyticus* (30.77%) and *S. capitis* (15.38%).

In this study, *S. xylosus* was the most prevalent uncommon CoNS among the nurses and physicians. *S. xylosus* is a commensal bacterium on the skin and mucous membranes of mammals and birds. It is more common in animals than humans (12,13). *S. haemolyticus* has the highest level of antibiotic resistance among the CoNS. *S. haemolyticus* has a tendency to develop multiple antibiotic resistance. Moreover, the expression of glycopeptide resistance is common among the strains. Based on the results of our study, *S. haemolyticus* isolates were susceptible to certain antibiotics. These rates are different from those reported by other authors (14-17). These variations may

be explained by differences in the local infections control measures and the sample size. *S. capitis* is a human commensal bacterium with the capacity of causing endocarditis on native and prosthetic heart valves (18). In the current study, *S. capitis* strains showed a high rate of antibiotic resistance. Similarly, Cui *et al.*, reported a high level of resistance to oxacillin, erythromycin and penicillin in *S. capitis* (19) but Sandoe *et al.*, reported methicillin-resistance as uncommon among the *S. capitis* isolates (18).

Although many studies have been conducted on the prevalence of CoNS nasal carriage in healthcare workers (20-22), there is not enough data to show the frequency of uncommon CoNS, and nearly all the isolates of CoNS are reported without any further identification at species level. However, many details have been released about coagulase positive staphylococci such as *S. aureus* (23).

The presence of uncommon CoNS nasal carriers among nurses and physicians has raised many concerns. Therefore, screening healthcare workers for uncommon coagulase negative staphylococcal colonization along with the identification and testing for susceptibility of cultured isolates are of paramount importance in strengthening nosocomial infection control and prevention measures.

### Acknowledgment

This research was supported by Tehran University of Medical Sciences, Tehran, Iran (grant number: 20105).

### References

1. Boyce JM. Preventing Staphylococcal Infections by Eradicating Nasal Carriage of *Staphylococcus aureus*. *Infect Control Hosp Epidemiol* 1996;17(12):775-9.
2. Khalili MB, Sharifi-Yazdi MK, Dargahi H, et al. Nasal Colonization rate of *Staphylococcus aureus* strains among Health Care Service Employee's of Teaching University Hospitals in Yazd. *Acta Med Iran* 2009;47(4):315-7.
3. Von-Eiff C, Peters G, Heilmann C. Pathogenesis of infections due to coagulase-negative staphylococci. *Lancet Infect Dis* 2002;2(11):677-85.
4. Rastegar-Lari A, Pourmand MR, Ohadian Moghadam S, et al. Prevalence of PVL-containing MRSA isolates among hospital staff nasal carriers. *Lab Med* 2011;42(5):283-86.
5. Klingenberg CA, Rønnestad AS, Anderson TG, et al. Persistent strains of coagulase-negative staphylococci in a neonatal intensive care unit: Virulence factors and invasiveness. *Clin Microbiol Infect* 2007;13(11):1100-11.
6. Kleeman KT, Bannerman TL, Kloos WE. Species Distribution of Coagulase-Negative Staphylococcal Isolates at a Community Hospital and Implications for Selection of Staphylococcal Identification Procedures. *J Clin Microbiol* 1993;31(5):1318-21.
7. Koziol-Montewka M, Szczepanic A, Baranowicz L, et al. The investigation of *Staphylococcus aureus* and coagulase-negative staphylococci nasal carriage among patients undergoing haemodialysis. *Microbiol Res* 2006;161(4):281-7.
8. Pourmand MR, Abdossamadi Z, Salari MH, et al. Slime layer formation and the prevalence of *mecA* and *aap* genes in *Staphylococcus epidermidis* isolates. *J Infect Dev Ctries* 2011;5(1):34-40.
9. Havaei SA, Ohadian Moghadam S, Pourmand MR, et al. Prevalence of genes encoding bi-component leukocidins among clinical isolates of methicillin-resistant *Staphylococcus aureus*. *Iranian J Publ Health* 2010;39(1):8-14.
10. Sefani S, Varaldo PE. Epidemiology of methicillin resistant staphylococci in Europe. *Clin Microbiol Infect* 2003;9(12):1179-86.
11. Coia JE, Duckwort GJ, Edwards DI, et al. Guidelines for the control and prevention of methicillin-resistant *Staphylococcus aureus* (MRSA) in healthcare facilities. *J Hosp Infect* 2006;63(Suppl 1):S1-44.
12. Nagase N, Sasaki A, Yamashita K, et al. Isolation and species distribution of staphylococci from animal and human skin. *J Vet Med Sci* 2002;64(3):245-50.
13. Kloos WE, Musselwhite MS. Distribution and persistence of *Staphylococcus* and *Micrococcus* species and other aerobic bacteria on human skin. *Appl Microbiol* 1975;30(3):381-5.
14. Fredheim EG, Klingenberg C, Rohde H, et al. Biofilm Formation by *Staphylococcus haemolyticus*. *J Clin Microbiol* 2009;47(4):1172-80.
15. Froggatt JW, Johnston JL, Galetto DW, et al. Antimicrobial resistance in nosocomial isolates of *Staphylococcus haemolyticus*. *Antimicrob Agents Chemother* 1989;33(4):460-6.
16. Schwalbe RS, Stapleton JT, Gilligan PH. Emergence of vancomycin resistance in coagulase-negative staphylococci. *N Engl J Med* 1987;316(15):927-31.
17. Biavasco F, Vignaroli C, Varaldo PE. Glycopeptide resistance in coagulase-negative staphylococci. *Eur J Clin Microbiol Infect Dis* 2000;19(6):403-17.
18. Sandoe JA, Kerr KG, Reynolds GW, et al. *Staphylococcus capitis* endocarditis: two cases and review of the literature. *Heart* 1999;82(3):e1.
19. Cui B, Smooker PM, Rouch DA, et al. Differences between Two Clinical *Staphylococcus capitis* Subspecies

- as Revealed by Biofilm, Antibiotic Resistance, and Pulsed-Field Gel Electrophoresis Profiling. *J Clin Microbiol* 2013;51(1):9-14.
20. Abera B, Kibret M, Mulu W. Knowledge and beliefs on antimicrobial resistance among physicians and nurses in hospitals in Amhara Region, Ethiopia. *BMC Pharmacol Toxicol* 2014;15(1):26.
  21. Akhtar N. Staphylococcal Nasal Carriage of Health Care Workers. *J Coll Physicians Surg Pak* 2010;20(7):439-43.
  22. Al-Abdli NE, Baiu SH. Nasal Carriage of Staphylococcus in Health Care Workers in Benghazi Hospitals. *Am J Microbiol Res* 2014;2(4):110-12.
  23. Pourmand MR, Memarian M, Hoseini M, et al. High prevalence of sea gene among clinical isolates of Staphylococcus aureus in Tehran. *Acta Med Iran* 2009;47(5):357-61.

Archive of SID