

Pleomorphic Ductal Carcinoma of Breast: Two Cases and Review Literature

Mansour Moghimi¹, Amir Aryanfar², Saeed Kargar³, Fatemeh Joukar²

¹ Department of Pathology, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

² School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

³ Department of Surgery, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

Received: 28 Feb. 2018; Accepted: 05 Nov. 2018

Abstract- We report two cases of pleomorphic breast carcinoma, a rare variant of high-grade invasive ductal breast carcinoma of no special type, which is a combination of pleomorphic and bizarre giant cells in an adenocarcinoma background. For gaining better insights, available literature is also reviewed.

© 2019 Tehran University of Medical Sciences. All rights reserved.

Acta Med Iran 2019;57(3):200-204.

Keywords: Pleomorphic; Breast; Carcinoma

Introduction

The most common cancer accounts for 29% of all cancer diagnosis, and the second cause of cancer death among females is breast cancer which is classified into two categories: lobular and ductal (ductal and lobular dominancy). Pleomorphic carcinoma is a rare variant of high-grade invasive ductal carcinoma of the breast with unfavorable prognosis (1,2). It is characterized by more than 50% of pleomorphic and bizarre giant cells in a background of adenocarcinoma with spindle and squamous differentiation based on World Health Organization (WHO) classification (3). Pleomorphism and high mitotic activity with atypical mitosis are the two main features of these cells (4). Due to its unusual morphology, pleomorphic carcinoma may be misdiagnosed with sarcoma; therefore, specific immunostaining for epithelial and mesenchymal markers can be useful in differentiation (5). Based on our knowledge, 84 pleomorphic carcinomas of breast cases have been reported in English.

Herein, we present two cases of pleomorphic carcinoma, and pertinent literature is reviewed.

Case Report

Case 1

A 55-year-old woman had a palpable, itchy, and erythematous mass in her left breast, suspected breast cancer, 4 months earlier. In spite of explaining the process and plan, she did not acquiesce the treatment. During this period of time, the lump became larger, painful, infected,

and ulcerated. Besides losing appetite, she showed significant weight loss. Furthermore, at the time of admission, she complained of purulent, odoriferous discharges from several fistules. In the primary observations, the skin was abnormal, firm with cream-gray colored areas on its surface with the maximum diameter of 13 cm. Additionally, nodules with a 0.5-1 cm diameter were observed.

She underwent a left total mastectomy. The specimen measuring 19×16×8 cm had semifirm consistency and showed total infiltration by tumor with cream-pink colored cut sections. Breast skin was involved, and multiple cutaneous ulcerations were seen. Also, there was no nipple due to malignant destruction. Microscopic examination revealed breast tissue infiltrated diffusely by the malignant highly pleomorphic epithelial neoplasm. The tumor was composed of many mono- and multinucleated giant cells with bizarre vesicular nuclei containing prominent eosinophilic nucleoli. In some regions, streams of tumoral spindle-shaped cells and also brisk mitotic figures with atypical form were present. These pleomorphic tumoral cells were described by immunostaining as strongly positive cells for CKAE1/3, Vimentin, and CD10 and negative for ER, PR, HER2, p53, S-100, CK5/6, and CK7. Ki-67 proliferation index was 40%. At the beginning of the follow-up, the bone scan was normal, and there was not any sign of metastasis. Despite the post-surgical stability, she had severe dry coughs. Chest X-ray demonstrated blunt pleural angles, coarse reticular opacities, and right lower lobe consolidation (70×60 mm). Mycobacterium tuberculosis infection was ruled out. Spiral CT-scan of

Corresponding Author: F. Joukar

School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
Tel: +98 913 4541125, Fax: +98 35 36236439, E-mail address: joukar.f@ssu.ac.ir

lungs and mediastinum with IV contrast injection was done which revealed bilateral pleural effusion, multiple peripheral and subpleural pulmonary nodules, and partial abnormal alveolar opacities of right lung middle zone and the majority of lower lobes that confirmed diffuse pulmonary metastasis. Multiple metastases to left supraclavicular lymph nodes (largest size 29×23 mm) compressing left subclavian vein were seen in Doppler sonography which caused left-hand edema. Later, she developed wound infection presenting as prurency, scales and pus discharges. She was expired about two months after the surgery.




Figure 1. Mastectomy specimen which is totally infiltrated by tumor and shows multiple cutaneous tumoral nodules with ulceration and sinus tracts




Figure 2. A. H and E. bizarre pleomorphic and giant tumoral cells B. CD10 C. CKEA1/3 D. vimentin E. ki-67

Case 2

A 40-year-old woman presented with a firm, palpable, painless, and non-itching mass in the upper inner quadrant of her left breast from 2.5 years ago which had grown in size. There was not any scale, discharge or nipple retraction on examinations, while peau d'orange

appearance was present. Ultrasonography reported a large, ill-defined hypoechoic lesion at 10-11 O'clock measuring 85×66 mm with central necrotic changes and increased the thickness of the skin. Furthermore, 3 hypoechoic lesions measuring 42×28 mm, 26×19 mm, and 21×14 mm were recognized in the left axillary region. Mammographic observations proved the ultrasonographic findings and noted severe edema and increased the thickness of the subcutaneous tissue. There was not any localized lesion in the areola or subareolar region. These findings were compatible with a malignant lesion with BI-RADS category 5. Primary pathologic observation of core needle biopsy reported a tumoral tissue consisted of cells with severe nuclear pleomorphism, high mitotic activity, and low ductal formation surrounded by fibrotic stroma. Six sessions of chemotherapy were effective, and the tumor size decreased, 4 cm at maximum diameter. Then, she was referred to our hospital for the surgery. The patient underwent total mastectomy with axillary lymph node dissection. All surgical margins, axillary lymph nodes, and nipple were free of tumor. Hematoxylin and Eosin (H and E) staining showed cysts lining with multinucleated bizarre neoplastic cells and filled with a fibrinous material. The cysts were infiltrated by inflammatory cells mostly lymphoplasm cells and hemosiderin-laden macrophages. These pleomorphic cells had large hyperchromatic nuclei with atypical mitosis. Furthermore, Mild fibroadenomatous change was observed. All surgical margins were free of tumor. Immunohistochemistry (IHC) study was performed and was positive for S-100 (some of the cells), Vimentin, CKEA1/3 and, and ER, while negative for HER2 and PR. These findings demonstrated pleomorphic carcinoma. She was discharged with a good condition and referred for the following radiotherapy.

Ten months after the last, 25th, session of radiotherapy, she has not experienced any recur or metastasis.




Figure 2. A. H and E B. S100 C. vimentin D. CKEA1/3

Discussion

Breast cancer, the most common cancer with a high mortality rate among women, can be classified into ductal and lobular types. Invasive ductal breast cancer is more common and accounts for about 75-85% of invasive breast cancer and more frequent in lower age groups in comparison with invasive lobular carcinoma (6-8). Although there were reports of pleomorphic bizarre giant cell in breast cancer before 2000 (9-11), Silver and Tavassoli described pleomorphic breast carcinoma as a unique morphological variant of high-grade ductal carcinoma by studying 26 cases and identifying some of their similar morphological features (12) that later led to the new WHO classification of breast tumors (13). In the latest WHO edition, pleomorphic breast carcinoma is a rare variant of high-grade invasive breast carcinoma of no special type that more than half of the texture should be a combination of pleomorphic and bizarre giant cells in a background of adenocarcinoma with a spindle or squamous differentiation (3). Although the entity has a poor prognosis, not all of the reported tumors behave badly. Presence of a spindle cell metaplastic component and tumor size greater than 5 cm in stages I-III disease decreased overall survival.

After presenting with a palpable mass, which is the most common symptom, most of the patients refer within a few weeks. Some others come after a recent increase in the size of a previous stable tumor; however, there are reports of delayed refer to like the first case of our study. In addition, some of the cases are found through mammographic screening or physical examination for other diseases (4,14).

Imaging features of pleomorphic carcinoma can be similar to the benign breast lesions, fibroadenoma, malignant phyllodes tumor, and inflammation. So, pathologic findings play a critical role in the diagnosis (4).

Several cases of pleomorphic carcinoma have been reported to display cyst formation. Yamada *et al.*, suggested 3 mechanisms for this finding: 1) carcinoma cells initially develop in the ductal wall and locally grow intraductally; 2) the cells block the lumen via stromal

involvement; 3) increasing inner pressure and/or possible necrotic changes (15).

Other breast tumors, such as invasive pleomorphic lobular carcinoma, invasive carcinoma with osteoclast-like giant cells, invasive carcinoma with chorioepithelioma features, metaplastic carcinoma with giant cells, mammary sarcoma with giant cells, and metastatic tumors can present pleomorphic tumor cells (13). Pleomorphic lobular carcinoma cells nuclei are hyperchromatic, irregular, and markedly pleomorphic with a tendency to arrange in a linear pattern. Unlike pleomorphic ductal carcinoma, these cells do not express E-Cadherin in IHC study. Moreover, CD68 and CK markers are useful to differentiate pleomorphic ductal carcinoma from invasive ductal carcinoma with osteoclast-like giant cells, which is CD68 positive and CK negative. β -HCG positivity also makes the diagnosis of invasive carcinoma with chorioepithelioma features more likely. Finally, IHC study with CK and EMA, which are epithelial markers, can be applied to differentiate pleomorphic carcinoma from mammary sarcoma. The multicentric pattern of breast involvement in metastatic carcinomas and patient history of a primary tumor can be useful for diagnosis and differentiation from pleomorphic carcinoma (16). IHC study in some of the patients was positive for Vimentin, S100 protein, p53, C-erbB-2. However, there are other markers expressed in a few reported cases (14,17).

Late diagnosis and lack of proper treatment of pleomorphic carcinoma can lead to a catastrophe; the same as our first patient with total breast tissue involvement, multiple sinus tracts, and several metastases. However, timely diagnosis through pathologic examination and specific IHC studies to rule out other similar tumors and appropriate following are associated with a better prognosis and longer survival.

Acknowledgments

The authors would like to appreciate Dr. Shokouh Taghipour Zahir for valuable comments and helping in finding patients' documents and Mr. Mojtaba Shirpour for language editing.

Table 1. Previous reports of pleomorphic carcinoma of the breast

Author, year	number of cases	age	tumor size/ diameter (cm)	Right /Left	stage/grade	metastasis	treatment
Silver <i>et al.</i> , 2000(14)	26	28-96 (mean=51)	0.5-15 (average, 5.4)	65% right	all stages	lymph nodes	biopsy and/or mastectomy (24) or lumpectomy (2)
Lenicek <i>et al.</i> , 2007(18)	1	83	5.8	right	NR	-	Radical mastectomy
Kurokawa <i>et al.</i> , 2009(19)	1	51	1	left	I	-	mastectomy, adjuvant chemotherapy
Nguyen <i>et al.</i> , 2010(2)	37	23-78 (mean = 52)	1.2-11.6 (average, 3.5)	71% left	all stages	lymph nodes, lung, brain, liver, bone, skin, sort tissue	total mastectomy (22) BCS (9) No surgery (3) Unknown (3)
Yamaguchi <i>et al.</i> , 2010(17)	1	17	7.6	left	T3N2M0/IIIA	lymph nodes, lung	mastectomy, chemotherapy, radiotherapy
Zhao <i>et al.</i> , 2010(4)	10	33-76 (mean = 50)	1-15 (average, 4.3)	60% right	I, II, IIb	lymph nodes, lumbar vertebrae, liver, lung, thoracic bone	radical mastectomy (9), conservative surgery (1)
Caruso <i>et al.</i> , 2011(20)	1 male	65	2.5	left	pT2 N1aMx/III	lymph nodes	mastectomy
Tacchini <i>et al.</i> , 2011(21)	1	44	1.2 & 0.6			-	
Cordoba <i>et al.</i> , 2012(5)	2	76, 46	5, <1	right, Both	NR	NR	total mastectomy, Segmental resection of breast
Tang <i>et al.</i> , 2014(22)	1	45	3.5	left	T2N1M0 IIB	lymph node	radical mastectomy, chemotherapy
Yamada <i>et al.</i> , 2013(15)	1	55	3.5	right	pT2N0M0 IIA	-	adjuvant chemotherapy, radiotherapy
Shousha, 2014 (23)	1	NR	5	left	III	-	mastectomy
Arasi <i>et al.</i> , 2016(16)	1	57	5.6×2.5	left	-	lymph node	modified radical mastectomy
Moghimi <i>et al.</i> , 2017	2	55, 40	19×16×8,	Both left	IV, IIIB	Lung, -	total mastectomy, neoadjuvant chemotherapy

L= lobular D= ductal NR= not reported

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics. *CA Cancer J Clin* 2016;66:7-30.
2. Nguyen C V, Falco R, Hunt KK, Nayeemuddin KM, Lester TR, Harrell RK, et al. Pleomorphic Ductal Carcinoma of the Breast: Predictors of Decreased Overall Survival Approval from the Institutional Review Board of. *Am J Surg Pathol* 2010;34:486-93.
3. Lakhani SR, Ellis IO, Schnitt SJ, Tan PH, van de Vijver MJ, eds. WHO Classification of Tumours of the Breast. 4th ed. International Agency for Research on Cancer, 2012 .
4. Zhao J, Lang R, Guo X, Chen L, Gu F, Fan Y, et al. Clinicopathologic characteristics of pleomorphic carcinoma of the breast. *Virchows Arch* 2010;456:31-7 .
5. Córdoba A, Del Rosario Mercado M, Vicente F, Ederra M, Torrea N, Guerrero-Setas D. Pleomorphic carcinoma of the breast with expression of macrophage markers: Report of two cases. *Pathol Int* 2012;62:491-5.
6. Arpino G, Bardou VJ, Clark GM, Elledge RM. Infiltrating

- lobular carcinoma of the breast: tumor characteristics and clinical outcome. *Breast Cancer Res* 2018;6:R149-56.
7. Mersin H, Yildirim E, Gülben K, Berberoğlu U. Is invasive lobular carcinoma different from invasive ductal carcinoma? *Eur J Surg Oncol* 2018;29:390-5.
 8. Jo S, Hiraoka M, Akuta K, Nishimura Y, Takahashi M, Nishida H, et al. Histopathological changes of human tumors following thermoradiotherapy. *Int J Radiat Oncol Biol Phys* 1989;17:1265-71.
 9. Douglas-Jones A, Barr WT. Breast carcinoma with tumor giant cells. Report of a case with fine needle aspiration cytology. *Acta Cytol* 1989;33:109-14.
 10. Gupta RK, Wakefield SJ, Lallu S, Fauck R, Simpson J, Dowle CS. Aspiration cytodiagnosis of breast carcinoma in pregnancy and lactation with immunohistochemical and electron microscopic study of an unusual mammary malignancy with pleomorphic giant cells. *Diagn Cytopathol* 1992;8:352-6.
 11. Gupta RK. Aspiration cytodiagnosis of a rare carcinoma of breast with bizarre malignant giant cells. *Diagn Cytopathol* 1996;15:66-9.
 12. Rossi S, Buscarini E, Garbagnati F, Di Stasi M, Quaretti P, Rago M, et al. Percutaneous treatment of small hepatic tumors by an expandable RF needle electrode. *AJR Am J Roentgenol* 1998;170:1015-22.
 13. Tavassoli F, Devilee P, eds. Pathology and genetics of tumours of the breast and female genital organs. 3rd ed. World Health Organization of Tumors, 2003.
 14. Silver SA, Tavassoli FA. Pleomorphic carcinoma of the breast: clinicopathological analysis of 26 cases of an unusual high-grade phenotype of ductal carcinoma. *Histopathology* 2003;36:505-14.
 15. Yamada S, Nabeshima A, Nagata Y, Tasaki T, Noguchi H, Kitada S, et al. Pleomorphic carcinoma of the breast associated with cyst formation: a unique surgical case focusing on cytological and immunohistochemical findings. *Cystic breast PC. Diagn Pathol* 2013;8:75.
 16. Arasi NE, Neelima G, Anunayi J, Kumar LS, Swaroopa MJ. Pleomorphic ductal carcinoma of the breast: A Rare Variant 2016;4:26-9.
 17. Yamaguchi R, Tanaka M, Yamaguchi M, Fukushima T, Kaneko Y, Otsuka H, et al. Pleomorphic carcinoma of the breast in a 17-year-old woman. *Med Mol Morphol* 2010;43:43-7.
 18. Leniček T, Szerda F, Demirović A, Mijić A, Krušlin B, Tomas D. Pleomorphic ductal carcinoma of the breast with predominant micropapillary features. *Pathol Int* 2018;57:694-7.
 19. Kurokawa K, Mouri Y, Asano A, Kamei K, Iwata Y, Isogai M, et al. Pleomorphic carcinoma with osteoclastic giant cells of the breast: Immunohistochemical differentiation between coexisting neoplastic and reactive giant cells. *Pathol Int* 2018;59:91-7.
 20. Caruso R, Palmeri R, Racchiusa S, Bonanno A, Lucianò R, Scardigno M, et al. Pleomorphic ductal carcinoma of the male breast: report of a rare case and review of literature. *Anticancer Res* 2011;31:3069-71.
 21. Tacchini D, Mastrogiulio M, Vassallo L, Ginori A. Pleomorphic giant cell ductal carcinoma of the breast. *Pathologica* 2011;103:68-70.
 22. Tang H, Liu F, Li H, Huang X, Zhao T. Pleomorphic carcinoma of breast: a case report and review of literature. *Int J Clin Exp Pathol* 2014;7:5215-20.
 23. Shousha S. Pleomorphic invasive ductal carcinoma of the breast in a patient with Huntington's disease. *Case Rep Pathol* 2014;2014:979137.