

ROLE OF HRCT IN DIAGNOSING ACTIVE PULMONARY TUBERCULOSIS

M. Bakhshayesh Karam MD, M. R. Masjedi MD, L. Fadaizadeh MD, P. Dokouhaki MD, S. Alinejad Tahery, S. J. Tabatabaie MD, S. Sadeghi MD

National Research Institute of Tuberculosis and Lung Disease, Maseh Daneshvary Hospital, Tehran, Iran.

• Abstract

Background-The purpose of this study is to show the role of high-resolution CT scan (HRCT) in diagnosing active pulmonary tuberculosis. Previous studies show that the CT finding of centrilobular nodules and a tree-in bud appearance in HRCT is more sensitive than chest X-ray in detection of early endobronchial spread. As documented by Im and colleagues, 97% of patients with active pulmonary TB had evidence of bronchogenic spread of the disease on HRCT.

Materials & Methods-One hundred and two patients clinically suspected of having active pulmonary TB with chest-X-ray appearances suggestive of the disease, underwent HRCT examination.

Results-HRCT results showed that 76 patients (74.5%) had active pulmonary TB and further clinical work-up indicated that the final diagnosis of the disease was confirmed in 52 patients (51%). According to binomial test the combination of the two main HRCT appearances, the centrilobular nodule and "tree-in-bud" signs confirms TB diagnosis accurately. The sensitivity of HRCT in diagnosing active pulmonary TB in this study was 96% and the negative predictive value was calculated as 93%.

Conclusion-We conclude that HRCT is a powerful and reliable diagnostic tool in TB diagnosis, which means that it can be used even before mycobacteriologic results are available.

Keywords • HRCT • active pulmonary tuberculosis

Introduction

The frequency of tuberculosis has risen dramatically throughout the world since 1985. In developing countries, pulmonary tuberculosis is still a common disease particularly among the socioeconomically disadvantaged, elderly, chronically debilitated, and prisoners. Thus prompt diagnosis and treatment of this disease is crucial.

The chest radiograph has historically been a major tool in tuberculosis diagnosis, and it is used in concert with tuberculin skin test as a means of detecting the disease. Stability of the radiographic findings is a primary component in the assessment of disease activity as defined by the American Thoracic Society^{1,2}. Accordingly, the chest radiograph has become increasingly important in disease detection, particularly in cases which patient reliability is not assured^{3,4}. In selected cases, high-resolution CT scan (HRCT) may identify indicators of active disease not seen on chest radiograph. In addition, CT is more sensitive in detection of subtle or occult parenchymal disease^{4,5}. Recent studies have suggested a possible role of CT in assessing disease activity^{6,7}. HRCT findings in active tuberculosis have been correlated with pathologic findings^{6,8}. As documented by Im and colleagues, 97 percent of patients of active tuberculosis had evidence of bronchogenic spread of disease on HRCT⁸⁻¹⁰.

The characteristic HRCT appearance of active endobronchial disease includes poorly defined centrilobular nodules of 5-8 mm in diameter, centrilobular branching structures, and the "tree-in bud" appearance. These lesions represent caseous material filling the bronchioles, alveolar ducts, and peribronchial alveoli as well as peribronchial granuloma. In the setting of acute infectious disease,

Archive of SID

they are characteristic of active tuberculosis⁸⁻¹³.

The purpose of this study is to show the role of HRCT in diagnosing active pulmonary tuberculosis, when other means of diagnosing tuberculosis e.g. culture, BAL (Broncho Alveolar Lavage), or TBLB (Trans Bronchial Lung Biopsy) fail, are unavailable or are time consuming.

Materials and Methods

One hundred and two patients clinically suspected of having active pulmonary tuberculosis and a chest radiograph with suggestive patterns of disease entered the study. These patients were sequentially enrolled during a cross sectional study from February 1999 through November 1999 at the Maseeh Daneshvary Hospital chest clinic. After meeting the above-mentioned criteria, the patients were included in our survey.

Our method for gathering information was through a questionnaire, which was completed according to signs and symptoms, together with chest X-ray reports and laboratory results.

First a HRCT was performed, and the patients were then admitted into hospital for further work-up (sputum smear and culture, and if needed, BAL and TBLB). Our final diagnosis was based on either of the following criteria:

1. Positive smear and/or culture results for tuberculosis bacilli.
2. Positive smear and/or BAL culture results for tuberculosis bacilli.
3. Positive pathologic TBLB results.

If all above-mentioned results were negative but the patient showed dramatic response to anti tuberculosis drugs, he or she was categorized as "smear- negative" pulmonary tuberculosis cases.

Chest radiographs were viewed in order to find typical active lesions (e.g. apical opacities).

All patients underwent HRCT with Siemens Somatom Plus CT scanner. Studies were performed with scans obtained at end-inspiratory lung volume. Intravenous contrast medium was not routinely utilized; instead, contrast media were selectively administered primarily to assess the mediastinum in patients with equivocal mediastinal pathology.

From the lung apices to the hemidiaphragms, 1.5-mm thick sections were obtained at 10mm intervals. All images were prospectively reconstructed using a high-resolution (bone) algorithm to evaluate the nature of the lung lesions.

At the end, the HRCT diagnosis was compared to that obtained from the clinical and paraclinical work-up of patient in order to evaluate the diagnostic yield of HRCT.

The analysis and comparison of rank values were performed through chi-square. P-values less than 0.05 were considered statistically significant. Sensitivity, specificity, positive and negative predictive value were also calculated.

Results

One hundred and two patients (38 females and 64 males) aged 1 to 83 years, with an average age of 47.15 ± 22.6 entered the study.

All had the clinical signs and symptoms of active pulmonary tuberculosis; comprising, 84.3% cough, 51% fever, 38.2% hemoptysis, 62.7% sputum, 31.6% night sweats, and 53.9% weight loss.

Archive of SID

Sixty one patients were sputum smear and culture negative, 9 were positive for both sputum smear and culture, 5 were sputum smear negative and culture positive, and 27 were diagnosed according to BAL and TBLB results. ([Table 1](#))

All cases had chest radiographs with suggestive pattern of active pulmonary tuberculosis, namely infiltration or cavitation in the upper lobes.

According to HRCT results, 76 patients (74.5%) had active pulmonary tuberculosis, while further clinical work-up confirmed the final diagnosis of active pulmonary tuberculosis in only 52 patients (51%; $P < 0.0001$, $\alpha = 0.05$; [Table 2](#)). The final diagnosis and HRCT diagnosis showed a strong positive correlation ($P < 0.0001$, $\alpha = 0.05$). The sensitivity of HRCT in diagnosing active pulmonary tuberculosis was 96%, the specificity was 50%, the positive predictive value was 67%, and the negative predictive value, 93%.

Because of low number of TBLB and BAL performed, no correlation was found between these and HRCT results.

In order to find the diagnostic yield of each of the characteristic radiologic manifestations of tuberculosis found on CT scan ([Table 1](#)), we performed binomial tests between the final diagnosis and each of the characteristic "tree-in-bud" and "centrilobular nodule" appearances.

Analysis of data showed that although none of these two parameters can individually be diagnostic ($P = 0.80$), their combination, however, accurately confirms the diagnosis of tuberculosis ($P = 0.082$).

Discussion

Post-primary tuberculosis is usually a disease of adolescence and adulthood^{7,14}. The earliest finding of post primary tuberculosis is a heterogeneous, poorly marginated opacity (exudative lesion), usually in the apical or posterior segment of the upper lobes. Cavitation is seen by chest radiography in 40%^{15,16} to 87%¹⁷ (mean 50%) of patients sometimes during the course of the disease. Chest CT has proved to be more accurate in detection of cavitation, particularly in cases complicated by extensive fibrosis and architectural distortion^{7,18,19}.

The most common complication of tuberculous cavitation is endobronchial spread which is detected radiographically in 19%¹⁵ to 58%¹⁷ and by HRCT in up to 98%³ of cases.

Thus, finding centrilobular nodules and a tree-in-bud appearance in HRCT is more sensitive than the chest radiograph in detection of early endobronchial disease⁸.

In selected cases, HRCT may identify indicators of active disease not seen on chest radiographs.⁸

The increased sensitivity of CT scan may also prompt diagnosis, even at a time when microbiology results are pending.

What we found in our study was that in 76 cases (74.5%), HRCT suggested the diagnosis of active pulmonary tuberculosis, before bacteriologic results were available. Further studies showed that 52 of these cases (51%) were in fact infected with *Mycobacterium tuberculosis* and thus underwent anti-tuberculosis therapy with promising results.

Chi-square analysis showed a strong positive correlation between the final diagnosis and HRCT diagnosis with a sensitivity of 96% and a specificity of 50%, which strongly confirms our hypothesis on the role of HRCT in diagnosing active pulmonary tuberculosis.

Archive of SID

Also, our results showed that although infiltration was the dominant HRCT manifestation, "centrilobular nodule" and "tree-in-bud" appearances were the main findings in the majority of active pulmonary tuberculosis cases.

In this study, we reached the conclusion that HRCT is a powerful and reliable tool in tuberculosis diagnosis, when other means of diagnosing tuberculosis (e.g., culture, BAL or TBLB) fail to settle the matter, are not available or time consuming.

References

1. American Thoracic Society. Diagnosis standards and classification of tuberculosis. *Am Rev Respir Dis* 1990; **142** (3): 725-35.
2. Smuts NA, Beyers N, Gie RP, et al. Value of the lateral chest radiograph in tuberculosis in children; *Pediatr Radiol* 1994; **24**: 478-80.
3. Page H, Adams MC. Radiologic manifestations of pulmonary tuberculosis; *Radiologic Clinics of North America*, 1995 ; **33** (4): 655-77.
4. Rom WN, Gray S. Tuberculosis. Boston, Little Brown company **1996**.
5. Webb WR, Muller NL, Naidich DP. *High Resolution CT of the Lung*; New York: Raven Press, **1992**.
6. Im JG, Itoh H, Lee KS, Han Mc. CT-Pathology correlation of Pulmonary Tuberculosis. *Crit Rev Diagn Imaging* 1995; **36** (3): 227-85.
7. Agrons GA, Markowitz RI, Kramer SS. Pulmonary tuberculosis in children. *Semin Roentgenol* 1993; **28**: 158-72.
8. Im I H, Young-Aos A, Lee JH, et al. Pulmonary tuberculosis: CT finding-early active disease and sequential change with anti-tuberculous therapy. *Radiology* 1993; **186**: 653-60.
9. Greeden JF, Webb WR, Warnock N. Centrilobular opacities in the lung on high resolution CT: diagnostic considerations and pathologic. *Am J Roentgenol* relation, 1994; **162**: 569-74.
10. Leek S, Srug KS, Lin TH, et al. Adult-onset pulmonary tuberculosis: Findings in chest radiographs and CT scans; *Am J Roentgenol* 1993; **160**: 753-8.
11. Hong SH, Im JG, Lee Js, et al. High Resolution CT appearance of miliary tuberculosis. *J Comput Assist Tomogr* 1998; **22**: 220.
12. Leeks, Kim YH, Kim WS, et al. Endobronchial tuberculosis; CT features. *J Comput Assist Tomogr* 1991; **15** (3): 424-8.
13. Im JG, Webb WR, Han Mc, et al. Apical opacity associated with pulmonary tuberculosis: High resolution CT findings; *Radiology* 1991; **178**: 727-31.
14. Buckner CB, Walker CW. Radiologic manifestations of adult tuberculosis; *J Thoracic Imag* 1990; **5**: 28-37.
15. Hadlock FP, Park SK, Awc RI, et al. Unusual radiographic findings in adult pulmonary tuberculosis. *Am J Roentgenol* 1980; **134**: 1015-18.
16. Woodring JW, Vandiviere HM, Fried Am, et al. Update: The radiographic features of pulmonary tuberculosis. *Am J Roentgenol* 1986; **148**: 497-506.
17. Christensen EE, Dietz GW, Ahn CH, et al. Initial roentgenographic manifestations of pulmonary mycobacterium tuberculosis, *M. kansasii* and *M. intracellulare* infections. *Chest* 1981; **80**: 132-36.
18. Kuhlman JE, Deutsch JH, Fishman EK, et al. CT features of thoracic mycobacterial disease; *Radiographics* 1990; **10**:413-31.
19. Naidich DT, McCauley DI, Leitman BS, et al. CT of pulmonary tuberculosis. In: Siegelman SS (ed); *Computed Tomography of the Chest*; New York; Churchill-Livingston, **1984**.

[AIM Home](#) | [Table of Contents](#)