

## Original Article

# Reference Values for Serum Creatinine with Jaffe-compensated Assay in Adult Iranian Subjects: Tehran Lipid and Glucose Study

Asghar Ghasemi PhD<sup>1,2</sup>, Iraj Azimzadeh BSc<sup>3</sup>, Saleh Zahediasl PhD<sup>1,2</sup>, Fereidoun Azizi MD<sup>2</sup>

## Abstract

**Background:** Chronic kidney disease is a worldwide public health problem and glomerular filtration rate (GFR), the best overall index of renal function, is most commonly estimated from serum creatinine concentrations. The aim of this study was to determine reference values for serum creatinine concentrations using data from a population-based study in Iran.

**Methods:** Serum creatinine was measured using the Jaffe method in 5247 men and women, aged 20–88 years, participants of the Tehran Lipid and Glucose Study. For calculating Jaffe compensated creatinine values in 382 samples, serum creatinine was measured using both the Jaffe and the enzymatic p-aminophenazone (PAP) methods. Linear regression analysis yielded a regression line equation of Jaffe-creatinine =  $0.863 \times \text{PAP-creatinine} + 38.9 \mu\text{mol/L}$  ( $r = 0.973$ ,  $n = 382$ ,  $P < 0.001$ ). CLSI/IFCC guidelines (International Federation of Clinical Chemistry/ Clinical and Laboratory Standards Institute), non-parametric method was used for determining creatinine reference values.

**Results:** Reference values for serum creatinine ranged between 47–98  $\mu\text{mol/L}$  (0.53–1.11 mg/dL), 37–68  $\mu\text{mol/L}$  (0.42–0.77 mg/dL), and 37–78  $\mu\text{mol/L}$  (0.42–0.88 mg/dL) in men, non-menopausal women, and menopausal women, respectively. Mean serum creatinine concentration was significantly higher in men compared to women for both age  $\leq 50$  years [ $70 \pm 11$  vs.  $50 \pm 10 \mu\text{mol/L}$  ( $0.79 \pm 0.12$  vs.  $0.57 \pm 0.11$  mg/dL),  $P < 0.001$ ] and age  $> 50$  years [ $73 \pm 12$  vs.  $55 \pm 12 \mu\text{mol/L}$  ( $0.83 \pm 0.14$  vs.  $0.62 \pm 0.14$  mg/dL),  $P < 0.001$ ].

**Conclusion:** Reference values for serum creatinine using the compensated Jaffe method are presented in Iranian subjects, values that could help assessment of kidney function.

**Keywords:** Jaffe assay, reference values, serum creatinine

**Cite this article as:** Ghasemi A, Azimzadeh I, Zahediasl S, Azizi F. Reference Values for Serum Creatinine with Jaffe-compensated Assay in Adult Iranian Subjects: Tehran Lipid and Glucose Study. *Arch Iran Med*. 2014; **17**(6): 394 – 399.

## Introduction

Chronic kidney disease is a worldwide public health problem with increasing prevalence and incidence.<sup>1,2</sup> The incidence of chronic kidney disease has been reported to be above 2% each year in the Iranian population.<sup>3</sup> Glomerular filtration rate (GFR), the best overall index of renal function, is most commonly estimated from serum creatinine concentration,<sup>1,4,5</sup> and any equation using serum creatinine level for estimating GFR is dependent on the serum creatinine assay.<sup>6,7</sup> In addition, serum creatinine values are also used for assessing liver function.<sup>8</sup> Serum creatinine concentration is affected by factors other than creatinine filtration, including sex, age, race, diet, muscle mass, and the analytical method.<sup>4,9,10</sup> The Jaffe method, used in most routine laboratories, has low specificity and overestimates serum creatinine by approximately 20–30% in physiological values, due to non-creatinine chromogens, mainly proteins.<sup>6,9,11,12</sup> Enzymatic creatinine methods are more specific<sup>9</sup> and have been widely adopted for routine clinical laboratory as alternatives to the alkaline

picrate methods.<sup>10</sup>

For serum creatinine, like other naturally occurring biochemical compounds, a reference interval needs to be provided.<sup>9,13</sup> Reports of reference intervals for serum creatinine levels in Asian populations, considering IFCC (International Federation of Clinical Chemistry) criteria with accurate description of the measurement method, are scant<sup>9</sup> and to our knowledge, there is no documented report on reference values of serum creatinine in Iran. The aim of this study was therefore to determine age- and sex-specific reference intervals for serum creatinine concentrations using data from a population-based study from Iran.

## Subjects and Methods

### Subjects

The Tehran Lipid and Glucose Study (TLGS) was initiated in 1999, aiming to determine the prevalence of non-communicable disease risk factors.<sup>14</sup> A multistage stratified cluster random sampling technique was used to select 15,005 persons, aged over 3 years, from District 13 of Tehran, which is representative of Tehran's population.<sup>15</sup> In the current study, subjects ( $n = 10,795$ ) were participants, aged  $\geq 20$  years, of phase 4 TLGS (June 2008 to September 2011). Excluded were pregnant women, hypertensive subjects, those with diabetes, history of cardiovascular disease, cancer, and diarrhea or those using any medications including steroids, diuretics, beta-blockers, digitals, calcium channel blockers, angiotensin converting enzyme inhibitors, aspirin and other anticoagulants, lipid lowering drugs, male or female hormones,

**Authors' affiliation:** <sup>1</sup>Endocrine Physiology Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>2</sup>Endocrine Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran. <sup>3</sup>Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

**Corresponding author and reprints:** Asghar Ghasemi PhD, Endocrine Physiology Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, No. 24, Parvaneh St., Velenjak, Tehran, Iran. P.O. Box: 19395-4763, Tehran, Iran. Tel: +98 21 22409309; Fax: +98 21 22416264; E-mail: Ghasemi@endocrine.ac.ir




Figure 1. Study population and exclusions

contraceptives (oral or injection), or drugs for thyroid disorders. Subjects with a history of hospitalization during the past 3 months and those with a history of significant weight loss during the past 6 months were also excluded. After application of the exclusion criteria, 5247 apparently healthy participants (2792 men and 2455 women), aged 20 to 88 years, remained for analysis. A separate analysis was performed for healthy menopausal women ( $n = 452$ ; age range 51–84 years) (Figure 1). The study was approved by the ethics committee of the Research Institute for Endocrine Sciences and written informed consent was obtained from each participant.

#### Anthropometric and clinical assessments

Details of data collection in the TLGS have been published previously<sup>15</sup>; in brief, weight and height were measured according to standard protocols. Body mass index (BMI) was calculated as weight (Kg) divided by square of height ( $m^2$ ). Blood pressure was measured twice after 15 minutes of rest and the mean of two measurements was reported.

#### Creatinine measurement

Blood samples were obtained in a sitting position after 12–14 hours overnight fasting and centrifuged, within 30 to 45 minutes of collection; all blood analyses were done at the TLGS research laboratory on the day of sample collection. Serum creatinine was measured using the photometric Jaffe method (Pars Azmoon Kit, Tehran, Iran) in which creatinine reacts with picrate in an alkaline medium to yield an orange-red color, read at 505 nm. In 382 samples, creatinine measurement was done with both Jaffe and enzymatic p-aminophenazone (PAP) methods. Intra-


assay CVs were 2.2% and 3.1% for the Jaffe and PAP method, respectively ( $n = 72$ ). Inter-assay CVs for normal creatinine concentration were 4.1% and 6.1% for Jaffe and PAP method, respectively and for high creatinine concentration were 1.3% and 1.7% for the Jaffe and PAP method, respectively ( $n = 17$ ). Bland-Altman method comparison was used for comparing creatinine measurements by the Jaffe and PAP methods.

#### Determining outliers


The Dixon outlier range statistic was used for determining outliers as it has been recommended by Clinical and Laboratory Standards Institute (CLSI) for reference intervals determined by the nonparametric procedure.<sup>16</sup> In the Dixon test, if the D/R ratio exceeds 1/3, the extreme value is considered as outlier and should be deleted, where D is the absolute difference between the most extreme value and the next most extreme value and R is the range of the values.

#### Determining serum creatinine reference values

We used the CLSI/IFCC guidelines, non-parametric method, to determine reference values.<sup>17,18</sup> The retrospective (*a posteriori*) selection of individuals from a population-based study was used as it is considered ideal for the study of exclusion and partitioning criteria according to IFCC.<sup>19</sup> For the IFCC non-parametric method, which is recommended for determining reference values,<sup>20</sup> values were sorted in ascending order and rank numbers were assigned to values. Rank numbers of the 0.025 and 0.975 fractiles were computed as  $0.025 \times (N + 1)$  and  $0.975 \times (N + 1)$ , respectively and considered as reference intervals.



**Figure 2. (A)** Linear regression analysis of serum creatinine measured by conventional Jaffe method, compared to the PAP enzymatic method. The analysis yielded a regression line equation of:  $y = 0.863x + 38.9 \mu\text{mol/L}$  ( $r = 0.973$ ,  $n = 382$ ,  $p < 0.001$ ). **(B)** Bland and Altman analysis of serum creatinine comparison data: correlation  $r = -0.460$  ( $P < 0.001$ ), slope  $= -0.121$  ( $P < 0.001$ ), intercept  $= 39.4$ . Statistically significant bias was found with a mean difference of  $28.3 \pm 4.6 \mu\text{mol/L}$ . To convert creatinine values from micromole per liter to milligram per deciliter divide by 88.4.



**Figure 3.** Comparison of serum creatinine concentration by sex and age. Serum Creatinine concentrations were significantly lower in women compared with men in all age groups ( $P < 0.001$ ). \*: significant difference with other groups. †: significant difference compared to men. To convert creatinine values from micromole per liter to milligram per deciliter divide by 88.4.

#### Definitions of variables

Diabetes was defined according to the American Diabetes Association as fasting serum glucose  $\geq 7.0$  mmol/L or 2-hour serum glucose  $\geq 11.1$  mmol/L and/or medical treatment.<sup>21</sup> Hypertension was defined as blood pressure  $\geq 140/90$  or using antihypertensive medication. Smoking was defined as using  $\geq 1$  cigarettes per day or using the water pipe. Women were considered menopausal if they had had no menstrual bleeding in the previous 12 months. History of cardiovascular disease included coronary heart disease (myocardial infarction, history of heart surgery, angioplasty, and hos-

pitalization in the coronary care unit) and cerebrovascular attack.

#### Statistical analysis

For comparing baseline variables between men and women, the independent sample *t*-test was used. Pearson correlation coefficient was used for calculating correlation between age and serum creatinine concentrations. Differences between serum creatinine concentrations in different age groups were compared by one-way analysis of variance and Tukey post hoc test was used for multiple comparisons. Two-sided *p* values less than 0.05 were considered statistically significant. SPSS (SPSS Inc., Chicago, IL, USA; Version 15) software was used for all statistical analyses except for the Bland-Altman method comparison for which GraphPad Prism version 5.00 for Windows (GraphPad Software, San Diego California USA) was used.

#### Results

This study was conducted on 5247 healthy subjects (2792 men and 2455 women), aged 20 to 88 years. Men were older than women ( $39.9 \pm 14.3$  vs.  $33.3 \pm 8.5$  years). Comparison between Jaffe and PAP method for measuring serum creatinine concentration showed a good correlation between the two methods with a regression line of:  $y$  (Jaffe-creatinine)  $= 0.863 \times$  (PAP-creatinine)  $+ 38.9 \mu\text{mol/L}$  ( $r = 0.973$ ,  $n = 382$ ,  $P < 0.001$ ); this equation was then used for calculating compensated creatinine values for the entire data (Figure 2A); although the mean difference was  $28.3 \pm 4.6 \mu\text{mol/L}$  ( $0.32 \pm 0.05$  mg/dL), a statistically significant bias was found and the magnitude of difference was lower with higher concentrations of serum creatinine (Figure 2B).

Mean serum creatinine concentration was significantly higher in men compared to women for both age  $\leq 50$  years [ $70 \pm 11$  vs.  $50 \pm 10 \mu\text{mol/L}$  ( $0.79 \pm 0.12$  vs.  $0.57 \pm 0.11$  mg/dL),  $P < 0.001$ ] and age  $> 50$  years [ $73 \pm 12$  vs.  $55 \pm 12 \mu\text{mol/L}$  ( $0.83 \pm 0.14$  vs.

**Table 1.** Reference intervals for serum creatinine concentration ( $\mu\text{mol/L}$ ) in healthy adult subjects by age and sex using the compensated Jaffe method<sup>a</sup>.

Age (years)	n	95% Reference intervals	Mean $\pm$ SD	Median	IQR	Min	Max
Men							
29–20	856	88–47	70 $\pm$ 10	68	79–68	37	98
39–30	633	88–47	71 $\pm$ 11	68	79–68	37	98
50–40	701	98–47	71 $\pm$ 11	68	78–68	27	98
>50	602	98–47	73 $\pm$ 12	78	78–69	27	98
All	2792	98–47	72 $\pm$ 11	68	79–68	27	98
Women							
29–20	972	68–27	49 $\pm$ 9	47	57–47	27	88
39–30	818	68–37	50 $\pm$ 10	47	57–47	27	78
50–40	665	68–37	51 $\pm$ 10	47	57–47	27	88
>50	452 <sup>b</sup>	78–37	55 $\pm$ 12	57	57–47	27	98
All	2455 <sup>c</sup>	68–37	50 $\pm$ 10	47	57–47	27	88

a = According to Clinical and Laboratory Standards Institute (CLSI)/ International Federation of Clinical Chemistry (IFCC) criteria, non-parametric method;  
b = Healthy menopausal women; c = Menopausal women are not included; IQR = interquartile range. To convert creatinine values from micromole per liter to milligram per deciliter divide by 88.4.

0.62  $\pm$  0.14 mg/dL),  $P < 0.001$ ]. In addition, in both genders, serum creatinine concentrations were significantly higher in subjects aged above 50 years (Figure 3). A weak but significant correlation was found between serum creatinine levels and age in both men ( $r = 0.128$ ,  $n = 2792$ ,  $P < 0.001$ ) and women ( $r = 0.196$ ,  $n = 2907$ ,  $P < 0.001$ ).

Reference values for serum creatinine according to age and sex are presented in Table 1. Overall, 95% reference values for serum creatinine concentration ranged between 47–98  $\mu\text{mol/L}$  (0.53–1.11 mg/dL) and 37–68  $\mu\text{mol/L}$  (0.42–0.77 mg/dL) in men and women, respectively. Upper reference limits of serum creatinine levels were higher in men, aged over 40 years and in post-menopausal women. In addition, median serum creatinine level was higher in men older than 50 years and in post-menopausal women.

## Discussion

This study presents reference values for serum creatinine concentrations according to the Jaffe compensated method in apparently healthy Iranian subjects from a population-based study. These values could be used for diagnostic and therapeutic purposes.

In our study, the mean difference between creatinine measurement by the conventional Jaffe method and the enzymatic one, as the most specific routine method commercially available,<sup>22</sup> was 28.3  $\mu\text{mol/L}$  (0.32 mg/dL). By subtracting this value, results were considered Jaffe compensated. In line with this result, subtracting a constant of 26  $\mu\text{mol/L}$  (0.29 mg/dL) or 27  $\mu\text{mol/L}$  (0.31 mg/dL) of creatinine as an average value has been done by some kit producers to solve the interference problem in Jaffe method<sup>22,23</sup>. Subtractions of 15  $\mu\text{mol/L}$  (0.17 mg/dL),<sup>24</sup> 18  $\mu\text{mol/L}$  (0.20 mg/dL),<sup>8</sup> or 21  $\mu\text{mol/L}$  (0.24 mg/dL)<sup>12</sup> have also been reported. In addition, creatinine-free serum samples, when measured by the Jaffe kinetic method, have on the average 26.5  $\mu\text{mol/L}$  (0.30 mg/dL) creatinine<sup>12,25</sup> and a positive difference of about 27  $\mu\text{mol/L}$  (0.31 mg/dL) has been reported between the Jaffe and the HPLC method for creatinine measurement.<sup>26,27</sup>

In our study, serum creatinine levels were higher in men compared to women in all age groups; a finding similar to those of

other reports,<sup>24,28,29</sup> which may be due to muscle mass.<sup>28</sup> In addition, in our study, serum creatinine levels were higher in subjects aged >50 year, revealing positive correlation between age and serum creatinine levels. In line with this result, Wang et al., reported higher serum creatinine values in subjects > 60 years<sup>24</sup> and it has been reported that serum creatinine increases as a function of age with 0.22  $\mu\text{mol/L}$  (0.003 mg/dL) per year in men and 0.14  $\mu\text{mol/L}$  (0.002 mg/dL) per year in women.<sup>22</sup>

We found reference values for serum creatinine concentration to be 47–98  $\mu\text{mol/L}$  (0.53–1.11 mg/dL) and 37–68  $\mu\text{mol/L}$  (0.42–0.77 mg/dL) in men and women, respectively. Reference values for serum creatinine concentrations in some countries are summarized in Table 2.<sup>12,22–25,28–31</sup> Our values, especially the upper reference limit which is medically more important than the lower limit,<sup>23</sup> are very close to those of China<sup>24</sup> and Kenya<sup>28</sup> and lower than those of European countries. In line with these results, serum creatinine concentrations in Asians have been reported to be less than those of the Caucasians, which may be due to less muscle mass in the former.<sup>32</sup> This issue raises the need for inclusion of an Asian ethnic factor for calculation of estimated GFR according to the MDRD formula.<sup>32</sup> Although the Nordic group suggest that common reference intervals for serum creatinine could be used,<sup>33</sup> differences have been reported in Asians between cities.<sup>34</sup> Upper limits of our reference values were higher in men and in menopausal women, aged over 50 years. Partition age for reporting creatinine reference values has been reported to be 50, 60, or 70 years and in line with our results, higher values have been reported with increasing age.<sup>24</sup>

The strengths of this study include a relatively large sample size used for determining reference values. In addition, our samples were obtained from individuals of a population-based study, which could provide the best reference intervals for use in preventive medicine.<sup>35</sup> As a limitation, we used the compensated Jaffe creatinine assay while the reference method for serum creatinine measurement is isotope dilution mass spectrometry<sup>7</sup>; in the compensated Jaffe creatinine assay recalibration consists of subtracting a constant value from the results of conventional Jaffe creatinine assay for non-specific chromogens; however, the levels of chro-



**Table 2.** Serum creatinine reference values ( $\mu\text{mol/L}$ ) in selected countries.

Country	Year	n	Sex (M/F)	Age (years)	Creatinine reference values ( $\mu\text{mol/L}$ )		Creatinine measurement method	Reference number
					Males	Females		
South Australia (Adelaide)	2000	562	293/269	18–70	62–106	44–80	Compensated Jaffe	25
Germany	2004	240	120/120	18–74	63–103	48–85	Compensated Jaffe	12
Nordic countries (Denmark, Finland, Iceland, Norway, and Sweden)	2004	1802	858/944	$\geq 18$	62–106	47–88	Compensated Jaffe	30
India	2006	1121	848/273	20–60	71–115	53–88	Enzymatic	31
Spain	2007	468	248/220	19–65	64–106	52–85	Compensated Jaffe	23
Belgium	2008	32658	14312/18346	20–70	56–103	42–82	Enzymatic	29
Italy	2010	16939	ND	18–79	57–103	40–88	Compensated Jaffe	22
China	2011	778	433/345	20–91	58–98	43–70	Enzymatic	24
Kenya	2012	360	206/154	18–50	56–99	48–85	ND	28
Iran	Present study	5247	2792/2455	20–88	47–98	37–68	Compensated Jaffe	Present study

ND = not determined. To convert creatinine values from micromole per liter to milligram per deciliter divide by 88.4.

mogens vary from one subject to another and low protein levels in samples lead to underestimation of creatinine values; these factors impact the measured levels of creatinine.<sup>8</sup>

In conclusion, the results of this study present reference intervals for serum creatinine concentration, according to the compensated Jaffe method, derived from a population-based study in Iran to be 47–98  $\mu\text{mol/L}$  (0.53–1.11 mg/dL) and 37–68  $\mu\text{mol/L}$  (0.42–0.77 mg/dL) in men and women, respectively. These values could be used for diagnostic and therapeutic purposes.

## Acknowledgments

This study was supported by Grant No.121 from the National Research Council of the Islamic Republic of Iran. The authors wish to thank Ms. N. Shiva for critical editing for English grammar and syntax of the manuscript.

## References

- Levey AS, Coresh J, Balk E, Kausz AT, Levin A, Steffes MW, et al. National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Ann Intern Med.* 2003; **139**: 137–147.
- Hosseinpanah F, Kasraei F, Nassiri AA, Azizi F. High prevalence of chronic kidney disease in Iran: a large population-based study. *BMC Public Health.* 2009; **9**: 44.
- Tohidi M, Hasheminiya M, Mohebi R, Khalili D, Hosseinpanah F, Yazdani B, et al. Incidence of chronic kidney disease and its risk factors, results of over 10 year follow up in an Iranian cohort. *PLoS One.* 2012; **7**: e45304.
- Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med.* 1999; **130**: 461–470.
- Selvin E, Manzi J, Stevens LA, Van Lente F, Lacher DA, Levey AS, et al. Calibration of serum creatinine in the National Health and Nutrition Examination Surveys (NHANES) 1988–1994, 1999–2004. *Am J Kidney Dis.* 2007; **50**: 918–926.
- Coresh J, Astor BC, McQuillan G, Kusek J, Greene T, Van Lente F, et al. Calibration and random variation of the serum creatinine assay as critical elements of using equations to estimate glomerular filtration rate. *Am J Kidney Dis.* 2002; **39**: 920–929.
- Hetu PO, Gingras ME, Vinet B. Development and validation of a rapid liquid chromatography isotope dilution tandem mass spectrometry (LC-IDMS/MS) method for serum creatinine. *Clin Biochem.* 2010; **43**: 1158–1162.
- Kuster N, Bargnoux AS, Pageaux GP, Cristol JP. Limitations of compensated Jaffe creatinine assays in cirrhotic patients. *Clin Biochem.* 2012; **45**: 320–325.
- Cerriotti F, Boyd JC, Klein G, Henny J, Queralto J, Kairisto V, et al. Reference intervals for serum creatinine concentrations: assessment of available data for global application. *Clin Chem.* 2008; **54**: 559–566.
- Myers GL, Miller WG, Coresh J, Fleming J, Greenberg N, Greene T, et al. Recommendations for improving serum creatinine measurement: a report from the Laboratory Working Group of the National Kidney Disease Education Program. *Clin Chem.* 2006; **52**: 5–18.
- Chromy V, Rozkosna K, Sedlak P. Determination of serum creatinine by Jaffe method and how to calibrate to eliminate matrix interference problems. *Clin Chem Lab Med.* 2008; **46**: 1127–1133.
- Junge W, Wilke B, Halabi A, Klein G. Determination of reference intervals for serum creatinine, creatinine excretion and creatinine clearance with an enzymatic and a modified Jaffe method. *Clin Chim Acta.* 2004; **344**: 137–148.
- Jorgensen LG, Brandslund I, Hyltoft Petersen P, Stahl M, de Fine Olivarius N. Creation of a low-risk reference group and reference interval of fasting venous plasma glucose. *Clin Chem Lab Med.* 2004; **42**: 817–823.
- Azizi F, Ghanbarian A, Momenan AA, Hadaegh F, Mirmiran P, Hedayati M, et al. Prevention of non-communicable disease in a population in nutrition transition: Tehran Lipid and Glucose Study phase II. *Trials.* 2009; **10**: 5.
- Azizi F, Rahmani M, Emami H, Mirmiran P, Hajipour R, Madjid M, et al. Cardiovascular risk factors in an Iranian urban population: Tehran lipid and glucose study (phase 1). *Soz Präventivmed.* 2002; **47**: 408–426.
- Horn PS, Feng L, Li Y, Pesce AJ. Effect of outliers and nonhealthy individuals on reference interval estimation. *Clin Chem.* 2001; **47**: 2137–2145.
- Horn PS, Pesce AJ, Copeland BE. A robust approach to reference interval estimation and evaluation. *Clin Chem.* 1998; **44**: 622–631.
- Solberg HE. Approved recommendation (1987) on the theory of reference values Part 5. Statistical treatment of collected reference values. Determination of reference limits. *J Clin Chem Clin Biochem.* 1987; **25**: 645–656.
- Petitclerc C, Solberg HE. International Federation of Clinical Chemistry (IFCC), Scientific Committee, Clinical Section, Expert Panel on Theory of Reference Values, and International Committee on Standardization in Haematology (ICSH), Standing Committee on Reference Values. Approved Recommendation (1987) on the theory of reference values. Part 2. Selection of individuals for the production of reference values. *J Clin Chem Clin Biochem.* 1987; **25**: 639–644.
- Henny J, Petitclerc C, Fuentes-Arderiu X, Petersen PH, Queralto JM, Schiele F, et al. Need for revisiting the concept of reference values. *Clin Chem Lab Med.* 2000; **38**: 589–595.
- American Diabetes Association. Standards of medical care in diabetes–2011. *Diabetes Care.* 2011. **34** (suppl 1): S11–S61.
- Arzideh F, Wosniok W, Haeckel R. Reference limits of plasma and serum creatinine concentrations from intra-laboratory data bases of several German and Italian medical centres: Comparison between direct and indirect procedures. *Clin Chim Acta.* 2010; **411**: 215–221.
- Fuentes-Arderiu X, Alvarez-Funes V, Coca-Fabregas L, Cruz-Placer M, Diaz-Fernandez J, Herrero-Bernal P, et al. Multicentre physiological reference values for the concentration of creatininium in plasma and diagnostic specificity of glomerular filtration rate estimated with the MDRD equation. *Clin Chem Lab Med.* 2007; **45**: 531–534.

24. Wang X, Xu G, Li H, Liu Y, Wang F. Reference intervals for serum creatinine with enzymatic assay and evaluation of four equations to estimate glomerular filtration rate in a healthy Chinese adult population. *Clin Chim Acta*. 2011; **412**: 1793–1797.
25. Mazzachi BC, Peake MJ, Ehrhardt V. Reference range and method comparison studies for enzymatic and Jaffe creatinine assays in plasma and serum and early morning urine. *Clin Lab*. 2000; **46**: 53 – 55.
26. Wuyts B, Bernard D, Van den Noortgate N, Van de Walle J, Van Vlem B, De Smet R, et al. Reevaluation of formulas for predicting creatinine clearance in adults and children, using compensated creatinine methods. *Clin Chem*. 2003; **49**: 1011–1014.
27. Chan MH, Ng KF, Szeto CC, Lit LC, Chow KM, Leung CB, et al. Effect of a compensated Jaffe creatinine method on the estimation of glomerular filtration rate. *Ann Clin Biochem*. 2004; **41**: 482 – 484.
28. Juma AA, Ngeranwa JJ, Njagi EN. Reference values for some renal function parameters for adult population in north-rift valley, kenya. *Indian J Clin Biochem*. 2012; **27**: 40 – 45.
29. Pottel H, Vrydags N, Mahieu B, Vandewynckele E, Croes K, Martens F. Establishing age/sex related serum creatinine reference intervals from hospital laboratory data based on different statistical methods. *Clin Chim Acta*. 2008; **396**: 49 – 55.
30. Martensson A, Rustad P, Lund H, Ossowicki H. Creatininium reference intervals for corrected methods. *Scand J Clin Lab Invest*. 2004; **64**: 439 – 441.
31. Verma M, Khadapkar R, Sahu PS, Das BR. Comparing age-wise reference intervals for serum creatinine concentration in a “Reality check” of the recommended cut-off. *Indian J Clin Biochem*. 2006; **21**: 90 – 94.
32. Hawkins RC. Differences in serum creatinine concentration between Caucasians, Chinese, Indians and Malays. *Clin Chim Acta*. 2010; **411**: 1393.
33. Rustad P, Felding P, Franzson L, Kairisto V, Lahti A, Martensson A, et al. The Nordic Reference Interval Project 2000: recommended reference intervals for 25 common biochemical properties. *Scand J Clin Lab Invest*. 2004; **64**: 271 – 284.
34. Ichihara K, Itoh Y, Lam CW, Poon PM, Kim JH, Kyono H, et al. Sources of variation of commonly measured serum analytes in 6 Asian cities and consideration of common reference intervals. *Clin Chem*. 2008; **54**: 356 – 365.
35. Hansen AM, Garde AH, Eller NH. Estimation of individual reference intervals in small sample sizes. *Int J Hyg Environ Health*. 2007; **210**: 471 – 478.

Archive of SID