

OCCURRENCE OF RICE INFECTION BY *Bipolaris sorokiniana* IN IRAN

S. NAEIMI^{1*}, V. KHOSRAVI² and T. TSUKIBOSHI³

(Received: 9. 3. 2011; Accepted: 7. 9. 2011)

Abstract

Diseased rice plants with foot rot as well as dark brown lesions on sheaths and stems were observed in a paddy field in Mazandaran province, Iran. Single lesions were excised and surface sterilized and then plated on water agar. The arising fungi were purified and refrigerated in agar slants. On the basis of morphological characteristics the fungus was identified as *Bipolaris sorokiniana*. The ITS sequence of the fungus was submitted to a BLAST search to find most similar sequences in GenBank. The search results showed highest similarity to eight strains of *B. sorokiniana* and *Cochliobolus sativus* (teleomorph). For phylogenetic comparison, the sequences of *Bipolaris* spp., *Cochliobolus* spp. together with other species belonging to closely related fungal genera were included. The resulted phylogram made with the neighbor-joining method using the program PAUP, showed that the Iranian strain (B54) formed a monophyletic group with seven *B. sorokiniana* and one *C. sativus* strain. To prove pathogenicity, several methods were done on two popular rice cultivars. This is the first record of *B. sorokiniana* on *Oryza sativa* in Iran.

Keywords: Pathogenicity, *Bipolaris sorokiniana*, *Oryza sativa*, spot blotch, Mazandaran province.

See Persian text for figures and tables (Pages ۳۵۳-۳۶۰).

*: Corresponding Author, Email: shnaeimi@yahoo.com

1. Res. Assist. Prof. of Plant Pathology, Amol Biological Control Research Laboratory, Iranian Res. Instit. of Plant Protec., Iran.
2. Scientific Member, Deputy of Iranian Rice Res. Institute, Amol, Mazandaran, Iran.
3. Prof. of Plant Pathology, Plant Pathology Laboratory, National Institute of Livestock and Grassland Science, National Agriculture and Food Research Organization, Japan.

References

- ASAD, S., IFTIKHAR, S., MUNIR, A. and AHMAD, I. 2009. Characterization of *Bipolaris sorokiniana* isolated from different agro-ecological zones of wheat production in Pakistan. **Pak. J. Bot.** 41(1): 301-308.
- CHAND, R., PANDEY, S.P., SINGH H.V., KUMAR, S. and JOSHI, A.K. 2003. Variability and its probable cause in the natural populations of spot blotch pathogen *Bipolaris sorokiniana* of wheat (*T. aestivum* L.) in India. **J. Plant Dis. Protect.** 110: 27-35.
- HERMOSA, M. R., GRONDONA, I., ITURRIAGA, E. A., DIAZ-MINGUEZ, J. M., CASTRO, C., MONTE, E. and GARCIA-ACHA, I. 2000. Molecular characterization of biocontrol isolates of *Trichoderma*. **Appl. Environ. Microbiol.** 66: 1890-1898.
- HETZLER, J., EYAL, J., FEHRMANN, H., MEHTA, Y.R., KUSHNIR, U., ZEKARIA-OREN, J. and COHEN, L. 1991. Interaction between *Cochliobolus sativus* and wheat cultivars. Pp. 266-283, *In*: Saunders, D.A. (Ed.), **Wheat for the Non-Traditional Warmer Areas**. Mexico, D.F., Mexico: CIMMYT.
- IFTIKHAR, S., ASAD, S., MUNIR, A., SULTAN, A. and AHMAD, I. 2009. Hosts of *Bipolaris sorokiniana*, the major pathogen of spot blotch of wheat in Pakistan. **Pak. J. Bot.** 41(3): 1433-1436.
- IRAM, S. and AHMAD, I. 2004. Diversity of *Bipolaris sorokiniana* isolates from wheat growing areas under rice-wheat cropping system in Punjab, Pakistan. **Pak. J. Bot.** 36(2): 439-444.
- JAISWAL, S. K., SWETA, PRASAD, L.C., SHARMA, S., KUMAR, S., PRASAD, R., PANDEY, S. P., CHAND, R. and JOSHI, A. K. 2007. Identification of molecular marker and aggressiveness for different groups of *Bipolaris sorokiniana* isolates causing spot blotch disease in wheat (*Triticum aestivum* L.). **Curr. Microbiol.** 55:135-141.
- KODAMA, F., TSUCHIYA, S., IWATA, T., IGARASHI, F. and SAWASAKI, A. 1979. Helminthosporium spot blotch of rice caused by *Helminthosporium sativum* P.K. & B. **Ann. Phytopath. Soc. Japan.** 45: 503-506.
- KUMAR, J., SCHAFER, P., HUCKELHOVEN, R., LANGEN, G., BALTRUSCHAT, H., STEIN, E., SUBRAMANIAN, N., KOGEL, K.H., KUMAR, J. and NAGARAJAN, S. 2002. *Bipolaris sorokiniana*, a cereal pathogen of global concern: cytological and molecular approaches towards better control. **Mol. Plant Pathol.** 3(4): 185-195.
- MANANDHAR, H.K., JØRGENSEN, H.J.L., MATHUR, S.B. and SMEDEGAARD-PETERSEN, V. 1998. Suppression of rice blast by preinoculation with avirulent *Pyricularia oryzae* and the nonrice pathogen *Bipolaris sorokiniana*. **Phytopathology** 88:735-739.
- MEHTA, Y.R. 1993. Spot blotch (*Bipolaris sorokiniana*). Pp 105-112, *In*: S.B., Mathur and B.M., Cunfer (Eds). **Seed Borne Disease and Seed Health Testing of Wheat**. Copenhagen: Institute of Seed Pathology for Developing Countries.
- NYVALL, R.F., and PERCICH, J.A. 1999. Development of fungal brown spot and spot blotch on cultivated wild rice in Minnesota. **Plant Dis.** 83:936-938.
- POLONI, A., PESSI, I.S., FRAZZON, A.P.G. and VAN DER SAND, S.T. 2009. Morphology, physiology, and virulence of *Bipolaris sorokiniana* isolates. **Curr. Microbiol.** 59:267-273.
- PREM, N., BISWAS, S.K., UPESH, K. and MOHD, R. 2009. Effect of Media, pH, temperature, host range and fungicides on *Bipolaris sorokiniana*. **Ann. Plant Protect. Sci.** 17 (2): 394-397.
- SAFAEE, D., OKHOVAT, S.M., HEDJAROUDE, GH. and YOUNESI, H. 2008. Diagnosis of *Bipolaris* form species involving root and crown rot of wheat, their distribution and disease severity in Kermanshah province. **J. Sci. Tech. Agr. Nat. Resour.** 43: 207-214.
- SIVANESAN, A. 1987. Graminicolous species of *Bipolaris*, *Curvularia*, *Drechslera*, *Exserohilum* and their teleomorphs. **Mycol. Pap.** 158: 1-261.
- TINLINE, R.D. 1988. *Cochliobolus sativus*, a pathogen of wide host range. **Adv. Plant Pathol.** 6:113-122.
- WALKER, J., KABLE, P.F., SMITH, A.M., DONALD, D.J., and BOEREMA, E.B. 1968. *Cochliobolus sativus* causing a leaf spot on rice in New South Wales. **Aust. J. Crop Sci.** 31:2, 82.

- WHITE, T.J., BRUNS, T., LEE, S. and Taylor, J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. Pp. 315–322, *In*: M. A. Innis, D. H. Gelfand, J. J. Sninsky, and T. J. White (ed.), **PCR Protocols. A Guide to Methods and Applications**. Academic Press, USA.
- ZHONG, S. and STEFFENSON, B.J. 2001. Virulence and molecular diversity in *Cochliobolus sativus*. **Phytopathology** 91:469–476.

Archive of SID