

COMPARISON AND DEVELOPMENT OF NEW METHOD FOR DETECTION OF *Erwinia amylovora* IN LATENT INFECTION PLANT MATERIAL *

C. MOSLEMKHANI** and L. SADEGHI¹

(Received: 24. 12. 2010; Accepted: 20. 7. 2011)

Abstract

Fire blight disease caused by *Erwinia amylovora* is deemed to be one of the most important and devastating disease affecting in Rosacea. Sensitive and rapid detection protocols are important tools in disease control. In this research sensitivity and specificity of four different detection methods with and with out enrichment were evaluated. 1CFU/ml *E. amylovora* could be detected by selective culture method in infected plant extract but not in latent infections. However, enrichment was increased the susceptibility of this method to detect *E. amylovora* in latent infections. Lateral flow immunochromatography detect 10^5 CFU/ml bacteria in infected plant extract and enrichment made this method to detect latent infections and increased the sensitivity to detect 1CFU/ml bacteria. Using specific primers, the standard and nested PCR were detected 1CFU/ml bacteria in infected plant material .

Keywords: *Erwinia amylovora*, Detection, PCR, Enrichment, Culture, Lateral flow immunochromatography.

See Persian text for figures and tables (Pages ۴۷۱-۴۷۴).

*: A Part of Research Project No. 2080888013 of Seed and Plant Certification and Registration Institute.

** : Corresponding Author, Email: moslemkhany@yahoo.com

1. Assis. Prof. and BSc. of Plant Pathology, Respectively, Seed and Plant Certification and Registration Institute.

References

- BARANAUSKAITE, L., JOGAITE, V. and JANKUVIENE, L. 2008. A three year study of fire blight in Lithuania. **Zemdirbyste -Agric.** 95 (3): 19–26
- BERESWILL, S., JOCK, S., BELLEMANN, P. and GEIDER, K. 1998. Identification of *Erwinia amylovora* by Growth Morphology on Agar Containing Copper Sulfate and by Capsule Staining with Lectin. **Plant Dis.** 82:158-164.
- CROSSE, J. E. and GOODMAN, R. N. 1973. A selective medium for and a definitive colony characteristic of *Erwinia amylovora*. **Phytopathology** 63:1425-1426.
- HENSON, J.M. and FRENCH, R. 1993. The polymerase chain reaction and plant disease diagnosis. **Annu. Rev. Phytopathol.** 31:81-109.
- ISHIMARU, C. and KLOS, E. J. 1984. New medium for detecting *Erwinia amylovora* and its use in epidemiological studies. **Phytopathology** 74: 13421345.
- LIOP, P., BONATERRA, A., PENALVER, J. and LOPEZ, MM. 2000. Development of a highly sensitive nested-PCR procedure using a single closed tube for detection of *Erwinia amylovora* in asymptomatic plant material. **Appl. Environ. Microbiol.** 66: 2071-2078.
- MCMANUS, P.S and JONES, A.C. 1995. Detection of *Erwinia amylovora* by nested PCR and PCR-dot blot and reverse blot hybridization. **Phytopathology** 85:618-623.
- MILLER, T. D. and SCHROTH, M. N. 1972. Monitoring the epiphytic population of *Erwinia amylovora* on pear with a selective medium. **Phytopathology** 62:1175-1182.
- VANDER ZWET, T. 2002. Present worldwide distribution of fire blight. **Acta Hort.** 590: 33–34.
- VANDER ZWET, T. and KEIL, H. L. 1979. Fire blight a bacterial disease of rosaceous plants. **Agricultural Handbook.** US Dept. of Agriculture, Washington DC. No 510.