short Article

EVALUATION OF PEROXIDASE ACTIVITY IN TWO RESISTANT AND SUSCEPTIBLE SESAME GERMPLASMS TO FUSARIUM DAMPING-OFF CAUSED BY *Fusarium oxysporum* f.sp. *sesami* *

A. FALLAHPORI¹, H. AMINIAN ^{1**} and S. A. ESMAILZADEH HOSSEINI²

(Received : 27.05.2011; Accepted : 07.11.2012)

Abstract

Sesame damping-off caused by *Fusarium oxysporum* f.sp. *sesami* is one of the most important diseases of *Sesamum indicum* L. in the world. In this work, the activity of peroxidase enzyme was assayed as a resistance mechanism in Asfij local germplasm (Bahabad, Yazd province) as resistant and Kahnoj local germplasm (Kerman province) as susceptible germplasm. Fifteen isolates of causal agent were isolated from sesame farms in Yazd province and one isolate with the highest pathogenicity potential was selected and used to determine hostrange. Among ten species of several plant families that were artificially inoculated with this pathogen, only sesame showed disease symptoms. Evaluation of peroxidase activity in tolerant and susceptible germplasms in 2, 4, 6, 8, 10 and 12 days after inoculation with *Fusarium oxysporum* f. sp. *sesami* showed that peroxidase enzyme activity increases in resistant germplasm with highest level in-4 days after inoculation. The activity of peroxidase was decreased 4 days after inoculation. In susceptible germplasm, enzyme activity was increased slightly in low quantity. In conclusion the results show that peroxidase activity play probable role in induction of plant resistance against outbreak of plant pathogens.

Keywords: Fusarium oxysporum f.sp. sesami, Local germplasm, Peroxidase.

See Persian text for figures and tables (Pages 117–11A).

^{*:} A Part of MSc. Thesis of The First Author, Submitted to University College of Aburaihan, The University of Tehran, Iran.

^{**:} Corresponding Author, Email: haminian@ut.ac.ir

^{1.} Former MSc. Student and Assoc. Prof of Plant Pathol., Respectively, University College of Aburaihan, The University of Tehran, Tehran, Iran.

^{2.} Res. Instructor of. Yazd Agricultural and Natural Resource Research Institute.

References

- BANIHASHEMI, Z. 1982. Evaluation of sesame germplasms for resistance to *Fusarium oxysporum*. Plant Pathol. 17: 75-79.
- BERNARD, M. A., SUMMERHURST, D. K., RAZEM, F. A. 2004. Oxidase, peroxidases and hydrogen peroxide : The suberin connection. **Phytochem. Rev.** 3: 113-126.
- BRADFORD, M. 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilization the principle of protein-dye binding. **Anal. Biochem.** 72: 248-254.
- BURGESS, L. W., SUMMERCELL, A., BULLOCK, S., GOTH, K. P. and BAKHOUSEe, D. 1994. Laboratory manual for Fusarium research. Department of Crop Science University of Sydney ND Royal Botanic Garden. 132 pp.
- CHENG, A. A. and Wu, F. R. 1991. Fusarium wilt of sesame in Taiwan. **Reporting TAINAN Station**. Council of Agriculture. 26: 53-60.
- DALLY, J. M., LUDDEN, P. and SEEVERS, P. 1971. Biochemical comparisons of resistance to wheat stem rust disease controlled by the sg6 or sr11 alleles. **Physiol. and Mol. Plant Pathol.** 1: 397-407.
- FALLAHPORI, A., AMINIAN, H., SAHEBANI, N. and ESMAILZADE HOSSEINI, S. A. 2010. Investigation of 10 sesame local germplasms to damping off caused by *Fusarium oxysporum* f.sp. *sesami*. Proc. 19th Iran. Plant Protec. Cong., Tehran, Iran. 307(Abst.).
- GAJHEDE, M. 2001. Plant peroxidase: substrate complexes with mechanistic implications. **Biochem. Soc.** 29: 91-195.
- JANDA, T., SZALAI, G., RIOS-GONZALES, K. and MOHAMMADI, M. 2003. Changes in specific activities of peroxidase, chitinase and phenylalanine ammonia-lyase and phenolic content in cucumber leaves inoculated with *Podosphaera fusca*, the causal agent of powdery mildew. Agric. Biol. Sci. 69: 545-553.
- KAVAK, K. and BOYDAK, E. 2006. Resistance levels of 26 sesame breeding lines to Fusarium wilt disease. Plant Pathol. 24: 141-146.
- KOBAYASHI, T. 1991. Cytogenetics of Sesame (Sesamum indicum). Elsevier Science Pub.Co. Inc., Amsterdam.
- MADHAIYAN, M., POONGUZHALI, S., SENTHIKUMER, M., SESHADRI, S., CHUNG, H., YONG, J., SUNDRAM, S. and SA, T. 2004. Growth promotion and induction of systemic resistance in rice cultivar Co-47 (*Oryzea sativa*) by *Methylbacterium spp.* Bot. Bull. Acad. Sinica 45: 315-324.
- MAGDA, M. A., SHOUSHA., A. WAFAA, G. and MAHDY, M. 2007. Biochemical alterations induced in tomato in response to *Fusarium oxysporum* infection purification and characterization of an acidic β-1,3-glucanasae. **Res. J. Agric. and Biol. Sci.** 3(6): 939-934.
- NELSON, PE., TOUSSOUN, TA. and MARASAS, WFO. 1983. Fusarium Species an Illustrated for Identification. The Pensylvania State University Press University Park. 193 pp.
- PINEDA, P. and AVILA, M. 1988. Alternative para el control de *Macrophomina phaseolina*, *Fusarium oxysporum*, pathogenos del ajonjoli (*Sesamum indicum* L.). Agronomia Tropical 38: 79-84.
- SEEVERS, P. M., DALLY, J. M. and CATEDRAL, F. F. 1971. The role of peroxidase isozymes in resistance to wheat stem rust disease. **Plant Physiol**. 48: 353-360.
- SIMOUSE, T.Y. and ROSS, A. F. 1995. Enhanced peroxidase activity associated with induction of resistance to tobacco mosaic virus in hypersensitive tobacco. **Phytopathology** 60: 383-384.
- YE, X. S., PAN, S. and KUC, J. 1990. Activity, isozyme pattern and cellular localization of peroxidase as related to systemic resistance of tobacco to blue mold (*Peronospora tabacina*) and to tobacco mosaic virus. **Physiol. and Biochem**. 80: 1295-1299.