Appl. Ent. Phytopath. Vol. 77, No. 1, Sep. 2009

Parasitoid wasp complex of *Phyllonorycter corylifoliella* (Lep.: Gracillariidae) in the Fars Province of Iran, and notes on their morphology and abundances

A. AMIRI¹, A. A. TALEBI^{1*}, P. NAVONE² and Z. YEFREMOVA³

1-Tarbiat Modares University, Tehran, Iran; 2- University of Torino, Italy
3- Ul'yanovsk State University, Russia
(Received: April 2008; Accepted: October 2008)

Abstract

The hawthorn red midget moth, *Phyllonorycter corylifoliella* (Lep.: Gracillariidae) is one of the important pests of apple trees in Fars Province. Parasitoid complex of this leafminer moth were investigated during 2006 and 2007. The leaves containing larvae and pupae were collected from five sites in Sarhad region and maintained in room conditions at temperature of 25±5°C until the parasitoids were emerged. A total of ten parasitoid species were reared: *Achrysocharoides suprafolius**, *Neochrysocharis longiventris**, *Zagrammosoma talitzkii, Sympiesis gordius, Sympiesis acalle**, *Sympiesis sericeicornis, Pnigalio agraules, Minotetrastichus frontalis**, *Baryscapus* sp. (Hym., Eulophidae) and *Pholetesor bicolor* (Hym., Braconidae). Among them, four species (marked with*) are new records for the fauna of Iran. The most common parasitoid was *A. suprafolius* (60.24% in 2006 and 68.98 % in 2007) followed by *Z. talitzkii* (14.20% in 2006 and 14.13% in 2007). Our new findings represent detailed evidence on parasitoid complex of *P. corylifoliella*. Notes on relative frequency, morphology, distribution and host range of the new parasitoid species records are included.

Key words: Phyllonorycter corylifoliella, Parasitoid, Apple, Fars Province.

^{*} Corresponding author: talebia@modares.ac.ir

چکیدہ

مینوز لکه تاولی روی برگ (Phyllonorycter corylifoliella (Lepidoptera: Gracillariidae) روی برگ (۲۰۹۵ مینوز از ۵ مکان یک آفت مهم در باغات سیب استان فارس است. پارازیتوئیدهای این آفت طی سالهای ۱۳۸۵ و ۲۸ مورد بررسی قرار گرفت. برگهای حاوی لارو و شفیرههای این گونه مینوز از ۵ مکان واقع در منطقه سرحد در شمال استان فارس جمع آوری و تا زمان ظهور پارازیتوئیدها در دمای ۵±۵۷ درجه سانتی گراه نگهداری شدند. پارازیتوئیدهای جمع آوری شده عبارتند از: Zagramosomma talitzkii Neochrysocharis longiventris ، Achrysocharoides suprafolus «Pnigalio agraules ، Sympiesis sericeicornis ، Sympiesis acalle ، Sympiesis gordius «Pnigalio agraules ، Sympiesis sericeicornis ، Sympiesis acalle ، Sympiesis forntalis » اوری شده عبارتند از خانواده عاده می اوری و تا و داده برای و داری را ز پر خانواده می می می می می از در میان آنها ٤ گونه (که با ستاره مشخص شدهاند) برای فون ایران گزارش جدید هستند. فراوانی نسبی گونهها، ویژگیهای مورفولوژیک گونههای جدید برای فون ایران، دامنه میزبانی و انتشار گونهها مورد بحث و بررسی قرار گرفته است.

Introduction

The genus *Phyllonorycter* includes 400 described species worldwide, with the greatest species richness in the palaearctic Region (285 species) (De Prins & De Prins, 2005). The *Phyllonorycter* species have been studied extensively and some of them are well known as pests of fruit orchards (Baggiolini, 1960; Potinger & Leroux, 1971). Outbreaks of leaf mining moths have occurred repeatedly in last decades and caused economic damage in apple orchards (Cross *et al.*, 1999). Heavy infestation can cause early leaf drop, reduction in terminal growth, small fruit size and premature ripening and fruit drop (Pottinger & Leroux, 1971). Until now, six species of the genus *Phyllonorycter* have been reported to feed in apple orchards in Europe: *Phyllonorycter blancardella* (F.), *P. mespilella* (Hübner), *P. pomonella* (Zeller), *P. gerasimovi* (Hering), *P. cerasicolella* (Herrich-Schäffer), and the howthorn red

^{*} **نشانی نگارندگان:** مهندس عباس امیری و دکتر علی اصغر طالبی، گروه حشره شناسی، دانشکده کشاورزی، دانـشگاه تربیت مدرس، صندوق پستی ۳۳۶–۱۴۱۱۵، تهران، ایـران؛ دکتـر پـائولو ناونــه، گـروه حــشره شناسـی و جانورشناسـی کاربردی، دانشگاه تورینو، ایتالیا؛ دکتر زویا یفرمووا، گروه جانورشناسی، دانشگاه ایالت آلیانوسک، روسیه.

midget moth (HRMM), *P. corylifoliella* (Hubner). Of these, only the larvae of HRMM make mines on the upper surface of leaves of rosaceous trees (Olivella, 1997). Acording to Radjabi (1986) three leafmining moth species of the genus *Phyllonorycter* including *P. blancardella*, *P. turanica* (Gerasimov) and HRMM with other related apple leafminer species such as *Leucoptera scitella* Zeller and *Stigmella malella* (Stainton) have been increasing important in majority regions of apple producing in Iran. The HRMM was firstly reported from Iran in 1970 (Radjabi, 1986) and gradually spread to across north, northwest and central regions (East Azarbaijan, Khorasan, Tehran, Markazi, Fars and Esfahan provinces) and probably distributed in other parts of Iran (Radjabi, 1986). Application of broad spectrum insecticides against apple key pests such as codling moth, *Cydia pomonella* (L.) (Lep., Tortricidae) and side effects of these compounds on beneficial arthropods and resistance to insecticides are the main reasons to increase the population density of leaf miner moths (Georghiou & Saito, 1983).

The HRMM has become an important pest in commercial apple orchards in Fars province (Amiri, 2008). It is frequent in apple orchards in Europe but has never been reported as an important pest in the area. The HRMM has four generations per year in the north of Fars Province conditions and hibernates as 5^{th} instar larvae (Amiri, 2008). Overwintering generation emerged in early spring and adult females deposited their eggs singly on the upper surface of the leaves (Radjabi, 1986; Amiri, 2008).

Natural enemies of leaf miner moths were investigated in the world and among them, hymenopterous parasitic wasps are valuable components in maintaining ecological balance of these insects (Lasalle & Gauld, 1993). Acording to Gates *et al.* (2002) eulophid wasps are consisted over 80% of leaf miner parasitoids. In northeastern Spain, the dominant parasitoids of HRMM were *Sympiesis gordius* (Walker), *S. acalle* (Walker) and *P. bicolor* (Nees) (Bellostas *et al.*, 1998) while, in Hungary, the more abundant parasitoid species was *S. sericeicornis* (Balazs, 1989).

Here, we report the results of a 2-year survey of the hymenopterous parasitoids of HRMM in the north of Fars Province of Iran. This research is a part of a larger project for biological control program against HRMM. As the first important step in this process, we try to identify, re-describe and illustrate newly recorded parasitoid species of HRMM from Fars province. Descriptions of the newly recorded species are provided to help Iranian researchers in identifying those species.

Materials and Methods

Parasitoid complex of HRMM were investigated during May to October 2006 and 2007 in the Sarhad region, (52°, 12' E and 30°, 43' N), located in the north of Fars province of Iran. Parasitoid complex were collected by weekly sampling of infested leaves in five apple orchards in the Sarhad region at five sites: Sedeh, Tange Boragh, Bande Bahram, Baseri, Bakan, within 80 Km radiuses. On each sampling date, 100 infested leaves were selected from each orchard. Samples were placed into clean plastic bags and transferred to the laboratory. Each sample (100 infested leaves) was placed in a plastic culture container (25×10×8 cm) and covered with fine-mesh nylon. The rearing boxes were maintained in room temperature (25±5 °C) for at least two months. The rearing boxes were checked daily, and parasitoid adults were collected. The specimens were preserved in 75% ethanol. The parasitoid species were confirmed by the fourth and fifth authors of this paper (Z. Yefremova and P. Navone). Some of the ethanol-preserved specimens were mounted on slides for microscopic study in Canada balsam using the techniques outlined in Noyes (1982).

The external morphology of the parasitoid was illustrated using a phase-contrast OLYMPUS BH2 microscope with a drawing tube. The morphological terminology used in the parasitoid species morphology is based on Gibson *et al.* (1997) and Noyes (2007). The synonym names of the newly recorded species were standardized according to Noyes (2007). The material is deposited mainly in the collection of the Department of Entomology, College of Agriculture, Tarbiat Modares University, Tehran Iran, with duplicate in the Department of Zoology, Ul'yanovsk State University, Russia, and in the University of Torino, Italy.

Result and discussion

A total of 10 hymenopterous parasitoid species were reared from HEMM in the studied area during May to October 2006 and 2007: *Achrysocharoides suprafolius* (Askew & Ruse), *Neochrysocharis longiventris* (Askew), *Zagrammosoma talitzkii* (Boucek), *Sympiesis gordius* (Walker), *Sympiesis acalle* (Walker), *Sympiesis sericeicornis* (Nees), *Pnigalio agraules* (Walker), *Minotetrastichus frontalis* (Nees), *Baryscapus* sp. (Hym., Eulophidae) and *Pholetesor bicolor* (Hym., Braconidae). Among them four parasitoid species: *A. suprafolius*, *S. acalle, N. longiventris* and *M. frontalis* are new records for Iran.

A. suprafolius is found to be the most common parasitoid (60.24 % and 68.98% in 2006 and 2007, respectively) followed by *Z. talitzkii* (14.20% and 14.13% in 2006 and 2007, respectively). The other parasitoids, e.g. *S. gordius*, *S. acalle*, *N. longiventris*, *P. agraules* had

moderately abundant and the remaining species, *Baryscapus* sp., *M. frontalis*, S. *sericeicornis*, and *A. bicolor* occurred in small numbers (Table 1). Previous studies on the leafminer moths have shown that chalcidoid wasps usually play an important role in regulating of these pests population and level of parasitism in some cases reach over 50% (Askew and Shaw, 1979; Gibogini *et al.*, 1996). Parasitoids of HRMM were identified in Northern Spain and their dominant parasitoids were *Symplesis gordius*, *S. acalle* and *Pholetesor bicolor* (Bellostas *et al.*, 1998).

Parasitoids	Number of parasitoid specimens		Relative frequency (%)	
	2006	2007	2006	2007
Achrysocharoides suprafolius	509	576	60.24	68.98
Zagrammosoma talitzkii	120	114	14.20	13.65
Sympiesis gordius	56	17	6.63	2.04
S. sericeicornis	2	-	0.24	_
S. acalle	47	31	5.56	3.71
Neochrysocharis longiventris	47	67	5.56	8.02
Pnigalio agraules	41	10	4.85	1.19
Baryscapus sp.	8	4	0.95	0.47
Minotetrastichus frontalis	7	3	0.83	0.36
Pholetesor bicolor	8	13	0.95	1.56

Table 1- Frequency of parasitoid complex of Phy	llonoeycter corylifoliella
in Fars province of Iran in 2006 a	and 2007.

Notes on the Newly Recorded Parasitoid Species

Achrysocharoides suprafolius (Askew, 1974)

Syn.: Enaysma suprafolia Askew, 1974

Diagnosis: Female: Body length 1.2-1.5 mm; vertex golden green, face brown, compound eyes red, frontal suture T-shape; mandibles tridentate; antenna 7-segmented, scrobe reticulated, antennal segments yellowish to white, scape length 3x as long as pedicel, funicle 3-segmented, clava 2-segmented (Fig. 5), antenna with three discoid anelli in both sex; thorax and abdomen metallic green, pronotal collar without carina, antero-median part of

propodeum smooth without carina (Fig. 12), thorax included propodeum 0.86x as long as abdomen; legs white and pretarsus pale brown; post marginal vein as long as stigmal vein, fore wing 2.04 as long as wide, ratio of submarginal/marginal/postmarginal/stigmal veins is 0.9: 2.3: 0.3: 0.27 (Fig. 1); **Male**: Body length 1.1-1.3 mm; resembles female but its antennal segment paler than female, width of antennal scape about 2x as female (Fig. 6).

Distribution and host range: *A. suprafolius* was the most abundant parasitoid of HRMM in the studied area (Table 1). This species is distributed in Germany, the Netherlands and United Kingdom (Noyes, 2007). *Achrysocharoides* species attack *Phyllonorycter* larvae that mine lower surfaces of leaves, except for *A. suprafolius*, which feeds only on the HRMM larvae (Askew & Ruse, 1974).

Sympiesis acalle (Walker, 1848)

Syn.: Astichus bimaculatipennis Girault, 1912; Entedon nubeculatus Ratzeburg, 1848; Eulophus acalle Walker, 1848; Eulophus bifasciatus Thomson 1878; Sympiesis bimaculata Crawford, 1913; Sympiesis bimaculatipennis (Girault, 1912); Sympiesis meteori Girault, 1916.

Diagnosis: Female: Body length 2.5-2.9 mm; body dark green with bluish tint; antennae 8 segmented in both sexes, scape yellowish to white, with black dorsal surface, funicle 4 segmented, clava 2-segmented (Fig. 7); thorax 2x as long as wide, propodeum with complete median carina (Fig. 13); coxa and femur in all legs brownish to black; fore wings with two dark strips, post marginal vein about 2x as long as stigmal vein (Fig. 2); **Male**: Body length 1.8–2 mm; three first segments of funicle with branch–like projections (Fig. 8), forewings without dark stripes.

Distribution and host range: This species was reported from Azerbaijan, Bosnia, Bulgaria, Canada, United Kingdom, Czech Republic, Germany, Hungary, Italy, Japan, Korea, Rumania, Russia and the Netherlands (Noyes, 2007). This species is a parasitoid of *Phyllonorycter* (Lepidoptera: Gracillariidae) and *Agromyza* (Diptera: Agromyzidae) (Boucek & Askew, 1968; Yefremova, 2007).

Neochrysocharis longiventris (Askew, 1979)

Syn.: Chrysonotomyia longiventris Askew, 1979

Diagnosis: Female: Body length 0.9–1.2 mm; Body dark green, anterior margin of clypeus almost straight, mandibles bidentate and brown; eyes with pubescence, hairs minute and visible only under high magnification; scape somewhat paler than flagellum and palest at base, pedicel in profile 1.6x as long as broad, clava 3-segmented with an apical spine, longer

than funicle segments plus anelli (Fig. 11); thorax somewhat flattened, scutellum very slightly longer than broad with one pair of long setae, thorax in combination with propodeum 1.4x as long as broad in dorsal view (Fig. 14); gaster with dark green reflection on the first tergite; ovipositor dark brown; wings hyaline with a small fuscous spot extending from stigma, venation brownish, fore wings 1.9x as long as broad (Fig. 3). coxa green metallic, *N. longiventris* is very similar to *Closterocerus formosus* in morphological chrachteristics but *N. longiventris* have relatively longer gaster; **Male**: Body length 0.7- 1.1 mm; resembles female, but gaster trapezoidal and its length about as long as rest of body, antennae rather longer and more slender.

Distribution and host range: *N. longiventris* was reported from Denmark, United Kingdom (Askew, 1979), Sweden (Hansson, 1990). It is a polyphagous species and reared as parasitoid of Coleoptera (Curculionidae) and Lepidoptera (such as *Phyllonorycter maestingella* Zeller, *P. quinnata* (Stainton), *P.tenella* (Joannis) (Noyes, 2007)). *N. longiventris* is recorded here for the first time as parasitoid of HRMM. Species of the genus *Neochrysocharis* developed as solitary or gregarious endoparasites in immature stage of mainly phytophagous insects (Hansson, 1990).

Minotetrastichus frontalis (Nees, 1834)

Syn.: Aprostocetus budensis (Erdös, 1954); Aprostocetus ecus (Walker, 1839); Cirrospilus ecus Walker, 1839; Entedon cyclogaster (Ratzeburg, 1844); Entedon rivillellae Rondani, 1877; Entedon xanthops (Ratzeburg, 1844); Eulophus cyclogaster Ratzeburg, 1844; Eulophus frontalis Nees, 1834; Eulophus xanthops Ratzeburg, 1844; Geniocerus budensis Erdös, 1954; Geniocerus cyclogaster (Ratzeburg, 1844); Geniocerus ecus (Walker, 1839); Geniocerus xanthops (Ratzeburg, 1844); Minotetrastichus ecus (Walker, 1839); Tetrastichus cimbicis Kostjukov, 1976; Tetrastichus cyclogaster (Ratzeburg, 1844); Tetrastichus cyclogaster obscurata Ruschka, 1924; Tetrastichus ecus (Walker, 1839)

Diagnosis: Female: Body length 0.8–0.9 mm; head whitish to yellow; antennae 8-segmented, funicle 3-segmented, clava 3-segmented (Fig. 10), space between ocelli dark brown, antennae pale brown, dorsal surface of pedicel dark brown; thorax metallic green, length of thorax including propodeum 0.72x as long as abdomen, length of thorax 1.6x as long as wide (Fig. 15); first three segments of abdomen whitish to yellow, but dorsal surface of three terminal segments is dark brown; legs white; length of fore wings 2.19x as long as wide, postmarginal vein considerably reduced (Fig 4); **Male**: Body length 0.67–0.8 mm; antennae 9-segmented, funicle 4-segmented with branch–like projection (Fig. 9).



Fig. 1-4. Forewings, 1. Achrysocharoides suprafolius ; 2. Symplesis acalle;
3. Neochrysocharis longiventris; 4. Minotetrastichus frontalis;
Scale bars=50 micrometers (original)



Fig. 5-11- Antennae, 5. Achrysocharoides suprafolius ♀, 6. ♂; 7. Symplesis acalle ♀,
8. ♂; 9. Minotetrastichus frontalis, ♂, 10. ♀; 11. Neochrysocharis longiventris ♀; scale bars=50 micrometers (original)



Fig. 12-15. Thorax, 12. Achrysocharoides suprafolius; 13. Sympiesis acalle; 14. Neochrysochaeis longiventris 15. Minotetrastichus frontalis; Scale bars=50 micrometers (original)

Distribution and host range: *M. frontalis* is recorded here for the first time from Iran. It is one of the most common and polyphagous parasitoids of various groups of leafminers. It is distributed in Europe, Canada and United States of America (Noyes, 2007). It was reared from *P. corylifoliella*, *P. blancardella* (Tomov, 2002), *P. robiniella* and *C. ohridella* (Grabenweger, 2003).

Z. talitzkii and *P. agraules* have recently been reported from Iran (Yefremova *et al.*, 2007). *P. agraules* was also reported from Austria, Bosnia, Bulgaria, France, Germany, Greece, Italy, North Africa, Russia, Spain, Turkey and Ukraine (Noyes, 2007). It is a solitary ectoparasitoid and a common species with a wide host range (Boucek & Askew, 1968). This parasitoid is one of the larger parasitoids attacking *Cameraria ohridella* (Grabenweger, 2003; Freise *et al.*, 2002). This species has already been reported as parasitoid of *P. coryliofoliella* (Noyes, 2007) and *P. robiniella* (Grabenweger, 2003).

Z. talitskii was the second dominant species in the parasitoid complex. This species is distributed in Italy, Kazakhstan, Moldova, Russia, Turkmenistan, Ukraine, eastern USSR (Noyes, 2007). This parasitoid has also been reported as parasitoid of HRMM in Moldavia and Ukraine (Boucek & Askew 1968). It is a common primary ectoparasitoid of leafminer moths (Gracillariidae, Lyonetiidae, Phyllocnistidae) and may also attack other lepidopterous and dipterous leafminers (Noyes, 2007; Yefremova, 2007).

In conclusion it should be noted that the reduction of broad-spectrum insecticides applications in apple orchards led to the rich diversity of parasitic wasps. Several studies have shown that parasitoid populations are much larger in untreated orchards than in orchards sprayed with insecticides (Cross *et al.*, 1999). It can be asserted that parasitoids have a great effect on population density of the leaf miner moths, but more attention should be given to the knowledge of the biology and ecology of parasitoid species to better use them in biological control programs.

Acknowledgments

The financial support for this work was provided by Tarbiat Modares University. The authors wish to thank to Prof. Dr. K. Kamali (Tarbiat Modares University, Tehran) and two anonymous reviewers for their valuable comments and suggestions on the manuscript and Dr. A. Lozan (Institute of Entomology, Ceske Budejovice, Czech Republic) for identification of *P. bicolor*. We also thank to Mr. Ahmad Amiri for valuable assistance with the collections of specimens.

References

- AMIRI, A. 2008. Biology, population dynamics and seasonal parasitism of *Phyllonorycter corylifoliella* (Lep.: Pyralidae) in north of Fars province, Iran. M.Sc. thesis, Tarbiat Modares University.
- ASKEW, R. R. 1979. Taxonomy of some European *Chrysontomyia* Ashmead (Hymenoptera: Eulophidae) with description of *C. longiventris* n. sp. and notes on distribution. *Entomol Scand.*, 10: 27-31.
- ASKEW, R. R. 1984. Species of *Pnigalio* and *Chrysocharis* (Hymenoptera: Eulophidae) parasitic on Tischeriidae (Lepidoptera), with the description of a new species. *Entomol. Gazette.*, 35: 103-109.
- ASKEW, R. R. and J. M. RUSE, 1974. Biology and taxonomy of species of the genus *Enaysma* Delucchi (Hym., Eulophidae, Entedoninae) with special reference to the British fauna. *Trans. Roy. Ent. Soc. Lond.*, 125 (3): 257-294.
- ASKEW, R. R. and M. R. SHAW, 1979. Mortality factor affecting the leafmining stage of *Phyllonorycter* (Lepidoptera: Gracillariidae) on oak and birch. The analysis of the mortality factors. *Zool. J. Linn. Soc.*, 76: 31-49.
- ASKEW, R. R. and M. R. SHAW, 1986. Parasitoid communities: their size, structure and development. in: J. Waage and Greathead (eds), Insect Parasitoids. Academic Press, London, pp. 225-264.
- BAGGIOLINI, M. 1960. Observations sur la biologie de deux mineuses du genre Lithocolletis: L. corylifoliella et L. blancardella (Lep. Gracillariidae), nuisibles aux arbres fruitiers en Suisse roman de. Mitt. Schw. Entomol. Ges. 32(4): 385-397.
- BALAZS, K. 1989. Zur Populations Dynamik von Miniermotten und ihren Parasiten in Apfelanlagen. Akademie Landwirtschafts-Wissenschaften DDR, Berlin 278, 185-191.
- BELLOSTAS, J. J., E. OLIVELLA, M. J. VERDU, M. J. SARASUA and J. AVILLA, 1998. Fauna parasitoidede *Phyllonorycter* spp. en plautaciones de manzano de Lleida. *Bol. Sanid. Veget. Plagas*, 24: 313-320.
- BOUCEK, Z. and R. R. ASKEW, 1968. Index of palaeartic Eulophidae (excl. Tetrastichinae).In: Delucchi. V, Remaudiere. G (eds) Index of entomophagous insect 3: 260pp, Le Francois, Paris, France.
- CROSS, J. V., M. G. SOLOMON, D. BABANDREIER, L. BLOMMERS, M. A. EASTERBROOK, C. N. JAY, G. JENSER, R. L. JOLLY, U. KUHLMANN, R.

LILLEY, E. OLIVELLA, S. TOEPFER and S. VIDAL, 1999. Biocontrol of Pests of Apples and Pears in Northern and Central Europe: 2. Parasitoids. *Biocon. Sci. Technol.*, 9: 277-314.

- DE PRINS, W. and J. DE PRINS, 2005. Family Gracillariidae. In Landry, B. (Ed.), *World Catalogue of Insects*, vol. 6. 502 pp. Apollo Books, Stenstrup
- FREISE, J., F. HEITLAND and I. TOSEVSKI, 2002. Parasitism of the horse chestnut leafminer, *Cameraria ohridella* Deschka and Dimic (Lep: Gracillariidae) in Serbia and Macedonia. J. Pest. Sci., 75:152-157.
- GATES, M. W., J. M. HERATY, D. L. WAGNER and J. B. WITHFIELD, 2002. Survey of the parasitic hymenoptera on leafminers in California. J. Hym. Res., 2: 213-270.
- GEORGHIOU, G. P. and T. SAITO, 1983. Pest resistance to pesticides. Plenum Press, New York and London. 809 pp.
- GIBOGINI, B., A. ALMA and A. AROZONE, 1996. Bio-ethological researches on the hymenoptera of the biocenosis of *Phyllonorycter robiniellus* (Clemens) (Lepidoptera: Gracillariidae). *Boll. Zool. Agr. Bachic.*, 28:13-22.
- GIBSON, G. A. P., J. T. HUBER and J. B. WOOLLEY, 1997. Annotated Keys to the Genera of Nearctic Chalcidoidea (Hymenoptera). National Research Council Canada, NRC Research Press, 794 pp
- GRABENWEGER, G. 2003. Poor control of the horse chestnut leafminer *Cameraria ohridella* (Lepidoptera: Gracillariidae) by native parasitoids: a synchronization problem. *Eur. J. Entomol.*, 101:189-192.
- HANSSON, C. 1983. Taxonomic notes on the genus Achrysocharoides Girault, 1913 (Hymenoptera: Eulophidae), with a redescription and a description of a new species. *Entomol. Scand.*, 14: 281-291.
- HANSSON, C. 1990. A taxonomic study on the Palaearctic species of *Chrysonotomyia*Ashmead and *Neochrysocaris* Kurdjumov (Hymenoptera: Eulophidae). *Entomol.Scand.*, 20: 29-52.
- LASALLE, J. and I. D. GAULD, 1993. Hymenoptera and Biodiversity. Cab International.Wallingford. Oxou OX 10 8DE, U.K. Nees ab F.senbeck, C. G. 1834. Hymenopterum Ichneomonidus afinium monograohiae genera europaea et species illustrantes, 2: 1-448.
- NOYES, J. 1982. Collecting and preserving chalcid wasps (Hymenoptera: Chalcidoidea). J. Nat. Hist., 16: 315-334

NOYES, J. 2007. Universal Chalcidoided Database. The natural History Meuseum, London,

UK, Available on the: http://www.nhm.ac.uk/ entomology/chalcidoids (Accessed May 2008).

- OLIVELLA, E. 1997. *Phyllonorycter mespilella*, (Hubner, 1805) new species for the Iberian fauna (Lepidoptera: Gracillariidae) SH ILAP, *Revista de Lepidopterologia*. 25, 37-42
- POTTINGER, R. P. and E. J. LEROUX, 1971. The biology and dynamics of *Lithocolletis* blancardella (Lepidoptera: Gracillariidae) on apple in Quebec. *Mem. Ent. Soc. Can.*, 77: 1-437.
- RADJABI, G. H. 1986. Insects attacking rosaceous fruit trees in Iran, Second Volume, Lepidoptera. Plant Pests and Diseases Research Institute, Tehran, Iran, 209pp.
- TOMOV, R. 2002. Species composition of parasitoids (Hymenoptera) on apple feeding *Phyllonorycter* (Lepidoptera: Gracillariidae) in Bulgaria. IN: Melika, G. and Thuroczy, C., eds., Pararasitic wasps: evolution, systematics, biodiversity and biological. International Symposium on Parasitic Hymenoptera: Taxonomy and Biological Control, Köszeg, Hungary 439 pp.
- YEFREMOVA, Z. A. 2007. The subfamilies Eulophinae, Euderinae and Entedoninae (Hymenoptera: Eulophidae) in Yemen. Fauna of Arabia, 23: 335-368.
- YEFREMOVA, Z. A., E. EBRAHIMI and E. YEGORENKOVA, 2007. The Subfamilies Eulophinae, Entedoninae and Tetrastichinae in Iran, with description of new species (Hymenoptera: Eulophidae). *Entomofauna*, 28: 321-356.

Address of the authors: Eng. A. AMIRI and Dr. A. A. TALEBI, Departement of Agricultural Entomology, Faculty of Agriculture, Tarbiat Modares University, P. O. Box 14115-336, Tehran, Iran; Dr. P. NAVANE, Di.Va.P.R.A. Entomologia e Zoologia applicate all'Ambiente, University of Torino, via L. da Vinci 44, 10095 Grugliasco (TO), Italy; Dr. Z. YEFREMOVA, Department of Zoology, Ul'yanovsk State University, pl. 100-letiya Lenina, 4, RUœ4327000 Ul'yanovsk, Russia.