# THE MAXIMUM NUMBER OF EDGES IN A STRONGLY MULTIPLICATIVE GRAPH

#### R. KAFSHGARZAFERANI

Communicated by Cheryl Praeger

ABSTRACT. We derive a formula for the maximum number of edges in a strongly multiplicative graph as a function of its order.

Recently, L.W. Beineke and S.M. Hegde [3] introduced the notion of a strongly multiplicative graph.

**Definition** (Beineke, Hegde [3]). A graph with n vertices is said to be strongly multiplicative if its vertices can be labeled 1, 2, ..., n, so that the values on the edges, obtained as the product of labels of the end vertices, all are distinct.

An interesting problem is to obtain a formula for the maximum number of edges  $\lambda(n)$  for a strongly multiplicative graph of order n. In [3], Beineke and Hegde gave an upper bound for  $\lambda(n)$ . In [2], C. Adiga et al. obtained a sharper upper bound for  $\lambda(n)$ . Then in [1], Adiga et al. established a formula for  $\lambda(n)$  in terms of the divisor function. We quote their result now.

MSC(2000): Primary 05C78

Keywords: Graph labelling, Strongly multiplicative graphs

Received: 4 May 2006, Revised: 21 July 2006

© 2005 Iranian Mathematical Society.

54 Kafshgar Zaferani

**Theorem** (Adiga et al. [1]).

$$\lambda(n) = \sum_{k=1}^{n(n-1)} g(k),$$

where

$$g(k) = \min\{1, f(k)\},\$$

$$f(k) = \begin{cases} \left[\frac{d(k)}{2}\right] & \text{if } 1 \le k \le n, \\ \left[\frac{d(k)}{2}\right] - d_n(k) & \text{if } n < k \le n(n-1) \end{cases}$$

and where d(k) denotes the number of distinct divisors of k, [x] denotes the largest integer less than or equal to x, and  $d_n(k)$  denotes the number of divisors of k greater than n.

In this note we derive the following formula for  $\lambda(n)$ .

Theorem.

$$\lambda(n) = \frac{n(n-1)}{2} + \sum_{m=2}^{n} \sum_{k=1}^{m-1} \left[ -\frac{\theta(m,k)}{\left[\sqrt{mk-1}\right] - k + 1} \right],$$

where

$$heta(m,k) = \sum_{s=k+1}^{\left[\sqrt{mk-1}\right]} \left[\frac{\left[\frac{mk}{s}\right]}{\frac{mk}{s}}\right].$$

**Proof.** Let 
$$\delta(n) = \lambda(n) - \lambda(n-1)$$
. Then 
$$\lambda(n) = \sum_{m=2}^{n} \delta(m). \tag{2.1}$$

Thus, in view of (2.1) it is enough to obtain a formula for  $\delta(m)$ . Consider the array of products

1.2 1.3 1.4 ... 1.
$$(n-1)$$
 1. $n$ 
2.3 2.4 ... 2. $(n-1)$  2. $n$ 
3.4 ... 3. $(n-1)$  3. $n$ 

$$(n-2).(n-1)$$
  $(n-2).n$ 

Let  $A_k$  denote the set of all elements of the  $k^{th}$  row. We count the number of terms in the last column which appear in other rows. If k.nis divisible by s  $(k \leq s-1)$ , then there exists an m < n such that k.n = s.m, and hence k.n repeats in the  $s^{th}$  row, i.e.  $k.n \in A_s$ . Observe that k.n may belong to  $A_s$ , where s is the largest integer such that  $k+1 < s < \sqrt{kn}$ . Thus the number of repetition of k.n in these rows is

$$\theta(n,k) = \sum_{s=k+1}^{\left[\sqrt{kn-1}\right]} \left[\frac{\left[\frac{nk}{s}\right]}{\frac{nk}{s}}\right].$$

By the definition of  $\theta(n,k)$ , it is clear that

$$0 \le \theta(n,k) \le \left[\sqrt{kn-1}\right] - k + 1$$

Thus

$$0 \le \theta(n,k) \le \left[\sqrt{kn-1}\right] - k + 1.$$

$$\left[-\frac{\theta(n,k)}{\left[\sqrt{nk-1}\right] - k + 1}\right] = \begin{cases} -1 & \text{if } kn \in \bigcup_{s=k+1}^{\sqrt{kn-1}} A_s, \\ 0 & \text{otherwise.} \end{cases}$$

This implies that

$$\delta(n) = (n-1) + \sum_{k=1}^{n-1} \left[ -\frac{\theta(n,k)}{\left[\sqrt{nk-1}\right] - k + 1} \right]. \tag{2.2}$$

On using (2.2) in (2.1), we complete the proof.

56 Kafshgar Zaferani

# Acknowledgment

I am thankful to the referee for his valuable suggestions and comments which improved the quality of the paper. I also wish to thank Dr C. Adiga for his suggestions and help during the preparation of this paper.

## REFERENCES

- [1] C. Adiga, H.N. Ramaswamy and D.D. Somashekara, On the strongly multiplicative graphs, South East Asian J.Math. and Math.Sci. 2 (2003), 45-47.
- [2] C. Adiga, H.N. Ramaswamy and D.D. Somashekara, A note on strongly multiplicative graphs, *Discuss. Math. Graph Theory* **24** (2004), 81-83.
- [3] L.W. Beineke and S.M. Hegde, Strongly multiplicative graphs, *Discuss. Math. Graph Theory* **21** (2001), 63-76.

### R. Kafshgarzaferani

Department of Studies in Mathematics University of Mysore MYSORE - 570 006, INDIA

e-mail:rkzaferan@yahoo.com