AMENABILITY AND WEAK AMENABILITY OF TRIANGULAR BANACH ALGEBRAS

A. R. MEDGHALCHI, M. H. SATTARI AND T. YAZDANPANAH

ABSTRACT. Let \mathcal{A} and \mathcal{B} be Banach algebras and let \mathcal{X} be a Banach \mathcal{A}, \mathcal{B} -module. Let $\mathcal{T} = \begin{bmatrix} \mathcal{A} & \mathcal{X} \\ \mathcal{B} \end{bmatrix}$ be the corresponding triangular Banach algebra. Forrest and Marcoux have studied the n-weak amenability of triangular Banach algebras. We show that when \mathcal{A} has a bounded approximate identity and \mathcal{X} is essential, then \mathcal{T} is weakly amenable if and only if \mathcal{A} and \mathcal{B} are weakly amenable. We also study the amenability of triangular Banach algebras and show that \mathcal{T} is amenable if and only if \mathcal{A} and \mathcal{B} are amenable and $\mathcal{X} = \{0\}$.

1. Introduction

Let \mathcal{A} and \mathcal{B} be Banach algebras and \mathcal{X} be a Banach \mathcal{A} , \mathcal{B} -module. That is, \mathcal{X} is a left Banach \mathcal{A} -module, a right Banach \mathcal{B} -module, (ax)b = a(xb) for $a \in \mathcal{A}$, $b \in \mathcal{B}$, $x \in \mathcal{X}$ and there exists a constant k0 such that

$$\parallel axb \parallel \leq k \parallel a \parallel \parallel x \parallel \parallel b \parallel.$$

 \mathcal{X} is said to be essential provided that for every $x \in \mathcal{X}$ there are $a \in \mathcal{A}$, $b \in \mathcal{B}$ and $y, z \in \mathcal{X}$ such that x = ay = zb. Let \mathcal{X}^* be the topological dual of \mathcal{X} . Then \mathcal{X}^* is a Banach \mathcal{B} , \mathcal{A} -module via the following actions

$$\langle x, bx^* \rangle = \langle xb, x^* \rangle, \qquad \langle x, x^*a \rangle = \langle ax, x^* \rangle$$

MSC(2000): Primary 46H25, 16E40

Keywords: Amenability, n-weak amenability, Triangular Banach Algebra

Received: 17 December 2003 , Revised: 04 October 2005 , Revised: 05 July 2006

© 2005 Iranian Mathematical Society.

for all $a \in \mathcal{A}$, $b \in \mathcal{B}$, $x \in \mathcal{X}$ and $x^* \in \mathcal{X}^*$.

For $x \in \mathcal{X}$ and $x^* \in \mathcal{X}^*$ we define $xx^* \in \mathcal{A}^*$ and $x^*x \in \mathcal{B}^*$ by

$$\langle a, xx^* \rangle = \langle ax, x^* \rangle, \ \langle b, x^*x \rangle = \langle xb, x^* \rangle \ (a \in \mathcal{A}, b \in \mathcal{B}).$$

Similarly for $x \in \mathcal{X}$, $F_2 \in \mathcal{A}^{**}$ and $G_2 \in \mathcal{B}^{**}$ we define $F_2 x \in \mathcal{X}^{**}$ and $xG_2 \in \mathcal{X}^{**}$ via the actions (c.f.[6])

$$\langle x^*, F_2 x \rangle = \langle xx^*, F_2 \rangle, \ \langle x^*, xG_2 \rangle = \langle x^*x, G_2 \rangle \ (x^* \in \mathcal{X}^*).$$

We may continue this process to higher order dual spaces of \mathcal{X} , and $\mathcal{X}^{(2n)}$ is a Banach \mathcal{A} , \mathcal{B} -module, $\mathcal{X}^{(2n-1)}$ is a Banach \mathcal{B} , \mathcal{A} -module, $\mathcal{A}^{(2n)}\mathcal{X}\subseteq \mathcal{X}^{(2n)}$, $\mathcal{X}\mathcal{B}^{(2n)}\subseteq \mathcal{X}^{(2n)}$, $\mathcal{X}\mathcal{X}^{(2n-1)}\subseteq \mathcal{A}^{(2n-1)}$ and $\mathcal{X}^{(2n-1)}\mathcal{X}\subseteq \mathcal{B}^{(2n-1)}$ for all n>0.

A Banach \mathcal{A} , \mathcal{B} -module \mathcal{X} is called non-degenerate if $\mathcal{A}x = \{0\}$ implies x = 0 and $x\mathcal{B} = \{0\}$ implies x = 0 for all $x \in \mathcal{X}$. When \mathcal{A} and \mathcal{B} have bounded approximate identities and \mathcal{X} is essential, then \mathcal{X} is a non-degenerate Banach \mathcal{A} , \mathcal{B} -module. Also when \mathcal{X} is essential, then \mathcal{X}^* is a non-degenerate Banach \mathcal{B} , \mathcal{A} -module.

Let \mathcal{X} be a Banach \mathcal{A} -bimodule. A derivation $\delta: \mathcal{A} \longrightarrow \mathcal{X}$ is a linear map such that $\delta(ab) = \delta(a)b + a\delta(b)$ for all $a, b \in \mathcal{A}$. The derivation δ is inner if it is of the form $\delta(a) = \delta_x(a) := ax - xa$ for some $x \in \mathcal{X}$. The linear space of all bounded derivations from \mathcal{A} to \mathcal{X} is denoted by $\mathcal{Z}^1(\mathcal{A}, \mathcal{X})$ and the linear subspace of all inner derivations by $\mathcal{N}^1(\mathcal{A}, \mathcal{X})$. The first Hochschild cohomology group of \mathcal{A} with coefficients in \mathcal{X} is defined to be the linear space $\mathcal{H}^1(\mathcal{A}, \mathcal{X}) = \mathcal{Z}^1(\mathcal{A}, \mathcal{X})/\mathcal{N}^1(\mathcal{A}, \mathcal{X})$ [15]. A Banach algebra \mathcal{A} is said to be amenable if $\mathcal{H}^1(\mathcal{A}, \mathcal{X}^*) = \{0\}$ for every Banach \mathcal{A} -bimodule \mathcal{X} (see, [1], [2], [3], [9], [10], [11], [12], [13]). A Banach algebra \mathcal{A} is weakly amenable if $\mathcal{H}^1(\mathcal{A}, \mathcal{A}^*) = \{0\}$ ([1], [14], [17], [18], [19]) and \mathcal{A} is called n-weakly amenable if $\mathcal{H}^1(\mathcal{A}, \mathcal{A}^{(n)}) = \{0\}$, where $\mathcal{A}^{(n)}$ is the n-th dual module of \mathcal{A} when $n \geq 1$ and is \mathcal{A} itself when n = 0 ([4]).

Forrest and Marcoux in [7] have studied a class of Banach algebras, which is called triangular Banach algebras. They have studied the n-weak amenability of triangular Banach algebras in [8]. They consider the cases where \mathcal{A} and \mathcal{B} have units and \mathcal{X} is unital Banach \mathcal{A} , \mathcal{B} -module.

Let \mathcal{A} and \mathcal{B} be Banach algebras and \mathcal{X} be a Banach \mathcal{A} , \mathcal{B} -module. We define the corresponding triangular Banach algebra $\mathcal{T} = \begin{bmatrix} \mathcal{A} & \mathcal{X} \\ \mathcal{B} \end{bmatrix}$ with the usual 2×2 matrix operations and obvious interval module actions, and the norm

$$\parallel \begin{bmatrix} a & x \\ 0 & b \end{bmatrix} \parallel = \parallel a \parallel + \parallel x \parallel + \parallel b \parallel.$$

In this paper \mathcal{A} and \mathcal{B} are Banach algebras, \mathcal{X} is a Banach \mathcal{A},\mathcal{B} - module and $\mathcal{T} = \begin{bmatrix} \mathcal{A} & \mathcal{X} \\ \mathcal{B} \end{bmatrix}$ is the corresponding triangular Banach algebra.

2. (2n-1)-weak amenability

Forrest and Marcoux in [8] proved the following theorem.

Theorem 2.1. If for every continuous derivation $D: \mathcal{T}$ there exist continuous derivations $\delta_1: \mathcal{A} \longrightarrow \mathcal{A}^{(2n-1)}$, $\delta_4: \mathcal{B} \longrightarrow \mathcal{B}^{(2n-1)}$ and an element $\phi_0 \in \mathcal{X}^{(2n-1)}$ such that for all $\begin{bmatrix} a & x \\ 0 & b \end{bmatrix} \in \mathcal{T}$

$$D(\begin{bmatrix} a & x \\ 0 & b \end{bmatrix}) = \begin{bmatrix} \delta_1(a) - x\phi_0 & \phi_0 a - b\phi_0 \\ 0 & \delta_4(b) + \phi_0 x \end{bmatrix},$$
then $\mathcal{H}^1(\mathcal{T}, \mathcal{T}^{(2n-1)}) \simeq \mathcal{H}^1(\mathcal{A}, \mathcal{A}^{(2n-1)}) \oplus \mathcal{H}^1(\mathcal{B}, \mathcal{B}^{(2n-1)}).$

Proof. See [8, Lemma 3.2, Theorem 3.4 and Theorem 3.7].

It is easy to see that module actions on $\mathcal{T}^{(2n-1)}$ and $\mathcal{T}^{(2n)}$ are as follows:

$$\begin{bmatrix} a & x \\ 0 & b \end{bmatrix} \cdot \begin{bmatrix} F_{2n} & \phi_{2n} \\ 0 & G_{2n} \end{bmatrix} = \begin{bmatrix} aF_{2n} & a\phi_{2n} + xG_{2n} \\ 0 & bG_{2n} \end{bmatrix},$$
$$\begin{bmatrix} F_{2n} & \phi_{2n} \\ 0 & G_{2n} \end{bmatrix} \cdot \begin{bmatrix} a & x \\ 0 & b \end{bmatrix} = \begin{bmatrix} F_{2n}a & F_{2n}x + \phi_{2n}b \\ 0 & G_{2n}b \end{bmatrix},$$

$$\begin{bmatrix} a & x \\ 0 & b \end{bmatrix} \cdot \begin{bmatrix} \theta_{2n-1} & \phi_{2n-1} \\ 0 & \varphi_{2n-1} \end{bmatrix} = \begin{bmatrix} a\theta_{2n-1} + x\phi_{2n-1} & b\phi_{2n-1} \\ 0 & b\varphi_{2n-1} \end{bmatrix},$$

$$\begin{bmatrix} a & x \\ 0 & b \end{bmatrix} \cdot \begin{bmatrix} \theta_{2n-1} & \phi_{2n-1} \\ 0 & \varphi_{2n-1} \end{bmatrix} = \begin{bmatrix} a\theta_{2n-1} + x\phi_{2n-1} & b\phi_{2n-1} \\ 0 & b\varphi_{2n-1} \end{bmatrix},$$
$$\begin{bmatrix} \theta_{2n-1} & \phi_{2n-1} \\ 0 & \varphi_{2n-1} \end{bmatrix} \cdot \begin{bmatrix} a & x \\ 0 & b \end{bmatrix} = \begin{bmatrix} \theta_{2n-1}a & \phi_{2n-1}a \\ 0 & \varphi_{2n-1}b + \phi_{2n-1}x \end{bmatrix}$$

for all
$$\begin{bmatrix} a & x \\ 0 & b \end{bmatrix} \in \mathcal{T}$$
, $\begin{bmatrix} \theta_{2n-1} & \phi_{2n-1} \\ 0 & \varphi_{2n-1} \end{bmatrix} \in \mathcal{T}^{(2n-1)}$ and $\begin{bmatrix} F_{2n} & \phi_{2n} \\ 0 & G_{2n} \end{bmatrix} \in \mathcal{T}^{(2n)}$.

With simple calculations we can prove the following lemmas which are left to reader.

Lemma 2.2. Let $\mathcal{A}(\mathcal{B})$ have a bounded approximate identity and let $T: \mathcal{A} \longrightarrow \mathcal{X}^*(T: \mathcal{B} \longrightarrow \mathcal{X}^*)$ be a bounded right \mathcal{A} -module (left \mathcal{B} module) homomorphism. Then there is $x_0^* \in \mathcal{X}^*$ such that T(a) = $x_0^*a\ (T(b) = bx_0^*)$ for all $a \in \mathcal{A}\ (b \in \mathcal{B})$.

Lemma 2.3. Let n be a positive integer and let $D: \mathcal{T} \longrightarrow \mathcal{T}^{(n)}$ be a derivation. Then

(i)
$$\begin{cases} \delta_{1}: \mathcal{A} \longrightarrow \mathcal{A}^{(n)} \\ \delta_{1}(a) = \pi_{1}(D(\begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix})) \end{cases} \quad and \quad \begin{cases} \delta_{4}: \mathcal{B} \longrightarrow \mathcal{B}^{(n)} \\ \delta_{4}(b) = \pi_{4}(D(\begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix})) \end{cases}$$

$$are \ bounded \ derivations,$$
(ii)
$$\begin{cases} T: \mathcal{A} \longrightarrow \mathcal{X}^{(n)} \\ T(a) = \pi_{2}(D(\begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix})) \end{cases} \quad and \quad \begin{cases} S: \mathcal{B} \longrightarrow \mathcal{X}^{(n)} \\ S(b) = \pi_{2}(D(\begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix})) \end{cases}$$

(ii)
$$\begin{cases} T: \mathcal{A} \longrightarrow \mathcal{X}^{(n)} \\ T(a) = \pi_2(D(\begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix})) \end{cases} \quad and \begin{cases} S: \mathcal{B} \longrightarrow \mathcal{X}^{(n)} \\ S(b) = \pi_2(D(\begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix})) \end{cases}$$

are right (left) A-module and left (right) B-module homomorphisms, respectively.

Proposition 2.5. Let \mathcal{A} have a bounded approximate identity and $\mathcal{A}^{(2n-1)}, \mathcal{B}^{(2n-1)}$ and $\mathcal{X}^{(2n-1)}$ be non-degenerate. Then $\mathcal{H}^1(\mathcal{T}, \mathcal{T}^{(2n-1)})$ $\simeq \mathcal{H}^1(\mathcal{A},\mathcal{A}^{(2n-1)}) \oplus \mathcal{H}^1(\mathcal{B},\mathcal{B}^{(2n-1)}).$

Proof. Let $D: \mathcal{T} \longrightarrow \mathcal{T}^{(2n-1)}$ be a derivation. By Lemmas 2.1, 2.3 and 2.4 there exist derivations $\delta_1: \mathcal{A} \longrightarrow \mathcal{A}^{(2n-1)}$ and $\delta_4: \mathcal{B} \longrightarrow \mathcal{B}^{(2n-1)}$ and $\phi_0 \in \mathcal{X}^{(2n-1)}$ such that

$$D(\begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}) = \begin{bmatrix} \delta_1(a) & \phi_0 a \\ 0 & 0 \end{bmatrix}, \quad \pi_4(D(\begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix})) = \delta_4(b).$$

Let
$$b \in \mathcal{B}$$
 and $D(\begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix}) = \begin{bmatrix} \theta & \phi \\ 0 & \delta_4(b) \end{bmatrix}$, then for all $a \in \mathcal{A}$,

$$\begin{bmatrix} \theta & \phi \\ 0 & \delta_4(b) \end{bmatrix} \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix} \begin{bmatrix} \delta_1(a) & \phi_0 a \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

It follows that $(\phi + b\phi_0)a = 0$ and $\theta a = 0$, $(a \in \mathcal{A})$. Since $\mathcal{X}^{(2n-1)}$ and $\mathcal{A}^{(2n-1)}$ are non-degenerate, we get $\phi = -b\phi_0$, $\theta = 0$ and $D(\begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix}) = \begin{bmatrix} 0 & -b\phi_0 \\ 0 & \delta_4(b) \end{bmatrix}$.

Let $a \in \mathcal{A}, x \in \mathcal{X}, b \in \mathcal{B}$ and $D(\begin{bmatrix} 0 & x \\ 0 & 0 \end{bmatrix}) = \begin{bmatrix} \theta & \phi \\ 0 & \varphi \end{bmatrix}$. We have

$$\begin{bmatrix} \theta & \phi \\ 0 & \varphi \end{bmatrix} \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & x \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \delta_1(a) & \phi_0 a \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Consequently $\phi a = 0$ and $(\theta + x\phi_0)a = 0$. Since $\mathcal{A}^{(2n-1)}$ and $\mathcal{X}^{(2n-1)}$ are non-degenerate, we obtain that $\phi = 0$ and $\theta = -x\phi_0$. A similar calculation shows that $\varphi = \phi_0 x$.

calculation shows that $\varphi = \phi_0 x$.

Therefore $D(\begin{bmatrix} a & x \\ 0 & b \end{bmatrix}) = \begin{bmatrix} \delta_1(a) - x\phi_0 & \phi_0 a - b\phi_0 \\ 0 & \delta_4(b) + \phi_0 x \end{bmatrix}$ for all $\begin{bmatrix} a & x \\ 0 & b \end{bmatrix} \in \mathcal{T}$ and the result follows from Lemma 2.2.

Corollary 2.6. Let \mathcal{A} have a bounded approximate identity, \mathcal{B} be a Banach algebra such that $\mathcal{B}^2 = \mathcal{B}$ and \mathcal{X} be an essential Banach \mathcal{A}, \mathcal{B} -module. Then

$$\mathcal{H}^1(\mathcal{T},\mathcal{T}^*)\simeq\mathcal{H}^1(\mathcal{A},\mathcal{A}^*)\oplus\mathcal{H}^1(\mathcal{B},\mathcal{B}^*).$$

Dales, Ghahramani and Gronbaek [4, proposition, 1.3] have shown that if \mathcal{A} is a weakly amenable Banach algebra, then \mathcal{A}^2 , the linear span of products of elements in \mathcal{A} , is dense in \mathcal{A} (c.f. [10], [11], [12]). Hence \mathcal{A}^* is non-degenerate.

Corollary 2.7. Let A or B have a bounded approximate identity and let X be essential. Then T is weakly amenable if and only if A and B are weakly amenable.

Proof. It is easy to see that \mathcal{A}^* , \mathcal{B}^* and \mathcal{X}^* are non-degenerate. \square

Theorem 2.8. Let A and B have bounded approximate identities. Let n > 0, X and $X^{(2n)}$ be essential Banach A, B-modules. Then

$$\mathcal{H}^1(\mathcal{T},\mathcal{T}^{(2n+1)})\simeq\mathcal{H}^1(\mathcal{A},\mathcal{A}^{(2n+1)})\oplus\mathcal{H}^1(\mathcal{B},\mathcal{B}^{(2n+1)}).$$

Proof. Without loss of generality, we can assume that $\{e_{\alpha}\}$, $\{f_{\alpha}\}$ and $\{\begin{bmatrix} e_{\alpha} & 0 \\ 0 & f_{\alpha} \end{bmatrix}\}$ be bounded approximate identities of \mathcal{A} , \mathcal{B} and \mathcal{T} , respectively. By Lemmas 2.1, 2.3 and 2.4 there exist derivations $\delta_1: \mathcal{A} \longrightarrow \mathcal{A}^{(2n+1)}$ and $\delta_4: \mathcal{B} \longrightarrow \mathcal{B}^{(2n+1)}$, $\phi_0 \in \mathcal{X}^{(2n+1)}$ and $\psi_0 \in \mathcal{X}^{(2n+1)}$ such that

$$D(\begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}) = \begin{bmatrix} \delta_1(a) & \phi_0 a \\ 0 & 0 \end{bmatrix}, D(\begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix}) = \begin{bmatrix} 0 & b\psi_0 \\ 0 & \delta_4(b) \end{bmatrix} (a \in \mathcal{A}, b \in \mathcal{B}).$$

By [3, Proposition, 2.9.7], we have $\psi_0 = -\phi_0$ and therefore

$$D(\begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix}) = \begin{bmatrix} 0 & -b\phi_0 \\ 0 & \delta_4(b) \end{bmatrix}.$$

Let x = ay = zb be an arbitrary element of \mathcal{X} and let

$$D(\begin{bmatrix}0&z\\0&0\end{bmatrix})=\begin{bmatrix}\theta_{2n+1}&\phi_{2n+1}\\0&\psi_{2n+1}\end{bmatrix},\,D(\begin{bmatrix}0&y\\0&0\end{bmatrix})=\begin{bmatrix}\theta_{2n+1}'&\phi_{2n+1}'\\0&\psi_{2n+1}'\end{bmatrix}.$$

Then

$$D(\begin{bmatrix} 0 & x \\ 0 & 0 \end{bmatrix}) = \begin{bmatrix} a\theta'_{2n+1} & 0 \\ 0 & \phi_0 x \end{bmatrix} = \begin{bmatrix} -x\phi_0 & 0 \\ 0 & \psi_{2n+1}b \end{bmatrix} = \begin{bmatrix} -x\phi_0 & 0 \\ 0 & \phi_0 x \end{bmatrix}.$$
 Therefore
$$D(\begin{bmatrix} a & x \\ 0 & b \end{bmatrix}) = \begin{bmatrix} \delta_1(a) - x\phi_0 & \phi_0 a - b\phi_0 \\ 0 & \delta_4(b) + \phi_0 x \end{bmatrix} \text{ and the proof is completed by Lemma 2.2}$$

3. (2n)-weak amenability

In [8] Forrest and Marcoux defined the following sets. For each positive integer n, we denote the centralizer of \mathcal{A} in $\mathcal{A}^{(2n)}$ as

$$\mathcal{Z}_{\mathcal{A}}(\mathcal{A}^{(2n)}) = \{ F_{2n} \in \mathcal{A}^{(2n)} \mid F_{2n}a = aF_{2n} \text{ for all } a \in \mathcal{A} \}.$$

For $F_{2n} \in \mathcal{A}^{(2n)}$ and $G_{2n} \in \mathcal{B}^{(2n)}$ we consider the map $\rho_{F_{2n},G_{2n}} : \mathcal{X} \longrightarrow \mathcal{X}^{(2n)}$ defined by $x \longmapsto F_{2n}x - xG_{2n}$. The set

$$\mathcal{ZR}_{\mathcal{A},\mathcal{B}}(\mathcal{X},\mathcal{X}^{(2n)}) = \{
ho_{F_{2n},G_{2n}}: \mathcal{X} \longrightarrow \mathcal{X}^{(2n)} \mid F_{2n} \in \mathcal{Z}_{\mathcal{A}}(\mathcal{A}^{(2n)}), G_{2n} \in \mathcal{Z}_{\mathcal{B}}(\mathcal{B}^{(2n)})\}$$

is called central Rosenblum operators on \mathcal{X} with coefficient in $\mathcal{X}^{(2n)}$. We also have $\operatorname{Hom}_{\mathcal{A},\mathcal{B}}(\mathcal{X},\mathcal{X}^{(2n)}) = \{\phi: \mathcal{X} \longrightarrow \mathcal{X}^{(2n)} \mid \phi \text{ is left } \mathcal{A}\text{-module and right } \mathcal{B}\text{-module homomorphism } \}.$

Forrest and Marcoux [8] proved the following theorem.

Theorem 3.1. Let n be a positive integer and let A and B be (2n)weakly amenable. Let for every continuous derivation $D: \mathcal{T} \longrightarrow \mathcal{T}^{(2n)}$, there exist derivations $\delta_1: \mathcal{A} \longrightarrow \mathcal{A}^{(2n)}$ and $\delta_4: \mathcal{B} \longrightarrow \mathcal{B}^{(2n)}$, an element $\phi_0 \in \mathcal{X}^{(2n)}$ and a continuous map $\rho: \mathcal{X} \longrightarrow \mathcal{X}^{(2n)}$ such that $\begin{bmatrix} a & x \\ 0 & b \end{bmatrix} =$ $\begin{bmatrix} \delta_1(a) & a\phi_0 - \phi_0 b + \rho(x) \\ 0 & \delta_4(b) \end{bmatrix}, \ \rho(ax) = \delta_1(a)x + a\rho(x) \ and \ \rho(xb) = \rho(x)b + \rho(x$ $\mathcal{H}^1(\mathcal{T}, \mathcal{T}^{(2n)}) \simeq \operatorname{Hom}_{\mathcal{A}, \mathcal{B}}(\mathcal{X}, \mathcal{X}^{(2n)}) / \mathcal{ZR}_{\mathcal{A}, \mathcal{B}}(\mathcal{X}, \mathcal{X}^{(2n)}).$

$$\mathcal{H}^1(\mathcal{T},\mathcal{T}^{(2n)}) \simeq \mathrm{Hom}_{\mathcal{A},\mathcal{B}}(\mathcal{X},\mathcal{X}^{(2n)})/\mathcal{ZR}_{\mathcal{A},\mathcal{B}}(\mathcal{X},\mathcal{X}^{(2n)}).$$

Now we have the following theorem.

Theorem 3.2. Let A or B have a bounded approximate identity, and $\mathcal{A}^{(2n)}$, $\mathcal{B}^{(2n)}$ and $\mathcal{X}^{(2n)}$ be non-degenerate. If \mathcal{A} and \mathcal{B} are (2n)-weakly amenable, then

$$\mathcal{H}^1(\mathcal{T}, \mathcal{T}^{(2n)}) \simeq \operatorname{Hom}_{\mathcal{A}, \mathcal{B}}(\mathcal{X}, \mathcal{X}^{(2n)}) / \mathcal{Z}R_{\mathcal{A}, \mathcal{B}}(\mathcal{X}, \mathcal{X}^{(2n)}).$$

Proof. Without loss of generality we may assume that \mathcal{A} has a bounded approximate identity. Let $D: \mathcal{T} \longrightarrow \mathcal{T}^{(2n)}$ be a derivation. It is easy to see that $\pi_4(D(\begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix})) = 0$ for all $a \in \mathcal{A}$. By Lemmas 2.1, 2.3 and 2.4 there exist derivations $\delta_1 : \mathcal{A} \longrightarrow \mathcal{A}^{(2n)}$ and $\delta_4 : \mathcal{B} \longrightarrow \mathcal{B}^{(2n)}$ and $\phi_0 \in \mathcal{X}^{(2n)}$ such that $D(\begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}) = \begin{bmatrix} \delta_1(a) & a\phi_0 \\ 0 & 0 \end{bmatrix}$ and $\pi_4(D(\begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix})) = \delta_4(b)$

for all $a \in \mathcal{A}, b \in \mathcal{B}$.

For $b \in \mathcal{B}$ let $D(\begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix}) = \begin{bmatrix} F_{2n} & \phi_{2n} \\ 0 & \delta_4(b) \end{bmatrix}$. Then for all $a \in \mathcal{A}$ we have $\begin{bmatrix} \delta_1(a) & a\phi_0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix} + \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} F_{2n} & \phi_{2n} \\ 0 & \delta_4(b) \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$

Since $\mathcal{A}^{(2n)}$ and $\mathcal{X}^{(2n)}$ are non-degenerate, $F_{2n}=0,\,\phi=-\phi_0 b$ and

$$D(\begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix}) = \begin{bmatrix} 0 & -\phi_0 b \\ 0 & \delta_4(b) \end{bmatrix}.$$

For $x \in \mathcal{X}$, let $D(\begin{bmatrix} 0 & x \\ 0 & 0 \end{bmatrix}) = \begin{bmatrix} F_{2n} & \phi_{2n} \\ 0 & G_{2n} \end{bmatrix}$. Then for each a in \mathcal{A} , b in \mathcal{B}

$$\begin{bmatrix} 0 & \phi_0 b \\ 0 & \delta_4(b) \end{bmatrix} \begin{bmatrix} 0 & x \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & b \end{bmatrix} \begin{bmatrix} F_{2n} & \phi_{2n} \\ 0 & G_{2n} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix},$$

$$\begin{bmatrix} F_{2n} & \phi_{2n} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & x \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \delta_1(a) & a\phi_0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Therefore $F_{2n} = G_{2n} = 0$. We define $\rho : \mathcal{X} \longrightarrow \mathcal{X}^{(2n)}$ by $x \longmapsto \pi_2(D(\begin{bmatrix} 0 & x \\ 0 & 0 \end{bmatrix}))$. A simple calculation shows that $\rho(ax) = \delta_1(a)x + a\rho(x)$ and $\rho(xb) = \rho(x)b + x\delta_4(b)$ for all $a \in \mathcal{A}, b \in \mathcal{B}$. By Theorem 3.1 the proof is completed.

Let $\mathcal A$ be a Banach algebra. We consider the triangular Banach algebra

$$\mathcal{T} = \mathcal{T}_2 \otimes \mathcal{A} = egin{bmatrix} \mathcal{A} & \mathcal{A} \ \mathcal{A} \end{bmatrix},$$

where \mathcal{T}_2 denotes the algebra of 2×2 upper triangular matrices.

Proposition 3.3. Let \mathcal{A} be a Banach algebra with a bounded approximate identity. If \mathcal{A} is (2n)-weakly amenable and $\mathcal{A}^{(2n)}$ is non-degenerate, then $\mathcal{T} = \mathcal{T}_2 \otimes \mathcal{A}$ is (2n)-weakly amenable.

Proof. By Theorem 3.2 it is sufficient to show that $\operatorname{Hom}_{\mathcal{A},\mathcal{A}}(\mathcal{A},\mathcal{A}^{(2n)})$ $\simeq Z_{\mathcal{A},\mathcal{A}}(\mathcal{A},\mathcal{A}^{(2n)})$. Let (e_{α}) be a bounded approximate identity of \mathcal{A} and let $\phi: \mathcal{A} \longrightarrow \mathcal{A}^{(2n)}$ be a \mathcal{A} -module homomorphism. There exist $E \in \mathcal{A}^{(2n)}$ and a subnet $\{\phi(e_{\beta})\}$ of $\{\phi(e_{\alpha})\}$ such that $\phi(e_{\beta}) \longrightarrow E$ in the weak* topology. A simple calculation shows that for every a in \mathcal{A} ; $\phi(a) = aE = Ea$. Therefore $E \in Z_{\mathcal{A}}(\mathcal{A}^{(2n)})$ and $\phi = \rho_{E,0} \in Z_{\mathcal{A}}, \mathcal{A}(\mathcal{A}, \mathcal{A}^{(2n)})$.

Theorem 3.4. Let \mathcal{A} and \mathcal{B} have bounded approximate identities. Let n be a positive integer, \mathcal{X} and $\mathcal{X}^{(2n-1)}$ be essential Banach modules. If \mathcal{A} and \mathcal{B} are (2n)-weakly amenable, then

$$\mathcal{H}^1(\mathcal{T},\mathcal{T}^{(2n)}) \simeq \mathrm{Hom}_{\mathcal{A},\mathcal{B}}(\mathcal{X},\mathcal{X}^{(2n)})/\mathcal{ZR}_{\mathcal{A},\mathcal{B}}(\mathcal{X},\mathcal{X}^{(2n)}).$$

Lemma 3.5. Suppose that \mathcal{T} is 2-weakly amenable. Then there exist $F_0 \in Z_{\mathcal{A}}(\mathcal{A}^{**})$ and $G_0 \in Z_{\mathcal{B}}(\mathcal{B}^{**})$ such that for every x in \mathcal{X} ; $\hat{x} = xG_0 - F_0x$ where \hat{x} is the canonical image of x in \mathcal{X}^{**} .

Proof. It is easy to see that $D: \mathcal{T} \longrightarrow \mathcal{T}^{**}$ defind by $\begin{bmatrix} a & x \\ 0 & b \end{bmatrix} \longrightarrow \begin{bmatrix} 0 & \hat{x} \\ 0 & 0 \end{bmatrix}$ is a continuous derivation. Therefore there are $F_0 \in \mathcal{A}^{**}, G_0 \in \mathcal{B}^{**}$ and $x_0^{**} \in \mathcal{A}^{**}$ such that $D = \delta_{\begin{bmatrix} F_0 & x_0^{**} \\ 0 & G_0 \end{bmatrix}}$. So that for every $a \in \mathcal{A}, b \in \mathcal{B}$ and $x \in \mathcal{X}$ we have

$$\begin{bmatrix} 0 & \hat{x} \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} a & x \\ 0 & b \end{bmatrix} \begin{bmatrix} F_0 & x_0^{**} \\ 0 & G_0 \end{bmatrix} - \begin{bmatrix} F_0 & x_0^{**} \\ 0 & G_0 \end{bmatrix} \begin{bmatrix} a & x \\ 0 & b \end{bmatrix}$$
$$= \begin{bmatrix} aF_0 - F_0 a & ax_0^{**} + xG_0 - F_0 x - x_0^{**} b \\ 0 & bG_0 - G_0 b \end{bmatrix}.$$

Hence $F_0 \in Z_{\mathcal{A}}(\mathcal{A}^{**})$, $G_0 \in Z_{\mathcal{B}}(\mathcal{B}^{**})$ and for every x in \mathcal{X} , $\hat{x} = xG_0 - F_0x$.

Proposition 3.6. Let $\mathcal{T} = \begin{bmatrix} \mathcal{A} & \mathcal{A}^{(m)} \\ \mathcal{A} \end{bmatrix}$ for some nonnegative integer m. Suppose that \mathcal{T} is (2n)-weakly amenable for some positive integer n. Then \mathcal{A} has a bounded approximate identity.

Proof. \mathcal{T} is 2-weakly amenable by [4, Proposition, 1.2]. So there exist $F_0, G_0 \in Z_{\mathcal{A}}(\mathcal{A}^{**})$ such that $\hat{x} = xG_0 - F_0x$ $(x \in \mathcal{A}^{(m)})$ by Lemma 3.5. If m is odd then for all $a^* \in \mathcal{A}^*$; $a^* = a^*(G_0 - F_0) = (G_0 - F_0)a^*$, and if m is even then for all $a \in \mathcal{A}$; $\hat{a} = a(F_0 - G_0) = (F_0 - G_0)a$. So in both cases it is easy to see that $G_0 - F_0$ is a mixed unit for \mathcal{A}^{**} and hence \mathcal{A} has a bounded approximate identity.

It is well known that for a Banach algebra \mathcal{A} its second dual \mathcal{A}^{**} is a Banach algebra when equipped with the first or second Arens products (for more details see [6]). Recall that a Banach algebra \mathcal{A} is called a dual Banach algebra if there is a closed submodule \mathcal{X} of \mathcal{A}^* such that $\mathcal{A} = \mathcal{X}^*$ (see [19, 4.4.1]).

Proposition 3.7. Let \mathcal{A} be a second dual of a Banach algebra or a dual Banach algebra and $\mathcal{T} = \begin{bmatrix} \mathcal{A} & \mathcal{A}^{(2m)} \\ \mathcal{A} \end{bmatrix}$ for some positive integer m. Suppose that \mathcal{T} is (2n)-weakly amenable for some positive integer n. Then \mathcal{A} has an identity.

Proof. Without loss of generality, we may assume that \mathcal{A} is a dual Banach algebra or the second dual of a Banach algebra with the first Arens product. \mathcal{T} is 2-weakly amenable by [4, Proposition, 1.2]. So by Lemma 3.5 there exist $F_0, G_0 \in Z_{\mathcal{A}}(\mathcal{A}^{(**)})$ such that for all $x \in \mathcal{A}^{(2m)}$; $\hat{x} = xG_0 - F_0x$. Therefore $\hat{a} = a(G_0 - F_0) = (G_0 - F_0)a$ for all a in \mathcal{A} . Suppose that $\pi: \mathcal{X} \longrightarrow \mathcal{X}^{(**)}$ is the canonical embedding, where \mathcal{X} is the predual of \mathcal{A} . Put $e = \pi^*(G_0 - F_0)$. For $a \in \mathcal{A}$ we have

$$\langle x, ea \rangle = \langle \pi(ax), G_0 - F_0 \rangle$$

$$= \langle a\pi(x), G_0 - F_0 \rangle$$

$$= \langle \pi(x), \hat{a} \rangle$$

$$= \langle x, a \rangle \ (x \in \mathcal{X}).$$

So e is a right identity for \mathcal{A} . Now if \mathcal{A} is a dual Banach algebra, similarly e is a left identity for \mathcal{A} , and if \mathcal{A} is the second dual of Banach algebra \mathcal{B} , then for $a \in \mathcal{A}$ there exists net $\{b_{\alpha}\}$ in \mathcal{B} such that $b_{\alpha} \longrightarrow a$ in the weak* topology.

Therefore e is a right identity for \mathcal{A} and so \mathcal{A} has an identity.

Corolollary 3.8. Let \mathcal{A} be the second dual of a Banach algebra or a dual Banach algebra and let $\mathcal{T} = \begin{bmatrix} \mathcal{A} & \mathcal{A} \\ \mathcal{A} \end{bmatrix}$. Suppose that \mathcal{T} is (2n)-weakly amenable for some positive integer n. Then \mathcal{T} is (2n)-weakly amenable for all positive integer n.

4. Amenability of the triangular Banach algebra \mathcal{T}

In this section we give a necessary and sufficient condition for the amenability of \mathcal{T} .

Theorem 4.1. If $\mathcal{T} = \begin{bmatrix} \mathcal{A} & \mathcal{X} \\ 0 & \mathcal{B} \end{bmatrix}$ has a bounded approximate identity, then \mathcal{A} and \mathcal{B} have bounded approximate identities and \mathcal{X} is neo-unital.

Proof. Let $\left\{ \begin{bmatrix} a_{\alpha} & x_{\alpha} \\ 0 & b_{\alpha} \end{bmatrix} \right\}$ be a bounded approximate identity for \mathcal{T} . For any $a \in \mathcal{A}$, we have

$$\left[\begin{array}{cc} a_{\alpha} & x_{\alpha} \\ 0 & b_{\alpha} \end{array}\right] \left[\begin{array}{cc} a & 0 \\ 0 & 0 \end{array}\right] = \left[\begin{array}{cc} a_{\alpha}a & 0 \\ 0 & 0 \end{array}\right] \longrightarrow \left[\begin{array}{cc} a & 0 \\ 0 & 0 \end{array}\right],$$

and hence $a_{\alpha}a \longrightarrow a$. Similarly $aa_{\alpha} \longrightarrow a$ and thus $\{a_{\alpha}\}$ is a bounded approximate identity for \mathcal{A} . Similarly $\{b_{\alpha}\}$ is a bounded approximate identity for \mathcal{B} . For any $x \in X$,

$$\left[\begin{array}{cc} a_{\alpha} & x_{\alpha} \\ 0 & b_{\alpha} \end{array}\right] \left[\begin{array}{cc} 0 & x \\ 0 & 0 \end{array}\right] = \left[\begin{array}{cc} 0 & a_{\alpha}x \\ 0 & 0 \end{array}\right] \longrightarrow \left[\begin{array}{cc} 0 & x \\ 0 & 0 \end{array}\right],$$

so that $a_{\alpha}x \longrightarrow x$ and thus by Cohn factorization theorem $\mathcal{X} = \mathcal{A}.\mathcal{X}$ and similarly $\mathcal{X} = \mathcal{X}.B$.

Now we prove the main theorem of this section.

Theorem 4.2. \mathcal{T} is amenable if and only if both \mathcal{A}, \mathcal{B} are amenable and $\mathcal{X} = 0$.

Proof. Let \mathcal{A} and \mathcal{B} be amenable and $\mathcal{X} = 0$. Since $\begin{bmatrix} \mathcal{A} & 0 \\ 0 & 0 \end{bmatrix} \simeq \mathcal{A}$, $\begin{bmatrix} \mathcal{A} & 0 \\ 0 & \mathcal{B} \end{bmatrix} / \begin{bmatrix} \mathcal{A} & 0 \\ 0 & 0 \end{bmatrix} \simeq \mathcal{B}$, the closed ideal $\begin{bmatrix} \mathcal{A} & 0 \\ 0 & 0 \end{bmatrix}$ of \mathcal{T} and the quotient algebra $\begin{bmatrix} \mathcal{A} & 0 \\ 0 & \mathcal{B} \end{bmatrix} / \begin{bmatrix} \mathcal{A} & 0 \\ 0 & 0 \end{bmatrix}$ are amenable and thus $\mathcal{T} = \begin{bmatrix} \mathcal{A} & 0 \\ 0 & \mathcal{B} \end{bmatrix}$ is amenable. For the converse, suppose that \mathcal{T} is amenable. Since $\begin{bmatrix} \mathcal{A} & \mathcal{X} \\ 0 & 0 \end{bmatrix}$ is a closed ideal of \mathcal{T} , the quotient algebra

 $\begin{bmatrix} \mathcal{A} & \mathcal{X} \\ 0 & \mathcal{B} \end{bmatrix} / \begin{bmatrix} \mathcal{A} & \mathcal{X} \\ 0 & 0 \end{bmatrix} \text{ is amenable. On the other hand } \begin{bmatrix} \mathcal{A} & \mathcal{X} \\ 0 & \mathcal{B} \end{bmatrix} \\ / \begin{bmatrix} \mathcal{A} & \mathcal{X} \\ 0 & 0 \end{bmatrix} \simeq B, \text{ thus } \mathcal{B} \text{ is amenable. Similarly, since } \begin{bmatrix} 0 & \mathcal{X} \\ 0 & \mathcal{B} \end{bmatrix} \text{ is a closed ideal of } \mathcal{T} \text{ and } \begin{bmatrix} \mathcal{A} & \mathcal{X} \\ 0 & \mathcal{B} \end{bmatrix} / \begin{bmatrix} 0 & \mathcal{X} \\ 0 & \mathcal{B} \end{bmatrix} \simeq \mathcal{A}, \text{ the Banach algebra } \mathcal{A} \text{ is amenable. Since } \mathcal{T} \text{ is amenable and } \begin{bmatrix} 0 & \mathcal{X} \\ 0 & 0 \end{bmatrix} \text{ is a closed ideal of } \mathcal{T} \text{ which is complemented in } \mathcal{T}, \begin{bmatrix} 0 & \mathcal{X} \\ 0 & 0 \end{bmatrix} \text{ is amenable and thus it has a bounded approximate identity. However this is not possible unless } \mathcal{X} = 0.$

${\bf Acknowledgment}$

A part of this paper was written while the third author visited the University of Alberta. He is pleased to thank the department there for their warm hospitality, and he wishes to express his gratitude to Professor Antony To-Ming Lau. We would also like to thank the referee for careful reading of the paper and giving many useful suggestions.

REFERENCES

- [1] W. G. Bade, P. G. Curtis and H. G. Dales, Amenability and weak amenability for Beurling and Lipschits algebras, *Proc. London Math. Soc.* **55** (3) (1987), 359-377.
- [2] F. F. Bonsall and J. Duncan, *Complete normed algebras*, Springer-Verlage, Berlin, 1973.
- [3] H.E. Dales, Banach Algebra and Automatic Continuity, London Mathematical Society Monographs, Vol. 24, Clarendon Press, Oxford, 2000.
- [4] H.G.Dales, F.Ghahramani and N.Gronbaek, Derivations into iterated duals of Banach algebras, Studia Math. 128 (1998), 19-54.
- [5] M. Despic and F. Ghahramani, Weak amenability of group algebras of locally compact groups, Canad. Math. Bull. 37 (1994), 165-167.
- [6] J. Duncan and S.A.R. Hosseiniun, The second dual of a Banach algebra, Proc. Roy. Soc. Edinburgh 84 A (1979), 309-325.
- [7] B.E. Forrest, L.W. Marcoux, Derivations of Triangular Banach Algebras, *Indiana Univ. Math. J.* 45 (1996), 441-462.
- [8] B.E. Forrest, L.W. Marcoux, Weak Amenability of Triangular Banach Algebras, *Trans. Amer. Math. Soc.* **345** (2002), 1435-1452.
- [9] F. Ghahramani, R. J. Loy and G. A. Willis, Amenability and weak amenability of second conjugate of Banach algebras, *Proc. Amer. Math. Soc.* 124 (5) (1996), 1489-1497.

- [10] N. Grønbæk, A characterization of weakly amenable Banach algebras, Studia Math. 94 (1989), 150-162.
- [11] N. Grønbæk, Weak amenability of group algebras, Bull. London Math. Soc. 23 (1991), 231-284.
- [12] N. Grønbæk, Weak and cyclic amenability for non-commutative Banach algebras, Proc. Edinburg Math. Soc. 35 (1992), 315-328.
- [13] U. Haagerup, All nuclear C^* -algebras are amenable, *Invent. Math.* **74** (1983), 305-319.
- [14] A. Ya. Helmeskii, The homology of Banach and topological algebras. Kluwer, Academic Press, Dordrecht, 1989.
- [15] B. E. Johnson, Cohomology in Banach algebras, Mem. Amer. Math. Soc. 127 (1972).
- [16] B. E. Johnson, Weak amenability of group algebras, Bull. Lodon Math. Soc. 23 (1991), 281-284.
- [17] B. E. Johnson, Derivation from $L^1(G)$ into $L^1(G)$ and $L^{\infty}(G)$, Harmonic analysis, Luxembourg, (1987), 191-198. Lecture Notes in Math., 1359, Springer-Verlage, Berlin, 1988.
- [18] T. W. Palmer, Banach Algebra and The General Theory of *-Algebras Vol.1, Cambridge University Press, Cambridge, 1994.
- [19] V.Runde, Lectures on amenability, Springer-Verlage, Berlin, Hedinberg, New York, 2001.

A.R. Medghalchi and M. H. Sattari

Department of Mathematics Teacher Training University 599, Taleghani Avenue Tehran 15614, IRAN

e-mail:a_medghalchi@saba.tmu.ac.ir e-mail:sattari2005@yahoo.com

T. Yazdanpanah

Department of Mathematics Faculty of Sciences Persian Gulf University 75168, Boushehr, IRAN e-mail:yazdanpanah@pguac.ir