

SEQUENTIALLY COHEN-MACAULAY GRAPHS OF FORM θ_{n_1, \dots, n_k}

F. MOHAMMADI* AND D. KIANI

Communicated by Siamak Yassemi

ABSTRACT. Let k be an integer greater than 2 and n_1, \dots, n_k be a sequence of positive integers with at most one of them being equal to 1. Let θ_{n_1, \dots, n_k} be a graph consisting of k paths, having only their endpoints in common. We characterize all sequentially Cohen-Macaulay graphs of this type. We also show for these types of graphs the notions of vertex decomposable, shellable and sequentially Cohen-Macaulay are equivalent.

1. Introduction

Let G be a finite simple graph. To G with vertex set $[n] = \{1, \dots, n\}$ and edge set $E(G)$, one can associate an ideal $\mathcal{I}(G) \subset R = K[x_1, \dots, x_n]$, called the edge ideal of G , which is generated by all monomials $x_i x_j$ such that $\{i, j\} \in E(G)$. Here, K is an arbitrary field. The independence complex Δ_G of a graph G is defined by

$$\Delta_G = \{A \subseteq V \mid A \text{ is an independent set in } G\},$$

where, A is an independent set in G if none of its elements are adjacent. Note that Δ_G is precisely the simplicial complex associated with $\mathcal{I}(G)$.

It is a well-known consequence of Menger's Theorem [5, Theorem 3.3.5] that each 3-connected graph has an induced subgraph of the form

MSC(2010): Primary: 05C75; Secondary: 13H10.

Keywords: Sequentially Cohen-Macaulay graph, shellable graph.

Received: 20 June 2009, Accepted: 23 August 2009.

*Corresponding author

© 2010 Iranian Mathematical Society.

$\theta_{p,q,r}$, for some natural numbers p, q and r . This was our motivation to study sequentially Cohen-Macaulay graphs of the form θ_{n_1, \dots, n_k} .

A graded R -module M is called *sequentially Cohen-Macaulay* (over K) if there exists a finite filtration of graded R -modules,

$$0 = M_0 \subset M_1 \subset \cdots \subset M_r = M,$$

such that each M_i/M_{i-1} is Cohen-Macaulay, and the Krull dimensions of the quotients are increasing; that is,

$$\dim(M_1/M_0) < \dim(M_2/M_1) < \cdots < \dim(M_r/M_{r-1}).$$

A graph G is said to be sequentially Cohen-Macaulay, if $R/\mathcal{I}(G)$ is a sequentially Cohen-Macaulay R -module.

On the other hand, a simplicial complex Δ is called *shellable*, in the sense of Björner and Wachs [1], if the facets (maximal faces) of Δ can be ordered as F_1, \dots, F_s such that for all $1 \leq i < j \leq s$, there exists some $v \in F_j \setminus F_i$ and some $l \in \{1, \dots, j-1\}$ with $F_j \setminus F_l = \{v\}$. A graph G is called shellable, if Δ_G is a shellable simplicial complex. In [12], Stanley showed that every shellable simplicial complex was sequentially Cohen-Macaulay, but the converse was not true.

Studying shellable or sequentially Cohen-Macaulay graphs has attracted significant attentions of researchers working in the borderline of combinatorial commutative algebra and algebraic combinatorics; see [1, 6, 7, 8, 10, 14, 16]. In [8], Francisco and Van Tuyl characterized all sequentially Cohen-Macaulay cycles. They showed that the n -cycle C_n was sequentially Cohen-Macaulay if and only if $n \in \{3, 5\}$ (see [8, Proposition 4.1]). In [6], Faridi showed that simplicial trees were sequentially Cohen-Macaulay. Moreover, in [10], sequentially Cohen-Macaulay cacti graphs (a cactus is a connected graph in which each edge belongs to at most one cycle) were characterized. In addition, in [14], Van Tuyl and Villarreal showed that a bipartite graph G was shellable if and only if it was sequentially Cohen-Macaulay (see [14, Theorem 3.8]).

Here, we determine all sequentially Cohen-Macaulay graphs of the form θ_{n_1, \dots, n_k} , where $\{n_1, \dots, n_k\} \neq \{2, 5\}$. For $\{n_1, \dots, n_k\} \neq \{2, 5\}$, we show in Theorem 2.6 that θ_{n_1, \dots, n_k} is sequentially Cohen-Macaulay if and only if $\{1, 2\} \subseteq \{n_1, \dots, n_k\}$ or $\{2, 3\} \subseteq \{n_1, \dots, n_k\}$ or $\{n_1, \dots, n_k\} = \{1, 4\}$. Moreover, as a result of this theorem, in Theorem 2.7 we show those graphs of the form θ_{n_1, \dots, n_k} , which satisfy each one of the latter relations, are sequentially Cohen-Macaulay if and only if they are shellable or vertex decomposable.

Finally, in Proposition 2.8, we show that for $\{n_1, \dots, n_k\} = \{2, 5\}$, the graph θ_{n_1, \dots, n_k} is not vertex decomposable. Therefore, we characterize all vertex decomposable graphs of the form θ_{n_1, \dots, n_k} in Theorem 2.9. In Proposition 2.10, by direct computation, we show that for $k = 3$ and $\{n_1, \dots, n_k\} = \{2, 5\}$, the graph θ_{n_1, \dots, n_k} is not even sequentially Cohen-Macaulay. This result and computational evidences from some other examples lead us to conjecture that all graphs of the form θ_{n_1, \dots, n_k} , for which $\{n_1, \dots, n_k\} = \{2, 5\}$, are not sequentially Cohen-Macaulay.

Characterizing vertex decomposable, shellable and sequentially Cohen-Macaulay graphs of the form θ_{n_1, \dots, n_k} with [13, Lemma 2.4] and [14, Theorem 2.9] enable us to get more examples of vertex decomposable, shellable and sequentially Cohen-Macaulay graphs.

2. Sequentially Cohen-Macaulay graphs of the form θ_{n_1, \dots, n_k}

Let k be an integer greater than 1 and n_1, \dots, n_k be a sequence of positive integers. Let θ_{n_1, \dots, n_k} be the graph constructed by k paths of length n_1, \dots, n_k , with only their endpoints being in common. By length of a path, we mean the number of edges in the path. Since the graphs are assumed simple, at most one of the n_i s in θ_{n_1, \dots, n_k} can be equal to one. If $k = 2$, then θ_{n_1, \dots, n_k} would be a cycle of length $n_1 + n_2$. The vertex decomposable and sequentially Cohen-Macaulay graphs of these types are completely studied in [8, 16]. Here, we assume $k > 2$ and characterize all vertex decomposable, shellable and sequentially Cohen-Macaulay graphs of the form θ_{n_1, \dots, n_k} .

Given a simplicial complex Δ on $[n]$, the *Alexander dual* complex Δ^\vee is defined by $\Delta^\vee = \{[n] \setminus F \mid F \notin \Delta\}$. Unless otherwise stated, when we discuss the Alexander dual Δ^\vee of a simplicial complex Δ , we assume that $[n] \setminus i \notin \Delta$, for all $i \in [n]$. Thus, Δ^\vee is again a simplicial complex on $[n]$.

Let $I = (x_{1,1} \cdots x_{1,s_1}, \dots, x_{t,1} \cdots x_{t,s_t})$ be a square-free monomial ideal. The ideal

$$I^\vee = (x_{1,1}, \dots, x_{1,s_1}) \cap \dots \cap (x_{t,1}, \dots, x_{t,s_t})$$

is called the Alexander dual of I . These two ideals are related in the following way. If I is the Stanley-Reisner ideal of a simplicial complex Δ , then the Stanley-Reisner ideal of its Alexander dual Δ^\vee is I^\vee .

Another related notion is componentwise linear ideals, introduced by Herzog and Hibi, to characterize sequentially Cohen-Macaulay ideals.

Let I be a graded ideal of R and let $I_{<d>}$ be the ideal generated by all homogeneous polynomials of degree d of I . A graded ideal I of R is called *componentwise linear* if $I_{<d>}$ has a linear resolution, for every d . Let I be a square-free monomial ideal in a polynomial ring. The ideal generated by the square-free monomials of degree d of I is denoted by $I_{[d]}$. Herzog and Hibi in [9, Proposition 1.5] showed that the square-free ideal I was componentwise linear if and only if $I_{[d]}$ had a linear resolution for every d .

Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. A subset $C \subseteq V(G)$ is a *minimal vertex cover* of G if: (1) every edge of G is incident with one vertex in C , and (2) there is no proper subset of C with the first property. In [8], Francisco and Van Tuyl showed that if $\mathcal{I}(G)$ was the ideal of a graph G , then

$$\mathcal{I}(G)_{[d]}^\vee = (\{x_{i_1} \cdots x_{i_d} \mid \{x_{i_1}, \dots, x_{i_d}\} \text{ is a vertex cover of } G \text{ of size } d\}).$$

In [9], Herzog and Hibi showed the following theorem to be used in the proof of Proposition 2.4.

Theorem A. Let I be a square-free monomial ideal in a polynomial ring. Then I^\vee is componentwise linear if and only if R/I is sequentially Cohen-Macaulay.

Let $N(v)$ be the set of all adjacent vertices of v and let $N[v] = N(v) \cup \{v\}$. Vertex decomposability was introduced by Provan and Billera [11] in the pure case, and extended to the non-pure case by Björner and Wachs [2]. We will use the following definition of vertex decomposable graphs which is an interpretation of the definition of vertex decomposable for the independence complex of a graph, as stated in [13, 16].

Definition 2.1. The independence complex of G is vertex decomposable if G is a totally disconnected graph (with no edges), or if

- $G \setminus v$ and $G \setminus N[v]$ are both vertex decomposable, and
- No independent set in $G \setminus N[v]$ is a maximal independent set in $G \setminus v$.

A vertex v which satisfies in these conditions is called a shedding vertex.

The graph G is called vertex decomposable if its independence complex is vertex decomposable. It is known that the any vertex decomposable graph is shellable and so is sequentially Cohen-Macaulay (see [16]).

For characterizing vertex decomposable, shellable and sequentially Cohen-Macaulay graphs of the form θ_{n_1, \dots, n_k} , we have to distinguish among some cases, depending on n_1, \dots, n_k , as follows.

Proposition 2.2. *If $\{1, 2\} \subseteq \{n_1, \dots, n_k\}$, then θ_{n_1, \dots, n_k} is vertex decomposable and so is shellable and sequentially Cohen-Macaulay.*

Proof. Two paths of length one and two form a triangle. Let v, u and w be its vertices such that $\deg(v) = 2$. The graphs $\theta_{n_1, \dots, n_k} \setminus \{u\}$ and $\theta_{n_1, \dots, n_k} \setminus N[u]$ are chordal and so they are vertex decomposable, by [16, Theorem 1]. For any independent set F in $\theta_{n_1, \dots, n_k} \setminus N[u]$, $F \cup \{v\}$ is an independent set in $\theta_{n_1, \dots, n_k} \setminus \{u\}$. Therefore, θ_{n_1, \dots, n_k} fulfills the conditions of Definition 2.1, which completes the proof. \square

Remark 2.3. If in the above proposition, one assumes $\{n_1, \dots, n_k\} = \{1, 2\}$, then the associated graph, θ_{n_1, \dots, n_k} , is chordal. These types of graphs are known to be vertex decomposable, by [16, Theorem 1].

A chordless path in a graph G is a path v_1, v_2, \dots, v_k in G with no edge $v_i v_j$ with $j \neq i + 1$. A simplicial k -path in G is a chordless path v_1, v_2, \dots, v_k which cannot be extended on both endpoints to a chordless path $v_0, v_1, \dots, v_k, v_{k+1}$ in G .

Proposition 2.4. *Let $\{2, 3\} \subseteq \{n_1, \dots, n_k\}$. Then, θ_{n_1, \dots, n_k} is vertex decomposable and consequently shellable and sequentially Cohen-Macaulay.*

Proof. Let $P_1 : u, x, v$ and $P_2 : u, y, z, v$ be two paths of length two and three in θ_{n_1, \dots, n_k} . Since the path $P : x, u, y$ is a simplicial 3-path, which is not a subgraph of any chordless C_4 , by [16, Lemma 4.3] we deduce that G is vertex decomposable. \square

Proposition 2.5. *Let $\{n_1, \dots, n_k\} = \{1, 4\}$. Then, θ_{n_1, \dots, n_k} is vertex decomposable and consequently shellable and sequentially Cohen-Macaulay.*

Proof. Each cycle other than C_5 in θ_{n_1, \dots, n_k} has a chord and so, by [16, Theorem 1], it is vertex decomposable. \square

The following theorem is one of the main results of this paper which characterizes all sequentially Cohen-Macaulay graphs of the form θ_{n_1, \dots, n_k} , where $\{n_1, \dots, n_k\} \neq \{2, 5\}$.

Theorem 2.6. *Let $n_1, \dots, n_k \neq \{2, 5\}$. Then, θ_{n_1, \dots, n_k} is sequentially Cohen-Macaulay if and only if one of the following holds:*

- (1) $\{1, 2\} \subseteq \{n_1, \dots, n_k\}$.
- (2) $\{2, 3\} \subseteq \{n_1, \dots, n_k\}$.
- (3) $\{1, 4\} \subseteq \{n_1, \dots, n_k\}$.

Proof. “If”. Suppose that one of (1) to (3) holds. Then, by Proposition 2.2, Proposition 2.4 and Proposition 2.5, the result holds.

“Only if”. Let $G = \theta_{n_1, \dots, n_k}$ be a sequentially Cohen-Macaulay graph. The proof is by induction on k . If $k = 2$, then the graph is a cycle and so the result holds by [8, Proposition 4.1]. Let $k > 2$, $n_1 \leq \dots \leq n_k$ and $P_i : x, x_{i,1}, \dots, x_{i,n_i-1}, y$, for $1 \leq i \leq k$, be the paths which construct G . If $n_t \geq 6$, for some $t \geq 3$, then

$$H = G \setminus \bigcup_{i \neq t}^k (N[x_{i,2}] \cup N[x_{i,n_i-2}])$$

has a component of the form $\theta_{n_1, \dots, n_{t-1}}$. So, by the induction hypothesis, (1) or (2) or (3) holds, for $\theta_{n_1, \dots, n_{t-1}}$. If (1) or (2) holds for $\theta_{n_1, \dots, n_{t-1}}$, then this holds, for θ_{n_1, \dots, n_k} . Let (3) holds for $\theta_{n_1, \dots, n_{t-1}}$, but $\{n_1, \dots, n_k\} \neq \{1, 4\}$. Let $S = \{j; n_j = 4\}$ and $H' = G \setminus \bigcup_{j \in S} N[x_{j,2}]$. Since $n_2 = 4$, then H' has no path of length two, three and four. By the induction hypothesis, H' is not sequentially Cohen-Macaulay, which is a contradiction by [14, Theorem 3.3].

So, we can assume that $n_k < 6$. Since G has no vertex of degree one, it is not a bipartite graph by [14, Lemma 2.8]. Therefore, for $n_k = 2$, we have $n_1 = 1$ and so (1) holds. Similarly, If $n_k = 3$, then $n_i = 2$, for some i , and so (2) holds. If $n_k = 4$, then $G \setminus N[x_{k,2}]$ is $\theta_{n_1, \dots, n_{k-1}}$. If (1), (2) or (3) holds, for $\theta_{n_1, \dots, n_{k-1}}$, then the similar statement holds for G . So, assume that $n_k = 5$. Since G is not bipartite, for some i we have $n_i = 2$ or 4. If $n_i = 4$ for some i , then $H = G \setminus N[x_{i,2}]$ is sequentially Cohen-Macaulay and so (1) or (2) holds, which completes the result.

Otherwise, the assumption $\{n_1, \dots, n_k\} \neq \{2, 5\}$ shows that $n_j = 1$ or 3, for some j , and so (1) or (2) holds. \square

Recently, Van Tuyl showed that in bipartite graphs, the three concepts vertex decomposability, shellability and sequentially Cohen-Macaulayness are equivalent; see [13, Theorem 2.10]. Using the proof of the above theorem, we have the same property for θ_{n_1, \dots, n_k} , where $\{n_1, \dots, n_k\} \neq \{2, 5\}$.

Theorem 2.7. *Let $n_1, \dots, n_k \neq \{2, 5\}$. Then, the followings are equivalent:*

- (i) θ_{n_1, \dots, n_k} is sequentially Cohen-Macaulay.
- (ii) θ_{n_1, \dots, n_k} is shellable.
- (iii) θ_{n_1, \dots, n_k} is vertex decomposable.

Proof. Note that (iii) \Rightarrow (ii) \Rightarrow (i) always holds for any graph. It is enough to show that for these type of graphs, (i) \Rightarrow (iii). Let θ_{n_1, \dots, n_k} be a sequentially Cohen-Macaulay graph. Then, Theorem 2.6 shows that θ_{n_1, \dots, n_k} satisfies one of the relations of Theorem 2.6. Therefore, by Proposition 2.2, Proposition 2.4 and Proposition 2.5, we deduce that θ_{n_1, \dots, n_k} is vertex decomposable. \square

In the following, we consider the case $\{n_1, \dots, n_k\} = \{2, 5\}$.

Proposition 2.8. *Let $\{n_1, \dots, n_k\} = \{2, 5\}$. Then, θ_{n_1, \dots, n_k} is not vertex decomposable.*

Proof. Let P_1, \dots, P_s be the paths of length two in $G = \theta_{n_1, \dots, n_k}$ and P_{s+1}, \dots, P_k be the paths of length five in G . Consider the labeling for G such that $P_j : u, x_j, v$, for $1 \leq j \leq s$, and $P_j : u, x_{j,1}, x_{j,2}, x_{j,3}, x_{j,4}, v$, for $s+1 \leq j \leq k$. We claim that no vertex of G is a shedding vertex to deduce that G is not vertex decomposable. For any $s+1 \leq j \leq k$, the independent set $\{u, x_{s+1,4}, \dots, x_{k,4}\}$ is maximal in both graphs $G \setminus x_{j,2}$ and $G \setminus N[x_{j,2}]$. For the other vertices of G , the similar arguments hold. Therefore, G is not vertex decomposable. \square

Proposition 2.8 and Theorem 2.6 imply the following characterization of the vertex decomposable graphs of the form θ_{n_1, \dots, n_k} .

Theorem 2.9. *Let n_1, \dots, n_k be a sequence of positive integers. Then, θ_{n_1, \dots, n_k} is vertex decomposable if and only if one of the followings holds:*

- The next result extends Proposition 2.8 to show that for $k = 3$, those graphs are not even sequentially Cohen-Macaulay.

Proof. Consider the labeling for $\theta_{2,2,5}$ and $\theta_{2,5,5}$ as given in Figure 1 and Figure 2. By [8, Lemma 2.3], the minimal generators of $\mathcal{I}(\theta_{2,2,5})^\vee$, correspond to the minimal vertex covers of $\theta_{2,2,5}$ and these minimal vertex covers correspond precisely to minimal prime ideals of $\mathcal{I}(\theta_{2,2,5})$. Therefore, by finding the minimal prime ideals of $\mathcal{I}(\theta_{2,2,5})$, the monomials $x_1x_2x_4x_6, x_1x_3x_4x_6, x_2x_4x_6x_7x_8, x_1x_3x_5x_6, x_2x_4x_5x_7x_8, x_2x_3x_5x_7x_8, x_1x_3x_5x_7x_8$, generate the ideal $\mathcal{I}(\theta_{2,2,5})^\vee$. With computation by CoCoA, we see that $\mathcal{I}(\theta_{2,2,5})_{[5]}^\vee$ has the minimal graded free resolution as

Thus, it does not have a linear resolution. Therefore, $\theta_{2,2,5}$ is not sequentially Cohen-Macaulay, by Theorem A.

$$\cdots \rightarrow R^{55}(-10)(+)R(-11) \rightarrow R^{121}(-9) \rightarrow R^{124}(-8) \rightarrow R^{49}(-7) \rightarrow R.$$

Figure 1




Figure 2

3

In view of Proposition 2.8 and Proposition 2.10, we conjecture that the answer to the following questions is positive.

Question 2.11. Let $K > 2$ and $\{n_1, \dots, n_k\} = \{2, 5\}$. Is θ_{n_1, \dots, n_k} not shellable? Is θ_{n_1, \dots, n_k} not sequentially Cohen-Macaulay?

Theorem 2.6 with [14, Theorem 2.9] enable us to get more examples of shellable and sequentially Cohen-Macaulay graphs.

Acknowledgments

We gratefully acknowledge the use of computer algebra systems CoCoA [4] which was invaluable in our work here. We also thank Professor Siamak Yassemi for valuable conversations and suggestions on these topics. The research of the first author was in part supported by a grant from IPM (No. 89050043). The research of the second author was in part supported by a grant from IPM (No. 87200116). Finally, we thank the referee for his or her extremely careful reading of our paper and very helpful corrections and suggestions for improvement. With the referee's comments we could extend our results from shellability to vertex decomposability.

REFERENCES

- [1] A. Björner and M. L. Wachs, Shellable nonpure complexes and posets I, *Trans. Amer. Math. Soc.* **348** (1996) 1299–1327.
- [2] A. Björner and M. L. Wachs, Shellable nonpure complexes and posets, II, *Trans. Amer. Math. Soc.* **349**(10) (1997) 3945–3975.
- [3] W. Bruns and J. Herzog, Cohen-Macaulay Rings, Cambridge University Press, 1993.
- [4] CoCoATeam, CoCoA: a system for doing Computations in Commutative Algebra, Available at <http://cocoa.dima.unige.it>
- [5] R. Diestel, Graph Theory, Springer-Verlag.
- [6] S. Faridi, Simplicial trees are sequentially Cohen-Macaulay, *J. Pure Appl. Algebra*. **190** (2003) 121–136.
- [7] C. A. Francisco and H. T. Hà, Whiskers and sequentially Cohen-Macaulay graphs, *J. Combin. Theory Ser. A*. **115** (2008) 304–316.
- [8] C. A. Francisco and A. V. Tuyl, Sequentially Cohen-Macaulay edge ideals, *Proc. Amer. Math. Soc.* **135** (2007) 2327–2337.
- [9] J. Herzog and T. Hibi, Componentwise linear ideals, *Nagoya Math. J.* **153** (1999) 141–153.
- [10] F. Mohammadi, D. Kiani and S. Yassemi, Shellable Cactus graphs, *Math. Scand.* **106** (2010) 161–167.
- [11] J. Scott Provan and L. J. Billera, Decompositions of simplicial complexes related to diameters of convex polyhedra, *Math. Oper. Res.* **5**(4) (1980) 576–594.

- [12] R. P. Stanley, *Combinatorics and Commutative Algebra*, Second edition, Progress in Mathematics **41**, Birkhäuser Boston, Inc., Boston, MA, 1996.
- [13] A. Van Tuyl, *Sequentially Cohen-Macaulay Bipartite Graphs: Vertex Decomposability and Regularity*, 2009, Preprint. arXiv:0906.0273v1.
- [14] A. Van Tuyl and R. Villarreal, Shellable graphs and sequentially Cohen-Macaulay bipartite graphs, *J. Combin. Theory, Ser. A.* **115**(5) (2008) 799–814.
- [15] M. L. Wachs, Obstructions to shellability, *Discrete Comput. Geom.* **22**(1) (1999) 95–103.
- [16] R. Woodroffe, Vertex decomposable graphs and obstruction to shellability, *Proc. Amer. Math. Soc.* **137**(10) (2009) 3235–3246.

Fatemeh Mohammadi and Dariush Kiani

Department of Pure Mathematics, Faculty of Mathematics and Computer Science, Amirkabir University of Technology (Tehran Polytechnic), 424 Hafez Ave., Tehran 15914, Iran.

and

School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box 19395-5746, Tehran, Iran.

Email: f_mohammadi@aut.ac.ir

Email: dkiani@aut.ac.ir

Archive of SID