

## SOME EQUIVALENCE CLASSES OF OPERATORS ON $\mathcal{B}(\mathcal{H})$

T. AGHASIZADEH AND S. HEJAZIAN\*

Communicated by Gholamhossein Eslamzadeh

**ABSTRACT.** Let  $\mathcal{L}(\mathcal{B}(\mathcal{H}))$  be the algebra of all linear operators on  $\mathcal{B}(\mathcal{H})$  and  $\mathcal{P}$  be a property on  $\mathcal{B}(\mathcal{H})$ . For  $\phi_1, \phi_2 \in \mathcal{L}(\mathcal{B}(\mathcal{H}))$ , we say that  $\phi_1 \sim_{\mathcal{P}} \phi_2$ , whenever  $\phi_1(T)$  has property  $\mathcal{P}$ , if and only if  $\phi_2(T)$  has this property. In particular, if  $\mathcal{I}$  is the identity map on  $\mathcal{B}(\mathcal{H})$ , then  $\phi \sim_{\mathcal{P}} \mathcal{I}$  means that  $\phi$  preserves property  $\mathcal{P}$  in both directions. Each property  $\mathcal{P}$  produces an equivalence relation on  $\mathcal{L}(\mathcal{B}(\mathcal{H}))$ . We study the relation between equivalence classes with respect to different properties such as being Fredholm, semi-Fredholm, compact, finite rank, generalized invertible, or having a specific semi-index.

### 1. Introduction

Let  $\mathcal{H}$  be an infinite-dimensional separable complex Hilbert space and  $\mathcal{B}(\mathcal{H})$  the algebra of all bounded linear operators on  $\mathcal{H}$ . We denote by  $\mathcal{F}(\mathcal{H})$  and  $\mathcal{K}(\mathcal{H})$  the ideals of all finite rank and compact operators in  $\mathcal{B}(\mathcal{H})$ , respectively. The Calkin algebra of  $\mathcal{H}$  is the quotient algebra  $\mathcal{C}(\mathcal{H}) = \mathcal{B}(\mathcal{H})/\mathcal{K}(\mathcal{H})$ . An operator  $T \in \mathcal{B}(\mathcal{H})$  is said to be a Fredholm operator if  $Im(T)$ , the range of  $T$ , is closed and both its kernel and co-kernel are finite-dimensional. We recall that  $T \in \mathcal{B}(\mathcal{H})$  is called upper (resp. lower) semi-Fredholm if  $Im(T)$  is closed and its kernel (resp.

---

MSC(2010): Primary: 47B49; Secondary: 47L05, 47A53.

Keywords: Fredholm operator, semi-Fredholm operator, generalized invertible operator, semi-index.

Received: 20 May 2009, Accepted: 30 December 2009.

\*Corresponding author

© 2011 Iranian Mathematical Society.

co-kernel) is finite-dimensional. An operator which is either upper semi-Fredholm or lower semi-Fredholm is called a semi-Fredholm operator. We denote by  $\mathcal{UF}(\mathcal{H})$ ,  $\mathcal{LF}(\mathcal{H})$ ,  $\mathcal{SF}(\mathcal{H})$ , and  $\mathcal{FR}(\mathcal{H})$  the sets of upper semi-Fredholm, lower semi-Fredholm, semi-Fredholm and Fredholm operators, respectively. By Atkinson's Theorem, [4, Theorem 1.4.16], if  $\mathcal{H}$  is an infinite-dimensional Hilbert space, then  $U \in \mathcal{B}(\mathcal{H})$  is Fredholm if and only if  $U + K(\mathcal{H})$  is invertible in the Calkin algebra  $\mathcal{C}(\mathcal{H})$ . The reader is referred to [4, 6] for more on Fredholm operators. Let  $A \in \mathcal{B}(\mathcal{H})$ . If there exists  $B \in \mathcal{B}(\mathcal{H})$  such that  $ABA = A$ , then  $A$  is called generalized invertible and  $B$  is said to be a generalized inverse of  $A$ . Note that  $A \in \mathcal{B}(\mathcal{H})$  is generalized invertible if and only if  $Im(A)$  is closed [5]. The set of generalized invertible elements of  $\mathcal{B}(\mathcal{H})$  is denoted by  $\mathcal{G}(\mathcal{H})$ .

The nullity (resp. defect) of an operator  $T \in \mathcal{B}(\mathcal{H})$  is defined to be  $dim(Ker(T))$  (resp.  $dim(coker(T))$ ), denoted by  $nul(T)$  (resp.  $def(T)$ ). Now, we define the function  $s-index : \mathcal{B}(\mathcal{H}) \rightarrow \{0, \infty\} \cup \mathbb{N}$  as follows:

$$s-index(T) = \begin{cases} \infty & T \in \mathcal{B}(\mathcal{H}) \setminus \mathcal{SF}(\mathcal{H}), \\ 0 & T \in \mathcal{FR}(\mathcal{H}), \\ nul(T) & T \in \mathcal{UF}(\mathcal{H}) \setminus \mathcal{FR}(\mathcal{H}), \\ def(T) & T \in \mathcal{LF}(\mathcal{H}) \setminus \mathcal{FR}(\mathcal{H}). \end{cases}$$

The number  $s-index(T)$  is called the semi-index of  $T$ . Note that for a Fredholm operator  $T$ , in general,  $s-index(T)$  does not coincide with the classical index of  $T$  which is defined by  $nul(T) - def(T)$ .

Let  $\mathcal{L}(\mathcal{B}(\mathcal{H}))$  be the set of all linear mappings on  $\mathcal{B}(\mathcal{H})$ . Recall that  $\phi \in \mathcal{L}(\mathcal{B}(\mathcal{H}))$  is said to be surjective up to finite rank operators if  $\mathcal{B}(\mathcal{H}) = Im(\phi) + \mathcal{F}(\mathcal{H})$ , and  $\phi$  is said to be surjective up to compact operators if  $\mathcal{B}(\mathcal{H}) = Im(\phi) + \mathcal{K}(\mathcal{H})$ . Obviously, if  $\phi$  is surjective up to finite rank operators, then it is surjective up to compact operators and each surjective linear map satisfies both of these properties.

Let  $\mathcal{P}$  be a property on  $\mathcal{B}(\mathcal{H})$ . For  $\phi_1, \phi_2 \in \mathcal{L}(\mathcal{B}(\mathcal{H}))$ , we say that  $\phi_1 \sim_{\mathcal{P}} \phi_2$ , whenever  $\phi_1(T)$  has property  $\mathcal{P}$ , if and only if  $\phi_2(T)$  has this property. It is easy to see that each property  $\mathcal{P}$  produces an equivalence relation on  $\mathcal{L}(\mathcal{B}(\mathcal{H}))$ . Throughout this paper, we use the following notations for some specific properties:

- (i) “ $f$ ” is the property of “being finite-rank”;
- (ii) “ $k$ ” is the property of “being compact”;
- (iii) “ $fr$ ” is the property of “being Fredholm”;
- (iv) “ $sf$ ” is the property of “being semi-Fredholm”;

- (v) “g” is the property of “being generalized invertible”;  
 (vi) “si” is the property of “having a specific semi-index”.

Let  $\mathcal{I}$  denote the identity operator of  $\mathcal{L}(\mathcal{B}(\mathcal{H}))$  and  $\phi \in \mathcal{L}(\mathcal{B}(\mathcal{H}))$ . Then,  $\phi \sim_{\mathcal{P}} \mathcal{I}$  means that  $\phi$  preserves the property  $\mathcal{P}$  in both directions, that is,  $\phi(T)$  has property  $\mathcal{P}$  if and only if  $T$  has this property. Mbekhta and Šemrl in [3] study those  $\phi$  which satisfy  $\phi \sim_g \mathcal{I}$  and  $\phi \sim_{sf} \mathcal{I}$ . In general, if  $\psi$  is a linear operator on  $\mathcal{B}(\mathcal{H})$ , which preserves property  $\mathcal{P}$  in both directions, then  $\psi\phi \sim_{\mathcal{P}} \phi$ , for all  $\phi \in \mathcal{L}(\mathcal{B}(\mathcal{H}))$ . Also, if  $v$  is a linear operator on  $\mathcal{B}(\mathcal{H})$ , which does not preserve property  $\mathcal{P}$  in both directions, then for each surjective linear operator  $\phi$ ,  $v\phi \not\sim_{\mathcal{P}} \phi$ .

In the next section, we study the equivalence classes with respect to the above properties. We show that for surjective up to finite rank mappings  $\phi_1$  and  $\phi_2$  in  $\mathcal{L}(\mathcal{B}(\mathcal{H}))$ ,  $\phi_1 \sim_g \phi_2$  implies that  $\phi_1 \sim_{sf} \phi_2$  and  $\phi_1 \sim_f \phi_2$ . Also, if  $\phi_1, \phi_2$  are linear mappings on  $\mathcal{B}(\mathcal{H})$ , which are surjective up to compact operators, and  $\phi_1 \sim_{sf} \phi_2$  or  $\phi_1 \sim_{fr} \phi_2$ , then  $\phi_1 \sim_k \phi_2$ . It is also proved that  $\phi_1 \sim_{si} \phi_2$  implies  $\phi_1 \sim_{sf} \phi_2$ . We give some examples to illustrate that some of the reverse implications do not hold, in general. We also prove that for surjective linear operators  $\phi_1, \phi_2 \in \mathcal{L}(\mathcal{B}(\mathcal{H}))$ ,  $\phi_1 \sim_{si} \phi_2$  and  $\phi_1 \sim_f \phi_2$  imply that  $Ker(\phi_1) = Ker(\phi_2)$ , and we give an example to show that the converse is not true, in general. Finally, it is proved that if  $\phi_1, \phi_2$  are bijections such that  $\phi_1 \sim_{sf} \phi_2$ , then  $\phi_1\phi_2^{-1}$  induces a map  $\tilde{\psi} : \mathcal{C}(\mathcal{H}) \rightarrow \mathcal{C}(\mathcal{H})$ , which is either an automorphism or an anti-automorphism, multiplied by an invertible element  $A \in \mathcal{C}(\mathcal{H})$ .

## 2. The Results

In the following lemma, (i)  $\Leftrightarrow$  (ii) comes from [2, Lemma 2.2] and (i)  $\Leftrightarrow$  (iii) comes from [3, Lemma 2.2].

**Lemma 2.1.** *Let  $K \in \mathcal{B}(\mathcal{H})$ . Then, the following are equivalent.*

- (i)  $K$  is compact.
- (ii) for every  $B \in \mathcal{FR}(\mathcal{H})$ , we have  $B + K \in \mathcal{FR}(\mathcal{H})$ .
- (iii) for every  $B \in \mathcal{SF}(\mathcal{H})$ , we have  $B + K \in \mathcal{SF}(\mathcal{H})$ .

Take  $C = \{T \in \mathcal{B}(\mathcal{H}) \mid \text{for every operator } A \in \mathcal{B}(\mathcal{H}) \text{ with } Im(A) \text{ not closed, there exists } \lambda \in \mathbb{C} \text{ such that } A + \lambda T \neq 0 \text{ and } Im(A + \lambda T) \text{ is closed}\}$ . It is proved in [1, Lemma 3.1] that  $C = \mathcal{SF}(\mathcal{H})$ .

We recall that if  $T \in \mathcal{G}(\mathcal{H})$ , then, for each finite rank operator  $F$ , we have  $T + F \in \mathcal{G}(\mathcal{H})$ .

**Theorem 2.2.** Let  $\phi_1, \phi_2 : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  be linear mappings. Then,

$$(i) \quad \phi_1 \sim_{si} \phi_2 \Rightarrow \phi_1 \sim_{sf} \phi_2;$$

if  $\phi_1, \phi_2$  are surjective up to finite rank operators, then

$$(ii) \quad \phi_1 \sim_g \phi_2 \Rightarrow \phi_1 \sim_{sf} \phi_2;$$

$$(iii) \quad \phi_1 \sim_g \phi_2 \Rightarrow \phi_1 \sim_f \phi_2;$$

if  $\phi_1, \phi_2$  are surjective up to compact operators, then

$$(iv) \quad \phi_1 \sim_{fr} \phi_2 \Rightarrow \phi_1 \sim_k \phi_2;$$

$$(v) \quad \phi_1 \sim_{sf} \phi_2 \Rightarrow \phi_1 \sim_k \phi_2.$$

*Proof.* (i) It is trivial by the definition of semi-index.

(ii) Suppose that  $\phi_1(T)$  is a semi-Fredholm operator. We show that  $\phi_2(T) \in C$ . Suppose that  $B \in \mathcal{B}(\mathcal{H})$  is not generalized invertible, or equivalently  $Im(B)$  is not closed. Since  $\phi_2$  is surjective up to finite rank operators, there exist  $A \in \mathcal{B}(\mathcal{H})$  and  $F \in \mathcal{F}(\mathcal{H})$  such that  $\phi_2(A) = B + F$ . Since  $F$  is finite rank,  $Im(\phi_2(A))$  is not closed and it follows that the range of  $\phi_1(A)$  is not closed. Since  $\phi_1(T)$  is semi-Fredholm, we have  $\phi_1(T) \in C$ . Thus, there exists  $\alpha \in \mathbb{C}$  such that  $Im(\phi_1(\alpha T + A))$  is closed. It follows that the range of  $\phi_2(\alpha T + A) = \alpha\phi_2(T) + B + F$  is also closed, which implies that  $Im(\alpha\phi_2(T) + B)$  is closed. Note that  $\alpha\phi_2(T) + B \neq 0$ . Otherwise,  $Im(\alpha\phi_2(T)) = Im(B)$  is not closed, which contradicts  $\phi_1 \sim_g \phi_2$ . Therefore,  $\phi_2(T) \in C$ .

(iii) Let  $\phi_1 \sim_g \phi_2$ . Suppose that  $\phi_1(T) \in \mathcal{F}(\mathcal{H})$ , but  $\phi_2(T)$  is not finite-rank. Therefore, the range of  $\phi_1(T)$  is closed, but it is not semi-Fredholm. By the fact that  $\phi_1 \sim_g \phi_2$ ,  $Im(\phi_2(T))$  is closed. Also, by (ii),  $\phi_2(T)$  is not semi-Fredholm. Take  $S = \phi_2(T)$ . Then, both  $Ker(S)$  and  $Im(S)^\perp$  are infinite-dimensional and we can define a bounded linear bijection  $S' : Ker(S) \rightarrow Im(S)^\perp$ . Extend  $S'$  on  $\mathcal{H}$  by  $S'(x) = 0$ , for all  $x \in Ker(S)^\perp$ , and denote this extension by  $S'$  as well. Since  $S$  is not finite rank,  $S'$  is not a semi-Fredholm operator on  $\mathcal{H}$ . Now, take  $\tilde{T} \in \mathcal{B}(\mathcal{H})$  and  $F \in \mathcal{F}(\mathcal{H})$  such that  $\phi_2(\tilde{T}) = S' + F$ . We have  $S + S'$  is a bijective bounded linear operator on  $\mathcal{H}$ , and hence it is Fredholm. Therefore,  $\phi_2(T + \tilde{T}) = S + S' + F \in \mathcal{FR}(\mathcal{H})$ . On the other hand,  $\phi_1(T + \tilde{T})$  is not semi-Fredholm. Otherwise,  $\phi_1(\tilde{T})$  must be semi-Fredholm and it follows by (ii) that  $S' + F = \phi_2(\tilde{T})$  is semi-Fredholm, which is not correct. Thus,  $\phi_1 \not\sim_{sf} \phi_2$ , a contradiction with (ii), and so  $\phi_1 \sim_f \phi_2$ .

(iv) Suppose that  $\phi_1(T)$  is compact. Let  $S$  be an arbitrary Fredholm operator. Since  $\phi_2$  is surjective up to compact operators, there exist  $A \in \mathcal{B}(\mathcal{H})$  and  $K \in \mathcal{K}(\mathcal{H})$  such that  $\phi_2(A) = S + K$ . Obviously,  $\phi_2(A) \in$

$\mathcal{FR}(\mathcal{H})$ , and since  $\phi_1 \sim_{fr} \phi_2$ , we have  $\phi_1(A)$  is Fredholm. On the other hand,  $\phi_1(T)$  is compact and so by Lemma 2.1,  $\phi_1(T + A) \in \mathcal{FR}(\mathcal{H})$ . Thus,  $\phi_2(T + A)$  is Fredholm, and it follows that  $\phi_2(T) + S = \phi_2(T + A) - K$  is also a Fredholm operator and Lemma 2.1 implies that  $\phi_2(T)$  is compact.

(v) The proof is similar to the one given in (iv).  $\square$

In what follows we give some examples to show that in Theorem 2.2 some of the reverse implications do not hold, in general.

**Example 2.3.** Suppose that  $\phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  is a surjective linear map and  $S \in \mathcal{B}(\mathcal{H})$  is a lower semi-Fredholm operator such that  $s\text{-index}(S) = 1$ . Since  $\phi$  is surjective, there exists  $T \in \mathcal{B}(\mathcal{H})$  such that  $\phi(T) = S$ . Now, if  $A \in \mathcal{B}(\mathcal{H})$  is a Fredholm operator with  $\text{def}(A) = 2$ , then  $\phi \sim_{sf} L_A\phi$ , but  $\phi \not\sim_{si} L_A\phi$ , since  $s\text{-index}(A\phi(T)) \geq 2 > 1$ . Here  $L_A : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  is the left multiplier operator defined by  $L_A(S) = AS$ .

**Example 2.4.** We show that, in general,  $\phi_1 \sim_{sf} \phi_2$  or  $\phi_1 \sim_{fr} \phi_2$  does not imply  $\phi_1 \sim_g \phi_2$ . Let  $S$  be a surjective bounded linear map on  $\mathcal{H}$  such that  $\dim(\text{Ker}(S)) = 1$ . Note that  $S \in \mathcal{FR}(\mathcal{H})$ . Let  $P : \mathcal{H} \rightarrow \text{Ker}(S)$  be the projection of  $\mathcal{H}$  onto  $\text{Ker}(S)$ . Take  $\phi_0 = L_S$ . Since  $S$  has a bounded right inverse,  $\phi_0$  is surjective. Extend  $\{P\}$  to a vector space basis  $\{T_\alpha\}$  for  $\mathcal{B}(\mathcal{H})$ . Suppose that  $K \in \mathcal{K}(\mathcal{H})$  has a non-closed range. Define a linear map  $\lambda : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{K}(\mathcal{H})$  by

$$\lambda(T_\alpha) = \begin{cases} K & T_\alpha = P \\ 0 & T_\alpha \neq P. \end{cases}$$

Now, define  $\phi_1 : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  by  $\phi_0(T) + \lambda(T)$ . Then, by Lemma 2.1,  $\phi_1 \sim_{sf} \phi_0$ . Also,  $\phi_1$  is surjective. To see this, take  $T \in \mathcal{B}(\mathcal{H})$ . There exists  $U \in \mathcal{B}(\mathcal{H})$  such that  $\phi_0(U) = T$ . Since  $\{T_\alpha\}$  is a vector space basis for  $\mathcal{B}(\mathcal{H})$ , there exist  $\beta_1, \dots, \beta_n \in \mathbb{C}$  and  $T_{\alpha_1}, \dots, T_{\alpha_n} \in \{T_\alpha\}$  such that  $U = \sum_{i=1}^n \beta_i T_{\alpha_i}$ . If for each  $1 \leq j \leq n$ ,  $T_{\alpha_j} \neq P$ , then  $\phi_1(U) = \phi_0(U) = T$ . Otherwise, if for some  $1 \leq j \leq n$ ,  $T_{\alpha_j} = P$ , then take  $U' = U - \beta_j T_{\alpha_j}$  and we have  $\phi_0(U) = \phi_0(U')$ , and therefore,  $\phi_1(U') = \phi_0(U') + \lambda(U') = T$ .

Finally,  $\phi_1(P) = K$ , which is not generalized invertible since  $\text{Im}(K)$  is not closed, while  $\phi_0(P) = 0$  is generalized invertible and this shows  $\phi_1 \not\sim_g \phi_0$ .

We do not know any example of two linear mappings  $\phi_1, \phi_2 \in \mathcal{L}(\mathcal{B}(\mathcal{H}))$ , which are surjective up to compact operators,  $\phi_1 \sim_k \phi_2$  but  $\phi_1 \not\sim_{fr} \phi_2$  or  $\phi_1 \not\sim_{sf} \phi_2$ . As seen in the above examples, the multiplier operator  $L_T$

for a suitable  $T$  plays an important role. But, here we show that if  $\phi_1$  and  $\phi_2$  satisfy the above mentioned conditions, they can not be related by  $\phi_2 = L_A R_B \phi_1 + \lambda$ , where  $\lambda$  is a linear mapping from  $\mathcal{B}(\mathcal{H})$  to  $\mathcal{K}(\mathcal{H})$ . Here,  $R_B$  denotes the right multiplier operator  $T \mapsto TB$  on  $\mathcal{B}(\mathcal{H})$ .

**Proposition 2.5.** *Let  $\phi_1, \phi_2 \in \mathcal{L}(\mathcal{B}(\mathcal{H}))$  be surjective up to compact operators and  $\phi_2 = L_A R_B \phi_1 + \lambda$ , where,  $\lambda : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{K}(\mathcal{H})$  is a linear mapping. If  $\phi_1 \sim_k \phi_2$ , then  $A$  and  $B$  are Fredholm operators, and hence  $\phi_1 \sim_{fr} \phi_2$  and  $\phi_1 \sim_{sf} \phi_2$ .*

*Proof.* Let  $\phi_2 = L_A R_B \phi_1 + \lambda$ . For  $i = 1, 2$ , consider  $\tau_i : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{C}(\mathcal{H})$  defined by  $\tau_i(T) = \pi \circ \phi_i(T)$ , where,  $\pi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{C}(\mathcal{H})$  is the canonical quotient map. It is easy to check that  $\tau_2(T) = a\tau_1(T)b$ , for all  $T \in \mathcal{B}(\mathcal{H})$ , where,  $a = \pi(A), b = \pi(B)$ .

The condition that  $\phi_1, \phi_2$  are surjective up to compact operators implies that  $\tau_1, \tau_2$  are surjective. The condition  $\phi_1 \sim_k \phi_2$  says that  $\tau_1(T) = 0$  if and only if  $\tau_2(T) = 0$  if and only if  $a\tau_1(T)b = 0$ . Since  $\tau_1$  is onto, this in turn says that with  $x \in \mathcal{C}(\mathcal{H})$ ,  $axb = 0$  if and only if  $x = 0$ .

Now,  $\tau_2$  is onto, and so  $azb = \pi(I)$ , for some  $z \in \mathcal{C}(\mathcal{H})$ . Thus, there exists  $Z \in \mathcal{B}(\mathcal{H})$  such that  $AZB$  is a Fredholm operator, which shows that  $A$  and  $B$  are semi-Fredholm. If  $A$  were not Fredholm, then (since  $a = \pi(A)$  is right invertible), we must have  $nul(A) = \infty$ . Let  $P \in \mathcal{B}(\mathcal{H})$  be the orthogonal projection of  $\mathcal{H}$  onto  $Ker(A)$ . Then,  $p = \pi(P) \neq 0$ , but  $apb = \pi(APB) = \pi(0B) = 0$ , which is a contradiction. Thus,  $A$  is Fredholm. Finally, since  $AZB$  is a Fredholm operator, we have that  $B^*Z^*A^*$  is also a Fredholm operator. The same argument implies that  $B^*$ , and hence  $B$  is a Fredholm operator.  $\square$

In the sequel, we explore the consequences when both  $\phi_1$  and  $\phi_2$  are in certain equivalence classes.

**Theorem 2.6.** *If  $\phi_1, \phi_2 : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  are surjective linear maps such that  $\phi_1 \sim_{si} \phi_2$  and  $\phi_1 \sim_f \phi_2$ , then  $Ker(\phi_1) = Ker(\phi_2)$ .*

*Proof.* Let  $T \in Ker(\phi_1)$ . Since  $\phi_1 \sim_f \phi_2$ ,  $\phi_2(T)$  is finite-rank, then it is not a semi-Fredholm operator. It follows that  $null(\phi_2(T)) = \infty = def(\phi_2(T))$ . Now, we can write  $Ker(\phi_2(T)) = \mathcal{M} \oplus \mathcal{N}$ , where  $\mathcal{M}$  and  $\mathcal{N}$  are infinite-dimensional closed subspaces of  $Ker(\phi_2(T))$ . Define a bounded linear bijection  $T' : \mathcal{N} \rightarrow Im(\phi_2(T))^\perp$ . Extend  $T'$  on  $\mathcal{H}$  by  $T'(x) = 0$ , for all  $x \in (Ker(\phi_2(T)))^\perp \oplus \mathcal{M}$  and denote this extension by  $T'$  as well. Clearly,  $T'$  is a semi-Fredholm operator and

$s\text{-index}(T') = \text{rank}(\phi_2(T))$ . We have  $\phi_2(T) + T'$  is surjective and  $\mathcal{M} \subseteq \text{Ker}(\phi_2(T) + T')$ . Thus, it is a semi-Fredholm operator with  $s\text{-index}(\phi_2(T) + T') = 0$ . Since  $\phi_2$  is surjective, there exists  $S \in \mathcal{B}(\mathcal{H})$  such that  $\phi_2(S) = T'$ . Therefore,  $\text{rank}(\phi_2(T)) = s\text{-index}(\phi_2(S)) = s\text{-index}(\phi_1(S)) = s\text{-index}(\phi_1(S + T)) = s\text{-index}(\phi_2(S + T)) = 0$ . It follows that  $\text{Ker}(\phi_1) \subseteq \text{Ker}(\phi_2)$ . The reverse inclusion follows similarly and we have the result.  $\square$

**Corollary 2.7.** *If  $\phi$  is a surjective linear map on  $\mathcal{B}(\mathcal{H})$  that preserves finite rank operators and semi-index property in both directions, then  $\phi$  is injective.*

As a consequence, by Theorem 2.2 (iii), if  $\phi$  is a surjective linear map on  $\mathcal{B}(\mathcal{H})$  that preserves generalized invertible operators and semi-index property in both directions, then it is injective.

**Remark 2.8.** (i) *In general, if surjective linear maps  $\phi_1, \phi_2 : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  have the same kernels, then it may happen that  $\phi_1 \approx_f \phi_2$ , and hence the converse of Theorem 2.6 does not hold. To see this, take  $e \in \mathcal{H}$  with  $\|e\| = 1$ . It is clear that  $I$  and  $e \otimes e$  are linearly independent. We can extend  $\{I, e \otimes e\}$  to a basis for the vector space  $\mathcal{B}(\mathcal{H})$ . Now, define  $\phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  such that  $\phi(I) = e \otimes e$ ,  $\phi(e \otimes e) = I$  and  $\phi(T) = T$ , for every  $T$  in the basis with  $T \neq I, T \neq e \otimes e$ . Extend  $\phi$  to a bijection on  $\mathcal{B}(\mathcal{H})$ , by linearity. Hence,  $\text{Ker}(\phi) = 0 = \text{Ker}(I)$ , but we do not have  $\phi \sim_f I$ , since  $\phi(I) = e \otimes e$ . It also follows that  $\phi \approx_g I$ , since otherwise by Theorem 2.2 (iii), we must have  $\phi \sim_f I$ .*

(ii) *The condition  $\phi_1 \sim_{si} \phi_2$  in Theorem 2.6 can not be omitted. Suppose that  $S \in \mathcal{B}(\mathcal{H})$  is surjective and  $\dim(\text{Ker}(S)) = 1$ . Thus,  $I \sim_f L_S$ ,  $I \sim_{si} L_S$ , and clearly  $\text{Ker}(I) \neq \text{Ker}(L_S)$ . We do not know any example of surjective linear operators  $\phi_1, \phi_2$  on  $\mathcal{B}(\mathcal{H})$  such that  $\phi_1 \sim_{si} \phi_2$ , but  $\phi \not\sim_f \phi_2$ . So, at this point we do not know whether the case  $\phi_1 \sim_{si} \phi_2$ ,  $\phi_1 \not\sim_f \phi_2$  happens or not.*

Now, we consider the case  $\text{Ker}(\phi_1) = \{0\} = \text{Ker}(\phi_2)$ .

**Proposition 2.9.** *Let  $\mathcal{P}$  be a property on  $\mathcal{B}(\mathcal{H})$ . If  $\phi_1, \phi_2 : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  are bijective linear maps such that  $\phi_1 \sim_{\mathcal{P}} \phi_2$ , then  $\phi_1 \phi_2^{-1}$  preserves property  $\mathcal{P}$  in both directions.*

*Proof.* Let  $\mathcal{M}_{\mathcal{P}} = \{T \in \mathcal{B}(\mathcal{H}) : T \text{ has property } \mathcal{P}\}$ . Then,  $\phi_1^{-1}(\mathcal{M}_{\mathcal{P}}) = \phi_2^{-1}(\mathcal{M}_{\mathcal{P}})$ , and hence  $\mathcal{M}_{\mathcal{P}} = \phi_1 \phi_2^{-1}(\mathcal{M}_{\mathcal{P}})$ . It follows that  $\phi_1 \phi_2^{-1}$  preserves  $\mathcal{P}$  in both directions.  $\square$

The following theorem was proved by Mbekhta and Šemrl [3, Theorem 1.2].

**Theorem 2.10.** *Let  $\mathcal{H}$  be an infinite-dimensional separable Hilbert space and  $\phi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  be a linear map preserving semi-Fredholm operators in both directions. Suppose that  $\phi$  is surjective up to compact operators. Then,*

$$\phi(\mathcal{K}(\mathcal{H})) \subseteq \mathcal{K}(\mathcal{H}),$$

*and the induced map  $\tilde{\phi} : \mathcal{C}(\mathcal{H}) \rightarrow \mathcal{C}(\mathcal{H})$  is either an automorphism, or an anti-automorphism multiplied by an invertible element  $a \in \mathcal{C}(\mathcal{H})$ .*

**Corollary 2.11.** *Suppose that  $\phi_1, \phi_2$  are bijective linear maps on  $\mathcal{B}(\mathcal{H})$  such that  $\phi_1 \sim_{sf} \phi_2$ . Take  $\psi = \phi_1 \phi_2^{-1}$ . Then,  $\psi(\mathcal{K}(\mathcal{H})) = \mathcal{K}(\mathcal{H})$  and the induced map  $\tilde{\psi}$  on  $\mathcal{C}(\mathcal{H})$  is an automorphism or an anti-automorphism multiplied by an invertible element  $a \in \mathcal{C}(\mathcal{H})$ .*

Note that, by Theorem 2.2 (ii), we have the same result for  $\phi_1 \sim_g \phi_2$ . Now, a question comes to mind: *Is it possible to identify the equivalence class of  $\phi$  with respect to a property  $\mathcal{P}$ ?*

**Remark 2.12.** (i) *Let  $\tau : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$  be the linear map which takes  $T$  to its transpose with respect to a given basis for  $\mathcal{H}$ . If  $\phi \in \mathcal{L}(\mathcal{B}(\mathcal{H}))$ , then it is easy to see that  $\tau\phi \sim_g \phi$ , and hence  $\tau\phi \sim_{sf} \phi$ ,  $\tau\phi \sim_k \phi$ , and  $\tau\phi \sim_f \phi$ . Also,  $\tau\phi \sim_{fr} \phi$  and  $\tau\phi \sim_{si} \phi$ .*

(ii) *Let  $A$  and  $B$  be Fredholm operators and  $\lambda : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{K}(\mathcal{H})$  be a linear map. If  $\phi_1, \phi_2 \in \mathcal{L}(\mathcal{B}(\mathcal{H}))$  are related as  $\phi_2 = L_A R_B \phi_1 + \lambda$  or  $\phi_2 = L_A R_B \tau \phi_1 + \lambda$ , then it is easy to see that  $\phi_1 \sim_{sf} \phi_2$ ,  $\phi_1 \sim_{fr} \phi_2$ .*

(iii) *Let  $A$  and  $B$  be Fredholm operators and  $\lambda : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{F}(\mathcal{H})$  be a linear map. If  $\phi_1, \phi_2 \in \mathcal{L}(\mathcal{B}(\mathcal{H}))$  are such that  $\phi_2 = L_A R_B \phi_1 + \lambda$  or  $\phi_2 = L_A R_B \tau \phi_1 + \lambda$ , then  $\phi_1 \sim_g \phi_2$ .*

(iv) *Let  $A, B \in \mathcal{B}(\mathcal{H})$  be invertible operators. If  $\phi_2 = L_A R_B \phi_1$  or  $\phi_2 = L_A R_B \tau \phi_1$ , then it is easy to see that  $\phi_1 \sim_{si} \phi_2$ .*

**Question 2.13.** *Let  $\phi_1 \sim_g \phi_2$ . Are there  $A, B \in \mathcal{FR}(\mathcal{H})$  and a linear map  $\lambda : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{F}(\mathcal{H})$  such that  $\phi_2 = L_A R_B \phi_1 + \lambda$  or  $\phi_2 = L_A R_B \tau \phi_1 + \lambda$ ?*

**Question 2.14.** *Let  $\phi_1 \sim_{sf} \phi_2$  or  $\phi_1 \sim_{fr} \phi_2$ . Are there  $A, B \in \mathcal{FR}(\mathcal{H})$  and a linear map  $\lambda : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{K}(\mathcal{H})$  such that  $\phi_2 = L_A R_B \phi_1 + \lambda$  or  $\phi_2 = L_A R_B \tau \phi_1 + \lambda$ ?*



**Question 2.15.** Let  $\phi_1 \sim_{si} \phi_2$ . Are there invertible operators  $A, B \in \mathcal{B}(\mathcal{H})$  such that  $\phi_2 = L_A R_B \phi_1$  or  $\phi_2 = L_A R_{B\tau} \phi_1$ ?

### Acknowledgments

The authors sincerely thank the referee for the valuable comments and for suggesting the proof of Proposition 2.5.

### REFERENCES

- [1] J. Hou and J. Cui, Linear maps preserving essential spectral functions and closeness of operator ranges, *Bull. Lond. Math. Soc.* **39** (2007) 575-582.
- [2] M. Mbekhta, Linear maps preserving the set of Fredholm operators, *Proc. Amer. Math. Soc.* **135** (2007) 3613-3619.
- [3] M. Mbekhta and P. Šemrl, Linear maps preserving semi-Fredholm operators and generalized invertibility, *Linear Multilinear Algebra* **57** (2009) 55-64.
- [4] G. J. Murphy, *C\*-Algebras and Operator Theory*, Academic Press, Inc., Boston, MA, 1990.
- [5] M. Z. Nashed (ed.), *Generalized inverses and applications*, Univ. Wisconsin Math. Res. Center Publ., No. 32, Academic Press, New York, 1976.
- [6] N. E. Wegge-Olsen, *K-Theory and C\*-Algebras, a Friendly Approach*, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1993.

#### T. Aghasizadeh

Department of Pure Mathematics, Ferdowsi University of Mashhad, P.O. Box 1159-91775, Mashhad, Iran

Email: to\_ag99@stu.um.ac.ir

#### S. Hejazian

Department of Pure Mathematics, Ferdowsi University of Mashhad, P.O. Box 1159-91775, Mashhad, Iran

Center of Excellence in Analysis on Algebraic Structures (CEAAS), Ferdowsi University of Mashhad, P.O. Box 1159-91775, Iran

Tusi Mathematical Research Group (TMRG), Mashhad, Iran

Email: hejazian@ferdowsi.um.ac.ir