

## ONE-POINT EXTENSIONS OF LOCALLY COMPACT PARACOMPACT SPACES

M. R. KOUSHESH

Communicated by Fereidoun Ghahramani

**ABSTRACT.** A space  $Y$  is called an *extension* of a space  $X$ , if  $Y$  contains  $X$  as a dense subspace. Two extensions of  $X$  are said to be *equivalent*, if there is a homeomorphism between them which fixes  $X$  point-wise. For two (equivalence classes of) extensions  $Y$  and  $Y'$  of  $X$  let  $Y \leq Y'$ , if there is a continuous function of  $Y'$  into  $Y$  which fixes  $X$  point-wise. An extension  $Y$  of  $X$  is called a *one-point extension*, if  $Y \setminus X$  is a singleton. An extension  $Y$  of  $X$  is called *first-countable*, if  $Y$  is first-countable at points of  $Y \setminus X$ . Let  $\mathcal{P}$  be a topological property. An extension  $Y$  of  $X$  is called a  *$\mathcal{P}$ -extension*, if it has  $\mathcal{P}$ .

In this article, for a given locally compact paracompact space  $X$ , we consider the two classes of one-point Čech-complete;  $\mathcal{P}$ -extensions of  $X$  and one-point first-countable locally- $\mathcal{P}$  extensions of  $X$ , and we study their order-structures, by relating them to the topology of a certain subspace of the outgrowth  $\beta X \setminus X$ . Here  $\mathcal{P}$  is subject to some requirements and include  $\sigma$ -compactness and the Lindelöf property as special cases.

### 1. Introduction

A space  $Y$  is called an *extension* of a space  $X$ , if  $Y$  contains  $X$  as a dense subspace. If  $Y$  is an extension of  $X$ , then the subspace  $Y \setminus X$  of

---

MSC(2010): Primary: 54D20; Secondary: 54D35, 54D40, 54D45, 54E50.

Keywords: Stone-Čech compactification, one-point extension, one-point compactification, locally compact, paracompact, Čech complete; first-countable.

Received: 16 March 2010, Accepted: 21 July 2010.

© 2011 Iranian Mathematical Society.

$Y$  is called the *remainder* of  $X$ . Extensions with a one-point remainder are called *one-point extensions*. Two extensions of  $X$  are said to be *equivalent*, if there exists a homeomorphism between them which fixes  $X$  point-wise. This defines an equivalence relation on the class of all extensions of  $X$ . The equivalence classes will be identified with individuals when this causes no confusion. For two extensions  $Y$  and  $Y'$  of  $X$  we let  $Y \leq Y'$ , if there exists a continuous function of  $Y'$  into  $Y$  which fixes  $X$  point-wise. The relation  $\leq$  defines a partial order on the set of extensions of  $X$  (see Section 4.1 of [16] for more details). An extension  $Y$  of  $X$  is called *first-countable*, if  $Y$  is first-countable at points of  $Y \setminus X$ , that is,  $Y$  has a countable local base at every point of  $Y \setminus X$ . Let  $\mathcal{P}$  be a topological property. An extension  $Y$  of  $X$  is called a  $\mathcal{P}$ -*extension*, if it has  $\mathcal{P}$ . If  $\mathcal{P}$  is compactness, then  $\mathcal{P}$ -extensions are called *compactifications*.

This work was mainly motivated by our previous work [9] (see [1], [7], [8], [11], [12] and [13] for related results) in which we have studied the partially ordered set of one-point  $\mathcal{P}$ -extensions of a given locally compact space  $X$  by relating it to the topologies of certain subspaces of its outgrowth  $\beta X \setminus X$ . In this article, we continue our studies by considering the classes of one-point Čech-complete  $\mathcal{P}$ -extensions and one-point first-countable locally- $\mathcal{P}$  extensions of a given locally compact paracompact space  $X$ . The topological property  $\mathcal{P}$  is subject to some requirements and include  $\sigma$ -compactness, the Lindelöf property and the linearly Lindelöf property as special cases.

We review some of the terminology, notation and well-known results that will be used in the sequel. Our definitions mainly come from the standard text [3] (thus, in particular, compact spaces are Hausdorff, etc.). Other useful sources are [5] and [16].

The letters  $\mathbf{I}$  and  $\mathbf{N}$  denote the closed unit interval and the set of all positive integers, respectively. For a subset  $A$  of a space  $X$  we let  $\text{cl}_X A$  and  $\text{int}_X A$  denote the closure and the interior of  $A$  in  $X$ , respectively. A subset of a space is called *clopen*, if it is simultaneously closed and open. A *zero-set* of a space  $X$  is a set of the form  $Z(f) = f^{-1}(0)$  for some continuous  $f : X \rightarrow \mathbf{I}$ . Any set of the form  $X \setminus Z$ , where  $Z$  is a zero-set of  $X$ , is called a *cozero-set* of  $X$ . We denote the set of all zero-sets of  $X$  by  $\mathcal{Z}(X)$  and the set of all cozero-sets of  $X$  by  $\text{Coz}(X)$ .

For a Tychonoff space  $X$  the *Stone-Čech compactification* of  $X$  is the largest (with respect to the partial order  $\leq$ ) compactification of  $X$  and is denoted by  $\beta X$ . The Stone-Čech compactification of  $X$  can be

characterized among all compactifications of  $X$  by either of the following properties:

- (1) Every continuous function of  $X$  to a compact space is continuously extendible over  $\beta X$ .
- (2) Every continuous function of  $X$  to  $\mathbf{I}$  is continuously extendible over  $\beta X$ .
- (3) For every  $Z, S \in \mathcal{Z}(X)$  we have

$$\text{cl}_{\beta X}(Z \cap S) = \text{cl}_{\beta X}Z \cap \text{cl}_{\beta X}S.$$

A Tychonoff space is called *zero-dimensional*, if it has an open base consisting of its clopen subsets. A Tychonoff space is called *strongly zero-dimensional*, if its Stone-Čech compactification is zero-dimensional. A Tychonoff space  $X$  is called *Čech-complete*, if its outgrowth  $\beta X \setminus X$  is an  $F_\sigma$  in  $\beta X$ . Locally compact spaces are Čech-complete, and in the realm of metrizable spaces  $X$ , Čech-completeness is equivalent to the existence of a compatible complete metric on  $X$ .

Let  $\mathcal{P}$  be a topological property. A topological space  $X$  is called *locally- $\mathcal{P}$* , if for every  $x \in X$  there exists an open neighborhood  $U_x$  of  $x$  in  $X$  such that  $\text{cl}_X U_x$  has  $\mathcal{P}$ .

A topological property  $\mathcal{P}$  is said to be *hereditary with respect to closed subsets*, if each closed subset of a space with  $\mathcal{P}$  also has  $\mathcal{P}$ . A topological property  $\mathcal{P}$  is said to be *preserved under finite (closed) sums of subspaces*, if a Hausdorff space has  $\mathcal{P}$ , provided that it is the union of a finite collection of its (closed)  $\mathcal{P}$ -subspaces.

Let  $(P, \leq)$  and  $(Q, \leq)$  be two partially ordered sets. A mapping  $f : (P, \leq) \rightarrow (Q, \leq)$  is said to be an *order-homomorphism* (*anti-order-homomorphism*, respectively), if  $f(a) \leq f(b)$  ( $f(b) \leq f(a)$ , respectively) whenever  $a \leq b$ . An order-homomorphism (anti-order-homomorphism, respectively)  $f : (P, \leq) \rightarrow (Q, \leq)$  is said to be an *order-isomorphism* (*anti-order-isomorphism*, respectively), if  $f^{-1} : (Q, \leq) \rightarrow (P, \leq)$  (exists and) is an order-homomorphism (anti-order-homomorphism, respectively). Two partially ordered sets  $(P, \leq)$  and  $(Q, \leq)$  are called *order-isomorphic* (*anti-order-isomorphic*, respectively), if there exists an order-isomorphism (anti-order-isomorphism, respectively) between them.

## 2. Motivations, notations and definitions

In this article we will be dealing with various sets of one-point extensions of a given topological space  $X$ . For the reader's convenience we list all these sets at the beginning.

**Notation 2.1.** Let  $X$  be a topological space. Denote

- $\mathcal{E}(X) = \{Y : Y \text{ is a one-point Tychonoff extension of } X\}$
- $\mathcal{E}^*(X) = \{Y \in \mathcal{E}(X) : Y \text{ is first-countable at } Y \setminus X\}$
- $\mathcal{E}^C(X) = \{Y \in \mathcal{E}(X) : Y \text{ is Čech-complete}\}$
- $\mathcal{E}^K(X) = \{Y \in \mathcal{E}(X) : Y \text{ is locally compact}\}$

and when  $\mathcal{P}$  is a topological property

- $\mathcal{E}_{\mathcal{P}}(X) = \{Y \in \mathcal{E}(X) : Y \text{ has } \mathcal{P}\}$
- $\mathcal{E}_{local-\mathcal{P}}(X) = \{Y \in \mathcal{E}(X) : Y \text{ is locally-}\mathcal{P}\}$ .

Also, we may use notations which are obtained by combinations of the above notations, e.g.

$$\mathcal{E}_{local-\mathcal{P}}^*(X) = \mathcal{E}^*(X) \cap \mathcal{E}_{local-\mathcal{P}}(X).$$

**Definition 2.2** ([10]). For a Tychonoff space  $X$  and a topological property  $\mathcal{P}$ , let

$$\lambda_{\mathcal{P}}X = \bigcup \{ \text{int}_{\beta X} \text{cl}_{\beta X} C : C \in \text{Coz}(X) \text{ and } \text{cl}_X C \text{ has } \mathcal{P} \}.$$

**Definition 2.3** ([14]). We say that a topological property  $\mathcal{P}$  satisfies *Mrówka's condition* (W), if it satisfies the following: If  $X$  is a Tychonoff space in which there exists a point  $p$  with an open base  $\mathcal{B}$  for  $X$  at  $p$  such that  $X \setminus B$  has  $\mathcal{P}$ , for each  $B \in \mathcal{B}$ , then  $X$  has  $\mathcal{P}$ .

Mrówka's condition (W) is satisfied by a large number of topological properties; among them are (regularity +) the Lindelöf property, paracompactness, metacompactness, subparacompactness, the para-Lindelöf property, the  $\sigma$ -para-Lindelöf property, weak  $\theta$ -refinability,  $\theta$ -refinability (or submetacompactness), weak  $\delta\theta$ -refinability,  $\delta\theta$ -refinability (or the submeta-Lindelöf property), countable paracompactness,  $[\theta, \kappa]$ -compactness,  $\kappa$ -boundedness, screenability,  $\sigma$ -metacompactness, Dieudonné completeness,  $N$ -compactness [15], realcompactness, almost realcompactness [4] and zero-dimensionality (see [10], [12] and [13] for proofs and [2], [17] and [18] for definitions).

In [11] we have obtained the following result.

**Theorem 2.4** ([11]). *Let  $X$  and  $Y$  be locally compact locally- $\mathcal{P}$  non- $\mathcal{P}$  spaces where  $\mathcal{P}$  is either pseudocompactness or a closed hereditary topological property which is preserved under finite closed sums of subspaces and satisfies Mrówka's condition (W). Then, the following are equivalent:*

- (1)  $\lambda_{\mathcal{P}}X \setminus X$  and  $\lambda_{\mathcal{P}}Y \setminus Y$  are homeomorphic.
- (2)  $(\mathcal{E}_{\mathcal{P}}(X), \leq)$  and  $(\mathcal{E}_{\mathcal{P}}(Y), \leq)$  are order-isomorphic.
- (3)  $(\mathcal{E}_{\mathcal{P}}^C(X), \leq)$  and  $(\mathcal{E}_{\mathcal{P}}^C(Y), \leq)$  are order-isomorphic.
- (4)  $(\mathcal{E}_{\mathcal{P}}^K(X), \leq)$  and  $(\mathcal{E}_{\mathcal{P}}^K(Y), \leq)$  are order-isomorphic, provided that  $X$  and  $Y$  are moreover strongly zero-dimensional.

There are topological properties, however, which do not satisfy the assumption of Theorem 2.4 ( $\sigma$ -compactness, for example, does not satisfy Mrówka's condition (W); see [10]). The purpose of this article is to prove the following version of Theorem 2.4. Specific topological properties  $\mathcal{P}$  which satisfy the requirements of Theorem 2.5 below are  $\sigma$ -compactness, the Lindelöf property and the linearly Lindelöf property. Note that in Theorem 3.19 of [9] we have shown that conditions (1) and (3) of Theorem 2.5 are equivalent, if  $\mathcal{P}$  is  $\sigma$ -compactness, and in Theorem 3.21 of [9] we have shown that conditions (1) and (2) of Theorem 2.5 are equivalent, if  $\mathcal{P}$  is the Lindelöf property. Thus, in some sense, Theorem 2.5 generalizes Theorems 3.19 and 3.21 of [9], and at the same time, brings them under a same umbrella.

**Theorem 2.5.** *Let  $X$  and  $Y$  be locally compact paracompact spaces and let  $\mathcal{P}$  be a closed hereditary topological property of compact spaces which is preserved under finite sums of subspaces and coincides with  $\sigma$ -compactness in the realm of locally compact paracompact spaces. Then, the following are equivalent:*

- (1)  $\lambda_{\mathcal{P}}X \setminus X$  and  $\lambda_{\mathcal{P}}Y \setminus Y$  are homeomorphic.
- (2)  $(\mathcal{E}_{\mathcal{P}}^C(X), \leq)$  and  $(\mathcal{E}_{\mathcal{P}}^C(Y), \leq)$  are order-isomorphic.
- (3)  $(\mathcal{E}_{local-\mathcal{P}}^*(X), \leq)$  and  $(\mathcal{E}_{local-\mathcal{P}}^*(Y), \leq)$  are order-isomorphic.

We now introduce some notation which will be widely used in this article.

**Notation 2.6.** Let  $X$  be a Tychonoff space  $X$ . For a subset  $A$  of  $X$  denote

$$A^* = \text{cl}_{\beta X} A \setminus X.$$

In particular,  $X^* = \beta X \setminus X$ .

**Remark 2.7.** Note that the notation given in Notation 2.6 can be ambiguous, as  $A^*$  can mean either  $\beta A \setminus A$  or  $\text{cl}_{\beta X} A \setminus X$ . However, since for  $C^*$ -embedded subsets these two notions coincide, this will not cause any confusion.

**Definition 2.8** ([7]). For a Tychonoff space  $X$ , let

$$\sigma X = \bigcup \{ \text{cl}_{\beta X} H : H \subseteq X \text{ is } \sigma\text{-compact} \},$$

**Notation 2.9.** Let  $X$  be a locally compact paracompact non-compact space. Then,  $X$  can be represented as

$$X = \bigoplus_{i \in I} X_i$$

for some index set  $I$ , with each  $X_i$ , for  $i \in I$ , being  $\sigma$ -compact and non-compact (see Theorem 5.1.27 and Exercise 3.8.C of [3]). For  $J \subseteq I$  denote

$$X_J = \bigcup_{i \in J} X_i.$$

Thus, using the notation of 2.6, we have

$$X_J^* = \text{cl}_{\beta X} \left( \bigcup_{i \in J} X_i \right) \setminus X.$$

**Remark 2.10.** Note that in Notation 2.9 the set  $X_J^*$  is homeomorphic to  $\beta X_J \setminus X_J$ , as  $\text{cl}_{\beta X} X_J$  is homeomorphic to  $\beta X_J$  (see Corollary 3.6.8 of [3]). Thus, when  $J$  is countable (since  $X_J$  is  $\sigma$ -compact and locally compact)  $X_J^*$  is a zero-sets in  $\text{cl}_{\beta X} X_J$  (see 1B of [19]). But,  $\text{cl}_{\beta X} X_J$  is clopen in  $\beta X$ , as  $X_J$  is clopen in  $X$  (see Corollary 3.6.5 of [3]) therefore,  $X_J^*$  is a zero-set in  $\beta X$ . Also, note that with the notation given in 2.9, we have

$$\sigma X = \bigcup \{ \text{cl}_{\beta X} X_J : J \subseteq I \text{ is countable} \}.$$

Note that  $\sigma X$  is open in  $\beta X$  and it contains  $X$ .

### 3. Partially ordered set of one-point extensions as related to topologies of subspaces of outgrowth

In Lemma 3.5 we establish a connection between one-point Tychonoff extensions of a given space  $X$  and compact non-empty subsets of its outgrowth  $X^*$ . Lemma 3.5 (and its preceding lemmas) is known (see e.g. [12]). It is included here for the sake of completeness.

**Lemma 3.1.** *Let  $X$  be a Tychonoff space and let  $C$  be a non-empty compact subset of  $X^*$ . Let  $T$  be the space which is obtained from  $\beta X$  by contracting  $C$  to a point  $p$ . Then, the subspace  $Y = X \cup \{p\}$  of  $T$  is Tychonoff and  $\beta Y = T$ .*

**Proof.** Let  $q : \beta X \rightarrow T$  be the quotient mapping. Note that  $T$  is Hausdorff, and thus, being a continuous image of  $\beta X$ , it is compact. Also, note that  $Y$  is dense in  $T$ . Therefore,  $T$  is a compactification of  $Y$ . To show that  $\beta Y = T$ , it suffices to verify that every continuous  $h : Y \rightarrow \mathbf{I}$  is continuously extendable over  $T$ . Let  $h : Y \rightarrow \mathbf{I}$  be continuous. Let  $G : \beta X \rightarrow \mathbf{I}$  continuously extend  $hq|_{(X \cup C)} : X \cup C \rightarrow \mathbf{I}$  (note that  $\beta(X \cup C) = \beta X$ , as  $X \subseteq X \cup C \subseteq \beta X$ , see Corollary 3.6.9 of [3]). Define  $H : T \rightarrow \mathbf{I}$  such that  $H|_{(\beta X \setminus C)} = G|_{(\beta X \setminus C)}$  and  $H(p) = h(p)$ . Then,  $H|_Y = h$ , and since  $Hq = G$  is continuous, the function  $H$  is continuous.

**Notation 3.2.** Let  $X$  be a Tychonoff space and let  $Y \in \mathcal{E}(X)$ . Denote by

$$\tau_Y : \beta X \rightarrow \beta Y$$

the (unique) continuous extension of  $\text{id}_X$ .

**Lemma 3.3.** *Let  $X$  be a Tychonoff space and let  $Y = X \cup \{p\} \in \mathcal{E}(X)$ . Let  $T$  be the space which is obtained from  $\beta X$  by contracting  $\tau_Y^{-1}(p)$  to the point  $p$ , and let  $q : \beta X \rightarrow T$  be the quotient mapping. Then,  $T = \beta Y$  and  $\tau_Y = q$ .*

**Proof.** We need to show that  $Y$  is a subspace of  $T$ . Since  $\beta Y$  is also a compactification of  $X$  and  $\tau_Y|_X = \text{id}_X$ , by Theorem 3.5.7 of [3], we have  $\tau_Y(X^*) = \beta Y \setminus X$ . For an open subset  $W$  of  $\beta Y$ , the set  $q(\tau_Y^{-1}(W))$

is open in  $T$ , as  $q^{-1}(q(\tau_Y^{-1}(W))) = \tau_Y^{-1}(W)$  is open in  $\beta X$ . Therefore,

$$Y \cap W = Y \cap q(\tau_Y^{-1}(W))$$

is open in  $Y$ , when  $Y$  is considered as a subspace of  $T$ . For the converse, note that if  $V$  is open in  $T$ , since

$$Y \cap V = Y \cap (\beta Y \setminus \tau_Y(\beta X \setminus q^{-1}(V)))$$

and  $\tau_Y(\beta X \setminus q^{-1}(V))$  is compact and thus closed in  $\beta Y$ , the set  $Y \cap V$  is open in  $Y$  in its original topology. By Lemma 3.1 we have  $T = \beta Y$ . This also implies that  $\tau_Y = q$ , as  $\tau_Y, q : \beta X \rightarrow \beta Y$  are continuous and coincide with  $\text{id}_X$  on the dense subset  $X$  of  $\beta X$ .

**Lemma 3.4.** *Let  $X$  be a Tychonoff space. Let  $Y_i \in \mathcal{E}(X)$ , for  $i = 1, 2$ , and denote by  $\tau_i = \tau_{Y_i} : \beta X \rightarrow \beta Y_i$  the continuous extension of  $\text{id}_X$ . Then, the following are equivalent:*

- (1)  $Y_1 \leq Y_2$ .
- (2)  $\tau_2^{-1}(Y_2 \setminus X) \subseteq \tau_1^{-1}(Y_1 \setminus X)$ .

**Proof.** Let  $Y_i = X \cup \{p_i\}$ , for  $i = 1, 2$ . (1) *implies* (2). Suppose that (1) holds. By the definition, there exists a continuous  $f : Y_2 \rightarrow Y_1$  such that  $f|X = \text{id}_X$ . Let  $f_\beta : \beta Y_2 \rightarrow \beta Y_1$  continuously extend  $f$ . Note that the continuous functions  $f_\beta \tau_2, \tau_1 : \beta X \rightarrow \beta Y_1$  coincide with  $\text{id}_X$  on the dense subset  $X$  of  $\beta X$ , and thus  $f_\beta \tau_2 = \tau_1$ . Note that  $X$  is dense in  $\beta Y_i$  (for  $i = 1, 2$ ), as it is dense in  $Y_i$ , and therefore,  $\beta Y_i$  is a compactification of  $X$ . Since  $f_\beta|X = \text{id}_X$ , by Theorem 3.5.7 of [3], we have  $f_\beta(\beta Y_2 \setminus X) = \beta Y_1 \setminus X$ , and thus  $f_\beta(p_2) \in \beta Y_1 \setminus X$ . But,  $f_\beta(p_2) = f(p_2)$ , which implies that  $f_\beta(p_2) \in Y_1 \setminus X = \{p_1\}$ . Therefore,

$$\begin{aligned} \tau_2^{-1}(p_2) &\subseteq \tau_2^{-1}(f_\beta^{-1}(f_\beta(p_2))) \\ &= (f_\beta \tau_2)^{-1}(f_\beta(p_2)) = \tau_1^{-1}(f_\beta(p_2)) = \tau_1^{-1}(p_1). \end{aligned}$$

(2) *implies* (1). Suppose that (2) holds. Let  $f : Y_2 \rightarrow Y_1$  be defined such that  $f(p_2) = p_1$  and  $f|X = \text{id}_X$ . We show that  $f$  is continuous, this will show that  $Y_1 \leq Y_2$ . Note that by Lemma 3.3, the space  $\beta Y_2$  is the quotient space of  $\beta X$  which is obtained by contracting  $\tau_2^{-1}(p_2)$  to a point, and  $\tau_2$  is its corresponding quotient mapping. Thus, in particular,  $Y_2$  is the quotient space of  $X \cup \tau_2^{-1}(p_2)$ , and therefore, to show that  $f$  is continuous, it suffices to show that  $f \tau_2|(X \cup \tau_2^{-1}(p_2))$  is continuous. We show this by verifying that  $f \tau_2(t) = \tau_1(t)$ , for each  $t \in X \cup \tau_2^{-1}(p_2)$ . This obviously holds if  $t \in X$ . If  $t \in \tau_2^{-1}(p_2)$ , then  $\tau_2(t) = p_2$ , and



thus  $f\tau_2(t) = p_1$ . But, since  $t \in \tau_2^{-1}(\tau_2(t))$ , we have  $t \in \tau_1^{-1}(p_1)$ , and therefore  $\tau_1(t) = p_1$ . Thus,  $f\tau_2(t) = \tau_1(t)$  in this case as well.

**Lemma 3.5.** *Let  $X$  be a Tychonoff space. Define a function*

$$\Theta : (\mathcal{E}(X), \leq) \rightarrow (\{C \subseteq X^* : C \text{ is compact}\} \setminus \{\emptyset\}, \subseteq)$$

by

$$\Theta(Y) = \tau_Y^{-1}(Y \setminus X),$$

for  $Y \in \mathcal{E}(X)$ . Then,  $\Theta$  is an anti-order-isomorphism.

**Proof.** To show that  $\Theta$  is well-defined, let  $Y \in \mathcal{E}(X)$ . Note that since  $X$  is dense in  $Y$ , the space  $X$  is dense in  $\beta Y$ . Thus,  $\tau_Y : \beta X \rightarrow \beta Y$  is onto, as  $\tau_Y(\beta X)$  is a compact (and therefore closed) subset of  $\beta Y$  and it contains  $X = \tau_Y(X)$ . Thus,  $\tau_Y^{-1}(Y \setminus X) \neq \emptyset$ . Also, since  $\tau_Y|_X = \text{id}_X$  we have  $\tau_Y^{-1}(Y \setminus X) \subseteq X^*$ , and since the singleton  $Y \setminus X$  is closed in  $\beta Y$ , its inverse image  $\tau_Y^{-1}(Y \setminus X)$  is closed in  $\beta X$ , and therefore it is compact. Now, we show that  $\Theta$  is onto, Lemma 3.4 will then complete the proof. Let  $C$  be a non-empty compact subset of  $X^*$ . Let  $T$  be the quotient space of  $\beta X$  which is obtained by contracting  $C$  to a point  $p$ . Consider the subspace  $Y = X \cup \{p\}$  of  $T$ . Then,  $Y \in \mathcal{E}(X)$ , and thus, by Lemma 3.1 we have  $\beta Y = T$ . The quotient mapping  $q : \beta X \rightarrow T$  is identical to  $\tau_Y$ , as it coincides with  $\text{id}_X$  on the dense subset  $X$  of  $\beta X$ . Therefore,

$$\Theta(Y) = \tau_Y^{-1}(p) = q^{-1}(p) = C.$$

**Notation 3.6.** For a Tychonoff space  $X$  denote by

$$\Theta_X : (\mathcal{E}(X), \leq) \rightarrow (\{C \subseteq X^* : C \text{ is compact}\} \setminus \{\emptyset\}, \subseteq)$$

the anti-order-isomorphism defined by

$$\Theta_X(Y) = \tau_Y^{-1}(Y \setminus X),$$

for  $Y \in \mathcal{E}(X)$ .

Lemmas 3.7 and 3.8 below are known results (see [9]).

**Lemma 3.7.** *Let  $X$  be a Tychonoff space. For  $Y \in \mathcal{E}(X)$  the following are equivalent:*

- (1)  $Y \in \mathcal{E}^*(X)$ .
- (2)  $\Theta_X(Y) \in \mathcal{Z}(\beta X)$ .

**Proof.** Let  $Y = X \cup \{p\}$ . (1) *implies* (2). Suppose that (1) holds. Let  $\{V_n : n \in \mathbf{N}\}$  be an open base at  $p$  in  $Y$ . For each  $n \in \mathbf{N}$ , let  $V'_n$  be an open subset of  $\beta Y$  such that  $Y \cap V'_n = V_n$ , and let  $f_n : \beta Y \rightarrow \mathbf{I}$  be continuous and such that  $f_n(p) = 0$  and  $f_n(\beta Y \setminus V'_n) \subseteq \{1\}$ . Let

$$Z = \bigcap_{n=1}^{\infty} Z(f_n) \in \mathcal{Z}(\beta Y).$$

We show that  $Z = \{p\}$ . Obviously,  $p \in Z$ . Let  $t \in Z$  and suppose to the contrary that  $t \neq p$ . Let  $W$  be an open neighborhood of  $p$  in  $\beta Y$  such that  $t \notin \text{cl}_{\beta Y} W$ . Then,  $Y \cap W$  is an open neighborhood of  $p$  in  $Y$ . Let  $k \in \mathbf{N}$  be such that  $V_k \subseteq Y \cap W$ . We have

$$\begin{aligned} t \in Z(f_k) \subseteq V'_k &\subseteq \text{cl}_{\beta Y} V'_k \\ &= \text{cl}_{\beta Y} (Y \cap V'_k) \\ &= \text{cl}_{\beta Y} V_k \subseteq \text{cl}_{\beta Y} (Y \cap W) \subseteq \text{cl}_{\beta Y} W \end{aligned}$$

which is a contradiction. This shows that  $t = p$  and therefore  $Z \subseteq \{p\}$ . Thus,  $\{p\} = Z \in \mathcal{Z}(\beta Y)$ , which implies that  $\tau_Y^{-1}(p) \in \mathcal{Z}(\beta X)$ .

(2) *implies* (1). Suppose that (2) holds. Let  $\tau_Y^{-1}(p) = Z(f)$  where  $f : \beta X \rightarrow \mathbf{I}$  is continuous. Note that by Lemma 3.3 the space  $\beta Y$  is obtained from  $\beta X$  by contracting  $\tau_Y^{-1}(p)$  to  $p$  with  $\tau_Y : \beta X \rightarrow \beta Y$  as the quotient mapping. Then, for each  $n \in \mathbf{N}$ , the set  $\tau_Y(f^{-1}([0, 1/n]))$  is an open neighborhood of  $p$  in  $\beta Y$ . We show that the collection

$$\left\{ Y \cap \tau_Y \left( f^{-1} \left( \left[ 0, \frac{1}{n} \right) \right) \right) : n \in \mathbf{N} \right\}$$

of open neighborhoods of  $p$  in  $Y$  constitutes an open base at  $p$  in  $Y$ . This will show (1). Let  $V$  be an open neighborhood of  $p$  in  $Y$ . Let  $V'$  be an open subset of  $\beta Y$  such that  $Y \cap V' = V$ . Then,  $p \in V'$  and thus

$$\bigcap_{n=1}^{\infty} f^{-1} \left( \left[ 0, \frac{1}{n} \right] \right) = Z(f) = \tau_Y^{-1}(p) \subseteq \tau_Y^{-1}(V').$$

By compactness we have  $f^{-1}([0, 1/k]) \subseteq \tau_Y^{-1}(V')$ , for some  $k \in \mathbf{N}$ . Therefore,

$$\begin{aligned} Y \cap \tau_Y \left( f^{-1} \left( \left[ 0, \frac{1}{k} \right] \right) \right) &\subseteq Y \cap \tau_Y \left( f^{-1} \left( \left[ 0, \frac{1}{k} \right] \right) \right) \\ &\subseteq Y \cap \tau_Y (\tau_Y^{-1}(V')) \subseteq Y \cap V' = V. \end{aligned}$$

**Lemma 3.8.** *Let  $X$  be a locally compact space. For  $Y \in \mathcal{E}(X)$  the following are equivalent:*

- (1)  $Y \in \mathcal{E}^C(X)$ .
- (2)  $\Theta_X(Y) \in \mathcal{Z}(X^*)$ .

**Proof.** Let  $Y = X \cup \{p\}$ . (1) *implies* (2). Suppose that (1) holds. Then,  $Y^*$  is an  $F_\sigma$  in  $\beta Y$ . Let  $Y^* = \bigcup_{n=1}^{\infty} K_n$  where each  $K_n$  is closed in  $\beta Y$ , for  $n \in \mathbf{N}$ . Then,

$$X^* = \tau_Y^{-1}(p) \cup \bigcup_{n=1}^{\infty} K_n$$

(recall that  $\beta Y$  is the quotient space of  $\beta X$  which is obtained by contracting  $\tau_Y^{-1}(p)$  to  $p$  and  $\tau_Y$  is its quotient mapping; see Lemma 3.3). For each  $n \in \mathbf{N}$ , let  $f_n : \beta X \rightarrow \mathbf{I}$  be continuous and such that

$$f_n(\tau_Y^{-1}(p)) = \{0\} \text{ and } f_n(K_n) \subseteq \{1\}.$$

Let  $f = \sum_{n=1}^{\infty} f_n/2^n$ . Then,  $f : \beta X \rightarrow \mathbf{I}$  is continuous and

$$\tau_Y^{-1}(p) = Z(f) \cap X^* \in \mathcal{Z}(X^*).$$

(2) *implies* (1). Suppose that (2) holds. Let  $\tau_Y^{-1}(p) = Z(g)$  where  $g : X^* \rightarrow \mathbf{I}$  is continuous. Then, using Lemma 3.3, we have

$$\begin{aligned} Y^* = X^* \setminus \tau_Y^{-1}(p) &= X^* \setminus Z(g) \\ &= g^{-1}((0, 1]) = \bigcup_{n=1}^{\infty} g^{-1}\left(\left[\frac{1}{n}, 1\right]\right) \end{aligned}$$

and each set  $g^{-1}([1/n, 1])$ , for  $n \in \mathbf{N}$ , being closed in  $X^*$ , is compact (note that since  $X$  is locally compact,  $X^*$  is compact) and thus closed in  $\beta Y$ . Therefore,  $Y^*$  is an  $F_\sigma$  in  $\beta Y$ , that is,  $Y$  is Čech-complete.

Then, the following lemma justifies our requirement on  $\mathcal{P}$  in Theorem 3.16. We simply need  $\lambda_{\mathcal{P}}X$  to have a more familiar structure.

**Lemma 3.9.** *Let  $\mathcal{P}$  be a topological property which is preserved under finite closed sums of subspaces. The following are equivalent:*

- (1) *The topological property  $\mathcal{P}$  coincides with  $\sigma$ -compactness in the realm of locally compact paracompact spaces.*
- (2) *For every locally compact paracompact space  $X$  we have*

$$\lambda_{\mathcal{P}}X = \sigma X.$$

**Proof.** (1) *implies* (2). Suppose that (1) holds. Let  $X$  be a locally compact paracompact space. Assume the notation of 2.9. Let  $J \subseteq I$  be countable. Then,  $X_J$  is  $\sigma$ -compact and thus (since it is also locally compact and paracompact) it has  $\mathcal{P}$ . Note that  $X_J$  is clopen in  $X$  thus it has a clopen closure in  $\beta X$ , therefore

$$\text{cl}_{\beta X} X_J = \text{int}_{\beta X} \text{cl}_{\beta X} X_J \subseteq \lambda_{\mathcal{P}} X$$

that is,  $\sigma X \subseteq \lambda_{\mathcal{P}} X$ . To see the reverse inclusion, let  $C \in \text{Coz}(X)$  be such that  $\text{cl}_X C$  has  $\mathcal{P}$ . Then, (since  $\text{cl}_X C$  being closed in  $X$  is also locally compact and paracompact)  $\text{cl}_X C$  is  $\sigma$ -compact. Therefore,

$$\text{int}_{\beta X} \text{cl}_{\beta X} C \subseteq \text{cl}_{\beta X} C \subseteq \sigma X$$

which shows that  $\lambda_{\mathcal{P}} X \subseteq \sigma X$ . Thus,  $\lambda_{\mathcal{P}} X = \sigma X$ .

(2) *implies* (1). Suppose that (2) holds. Let  $X$  be a locally compact paracompact space. By the assumption we have  $\lambda_{\mathcal{P}} X = \sigma X$ . We verify that  $X$  has  $\mathcal{P}$  if and only if  $X$  is  $\sigma$ -compact. Assume the notation of Notation 2.9. Suppose that  $X$  has  $\mathcal{P}$ . Then,  $\lambda_{\mathcal{P}} X = \beta X$  and thus  $\sigma X = \beta X$ . Now, by compactness, we have

$$\beta X = \text{cl}_{\beta X} X_{J_1} \cup \cdots \cup \text{cl}_{\beta X} X_{J_n},$$

for some  $n \in \mathbf{N}$  and some countable  $J_1, \dots, J_n \subseteq I$ . Therefore,

$$X = X_{J_1} \cup \cdots \cup X_{J_n}$$

is  $\sigma$ -compact. For the converse, suppose that  $X$  is  $\sigma$ -compact. Then,  $\sigma X = \beta X$  and (since  $\lambda_{\mathcal{P}} X = \sigma X$ ) we have  $\beta X = \lambda_{\mathcal{P}} X$ . Thus, by compactness, we have

$$\beta X = \text{int}_{\beta X} \text{cl}_{\beta X} C_1 \cup \cdots \cup \text{int}_{\beta X} \text{cl}_{\beta X} C_n,$$

for some  $n \in \mathbf{N}$  and some  $C_1, \dots, C_n \in \text{Coz}(X)$  such that  $\text{cl}_X C_i$  has  $\mathcal{P}$ , for  $i = 1, \dots, n$ . Now, using our assumption, the space

$$X = \text{cl}_X C_1 \cup \cdots \cup \text{cl}_X C_n$$

being a finite union of its closed  $\mathcal{P}$ -subspaces, has  $\mathcal{P}$ .

**Lemma 3.10.** *Let  $X$  be a locally compact paracompact space and let  $\mathcal{P}$  be a closed hereditary topological property of compact spaces which is preserved under finite sums of subspaces and coincides with  $\sigma$ -compactness in the realm of locally compact paracompact spaces. For  $Y \in \mathcal{E}(X)$  the following are equivalent:*

- (1)  $Y \in \mathcal{E}_{\mathcal{P}}^C(X)$ .
- (2)  $\Theta_X(Y) \in \mathcal{Z}(X^*)$  and  $\beta X \setminus \lambda_{\mathcal{P}} X \subseteq \Theta_X(Y)$ .

Thus, in particular

$$\Theta_X(\mathcal{E}_{\mathcal{P}}^C(X)) = \{Z \in \mathcal{Z}(X^*) : \beta X \setminus \lambda_{\mathcal{P}} X \subseteq Z\} \setminus \{\emptyset\}.$$

**Proof.** Let  $Y = X \cup \{p\}$ . (1) *implies* (2). Suppose that (1) holds. By Lemma 3.8 we have  $\tau_Y^{-1}(p) \in \mathcal{Z}(X^*)$ . Note that by Lemma 3.9 we have  $\lambda_{\mathcal{P}} X = \sigma X$ . Let  $t \in \beta X \setminus \sigma X$  and suppose to the contrary that  $t \notin \tau_Y^{-1}(p)$ . Let  $f : \beta X \rightarrow \mathbf{I}$  be continuous and such that  $f(t) = 0$  and  $f(\tau_Y^{-1}(p)) = \{1\}$ . Since  $\tau_Y(f^{-1}([0, 1/2]))$  is compact, the set

$$T = X \cap f^{-1}\left(\left[0, \frac{1}{2}\right)\right) = Y \cap \tau_Y\left(f^{-1}\left(\left[0, \frac{1}{2}\right)\right)\right)$$

being closed in  $Y$ , has  $\mathcal{P}$ . But,  $T$ , being closed in  $X$ , is locally compact and paracompact, and thus, having  $\mathcal{P}$ , it is  $\sigma$ -compact. Therefore, by definition of  $\sigma X$  we have  $\text{cl}_{\beta X} T \subseteq \sigma X$ . But, since

$$\begin{aligned} t \in f^{-1}\left(\left[0, \frac{1}{2}\right)\right) &\subseteq \text{cl}_{\beta X} f^{-1}\left(\left[0, \frac{1}{2}\right)\right) \\ &= \text{cl}_{\beta X}\left(X \cap f^{-1}\left(\left[0, \frac{1}{2}\right)\right)\right) \\ &\subseteq \text{cl}_{\beta X}\left(X \cap f^{-1}\left(\left[0, \frac{1}{2}\right]\right)\right) = \text{cl}_{\beta X} T \end{aligned}$$

we have  $t \in \sigma X$ , which contradicts the choice of  $t$ . Thus,  $t \in \tau_Y^{-1}(p)$  and therefore  $\beta X \setminus \sigma X \subseteq \tau_Y^{-1}(p)$ .

(2) *implies* (1). Suppose that (2) holds. Note that since  $X$  is locally compact, the set  $X^*$  is closed in (the normal space)  $\beta X$  and thus, since  $\tau_Y^{-1}(p) \in \mathcal{Z}(X^*)$  (using the Tietze-Urysohn Theorem) we have  $\tau_Y^{-1}(p) = Z \cap X^*$ , for some  $Z \in \mathcal{Z}(\beta X)$ . Note that by Lemma 3.9 we have  $\lambda_{\mathcal{P}} X = \sigma X$ . Now, since  $\beta X \setminus \sigma X \subseteq \tau_Y^{-1}(p) \subseteq Z$  we have  $\beta X \setminus Z \subseteq \sigma X$ . Therefore, assuming the notation of 2.9 (since  $\beta X \setminus Z$ , being a cozero-set in  $\beta X$ , is  $\sigma$ -compact) we have

$$\beta X \setminus Z \subseteq \bigcup_{n=1}^{\infty} \text{cl}_{\beta X} X_{J_n} \subseteq \text{cl}_{\beta X} X_J$$

where  $J_1, J_2, \dots \subseteq I$  are countable and  $J = J_1 \cup J_2 \cup \dots$ . But,

$$Y = \tau_Y(Z) \cup (X \setminus Z) \subseteq \tau_Y(Z) \cup X_J$$

and thus we have

$$(3.1) \quad Y = \tau_Y(Z) \cup X_J.$$

Now, since  $X_J$  has  $\mathcal{P}$ , as it is  $\sigma$ -compact (and being closed in  $X$ , it is locally compact and paracompact) and  $\tau_Y(Z)$  has  $\mathcal{P}$ , as it is compact, from (3.1) it follows that the space  $Y$ , being a finite union of its  $\mathcal{P}$ -subspaces, has  $\mathcal{P}$ . The fact that  $Y$  is Čech-complete follows from Lemma 3.8.

The following generalizes Lemma 3.18 of [9].

**Lemma 3.11.** *Let  $X$  be a locally compact paracompact space and let  $\mathcal{P}$  be a closed hereditary topological property of compact spaces which is preserved under finite sums of subspaces and coincides with  $\sigma$ -compactness in the realm of locally compact paracompact spaces. For  $Y \in \mathcal{E}(X)$  the following are equivalent:*

- (1)  $Y \in \mathcal{E}_{local-\mathcal{P}}^*(X)$ .
- (2)  $\Theta_X(Y) \in \mathcal{Z}(\beta X)$  and  $\Theta_X(Y) \subseteq \lambda_{\mathcal{P}}X$ .

Thus, in particular

$$\Theta_X(\mathcal{E}_{local-\mathcal{P}}^*(X)) = \{Z \in \mathcal{Z}(\beta X) : Z \subseteq \lambda_{\mathcal{P}}X \setminus X\} \setminus \{\emptyset\}.$$

**Proof.** Let  $Y = X \cup \{p\}$ . (1) implies (2). Suppose that (1) holds. Since  $Y \in \mathcal{E}^*(X)$ , by Lemma 3.7 we have  $\tau_Y^{-1}(p) \in \mathcal{Z}(\beta X)$ . Let  $\tau_Y^{-1}(p) = Z(f)$ , for some continuous  $f : \beta X \rightarrow \mathbf{I}$ . Since  $Y$  is locally- $\mathcal{P}$ , there exists an open neighborhood  $V$  of  $p$  in  $Y$  such that  $\text{cl}_Y V$  has  $\mathcal{P}$ . Let  $V'$  be an open subset of  $\beta Y$  such that  $Y \cap V' = V$ . Then,  $p \in V'$ , and thus since

$$\bigcap_{n=1}^{\infty} f^{-1}\left(\left[0, \frac{1}{n}\right]\right) = Z(f) = \tau_Y^{-1}(p) \subseteq \tau_Y^{-1}(V')$$

by compactness, we have  $f^{-1}([0, 1/k]) \subseteq \tau_Y^{-1}(V')$ , for some  $k \in \mathbf{N}$ . Now, for each  $n \geq k$ , since

$$\begin{aligned} Y \cap \tau_Y\left(f^{-1}\left(\left[0, \frac{1}{n}\right]\right) \setminus f^{-1}\left(\left[0, \frac{1}{n+1}\right]\right)\right) &\subseteq Y \cap \tau_Y\left(f^{-1}\left(\left[0, \frac{1}{k}\right]\right)\right) \\ &\subseteq Y \cap \tau_Y(\tau_Y^{-1}(V')) \\ &\subseteq Y \cap V' = V \subseteq \text{cl}_Y V \end{aligned}$$

the set

$$\begin{aligned} K_n &= X \cap \left(f^{-1}\left(\left[0, \frac{1}{n}\right]\right) \setminus f^{-1}\left(\left[0, \frac{1}{n+1}\right]\right)\right) \\ &= Y \cap \tau_Y\left(f^{-1}\left(\left[0, \frac{1}{n}\right]\right) \setminus f^{-1}\left(\left[0, \frac{1}{n+1}\right]\right)\right) \end{aligned}$$

being closed in  $\text{cl}_Y V$ , has  $\mathcal{P}$ , and therefore (since being closed in  $X$  it is locally compact and paracompact) it is  $\sigma$ -compact. (It might be helpful to recall that by Lemma 3.3 the space  $\beta Y$  is obtained from  $\beta X$  by contracting  $\tau_Y^{-1}(p)$  to  $p$  with  $\tau_Y$  as its quotient mapping.) Thus, the set

$$X \cap f^{-1}\left(\left[0, \frac{1}{k}\right]\right) = \bigcup_{n=k}^{\infty} K_n$$

is  $\sigma$ -compact, and therefore, by the definition of  $\sigma X$ , we have

$$\text{cl}_{\beta X}\left(X \cap f^{-1}\left(\left[0, \frac{1}{k}\right]\right)\right) \subseteq \sigma X.$$

But,

$$\begin{aligned} Z(f) \subseteq f^{-1}\left(\left[0, \frac{1}{k}\right]\right) &\subseteq \text{cl}_{\beta X} f^{-1}\left(\left[0, \frac{1}{k}\right]\right) \\ &= \text{cl}_{\beta X}\left(X \cap f^{-1}\left(\left[0, \frac{1}{k}\right]\right)\right) \\ &\subseteq \text{cl}_{\beta X}\left(X \cap f^{-1}\left(\left[0, \frac{1}{k}\right]\right)\right) \end{aligned}$$

from which it follows that  $\tau_Y^{-1}(p) \subseteq \sigma X$ . Finally, note that by Lemma 3.9 we have  $\lambda_{\mathcal{P}} X = \sigma X$ .

(2) *implies* (1). Suppose that (2) holds. By Lemma 3.7 we have  $Y \in \mathcal{E}^*(X)$ . Therefore, it suffices to verify that  $Y$  is locally- $\mathcal{P}$ . Also, since by the assumption  $X$  is locally compact, it is locally- $\mathcal{P}$ , as  $\mathcal{P}$  is assumed to be a topological property of compact spaces. Thus, we only need to verify that  $p$  has an open neighborhood in  $Y$  whose closure in  $Y$  has  $\mathcal{P}$ . Let  $g: \beta X \rightarrow \mathbf{I}$  be continuous and such that  $Z(g) = \tau_Y^{-1}(p)$ . Then, since

$$\bigcap_{n=1}^{\infty} g^{-1}\left(\left[0, \frac{1}{n}\right]\right) = Z(g) \subseteq \lambda_{\mathcal{P}} X$$

by compactness (and since  $\lambda_{\mathcal{P}} X$  is open in  $\beta X$ ) we have  $g^{-1}([0, 1/k]) \subseteq \lambda_{\mathcal{P}} X$ , for some  $k \in \mathbf{N}$ . Note that by Lemma 3.9 we have  $\lambda_{\mathcal{P}} X = \sigma X$ . Assume the notation of Notation 2.9. By compactness, we have

$$g^{-1}\left(\left[0, \frac{1}{k}\right]\right) \subseteq \text{cl}_{\beta X} X_{J_1} \cup \cdots \cup \text{cl}_{\beta X} X_{J_n} = \text{cl}_{\beta X} X_J$$

where  $n \in \mathbf{N}$ , the sets  $J_1, \dots, J_n \subseteq I$  are countable and  $J = J_1 \cup \cdots \cup J_n$ . The set  $X \cap g^{-1}([0, 1/k]) \subseteq X_J$ , being closed in the latter ( $\sigma$ -compact

space) is  $\sigma$ -compact, and therefore (since being closed in  $X$ , it is locally compact and paracompact) it has  $\mathcal{P}$ . Let

$$V = Y \cap \tau_Y \left( g^{-1} \left( \left[ 0, \frac{1}{k} \right) \right) \right).$$

Then,  $V$  is an open neighborhood of  $p$  in  $Y$ . We show that  $\text{cl}_Y V$  has  $\mathcal{P}$ . But, this follows, since

$$\begin{aligned} \text{cl}_Y V \subseteq Y \cap \tau_Y \left( g^{-1} \left( \left[ 0, \frac{1}{k} \right] \right) \right) &= \left( X \cap \tau_Y \left( g^{-1} \left( \left[ 0, \frac{1}{k} \right] \right) \right) \right) \cup \{p\} \\ &= \left( X \cap g^{-1} \left( \left[ 0, \frac{1}{k} \right] \right) \right) \cup \{p\} \end{aligned}$$

and the latter, being a finite union of its  $\mathcal{P}$ -subspaces (note that the singleton  $\{p\}$ , being compact, has  $\mathcal{P}$ ) has  $\mathcal{P}$ , and thus, its closed subset  $\text{cl}_Y V$ , also has  $\mathcal{P}$ .

Lemmas 3.12–3.14 are from [8].

**Lemma 3.12.** *Let  $X$  be a locally compact paracompact space. If  $Z \in \mathcal{Z}(\beta X)$  is non-empty, then  $Z \cap \sigma X \neq \emptyset$*

**Proof.** Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence in  $\sigma X$ . Assume the notation of 2.9. Then,  $\{x_n : n \in \mathbf{N}\} \subseteq \text{cl}_{\beta X} X_J$ , for some countable  $J \subseteq I$ . Therefore,  $\{x_n : n \in \mathbf{N}\}$  has a limit point in  $\text{cl}_{\beta X} X_J \subseteq \sigma X$ . Thus,  $\sigma X$  is countably compact, and therefore is pseudocompact, and  $v(\sigma X) = \beta(\sigma X) = \beta X$  (note that the latter equality holds, as  $X \subseteq \sigma X \subseteq \beta X$ ). The result now follows, as for any Tychonoff space  $T$ , any non-empty zero-set of  $vT$  meets  $T$  (see Lemma 5.11 (f) of [16]).

**Lemma 3.13.** *Let  $X$  be a locally compact paracompact space. If  $Z \in \mathcal{Z}(X^*)$  is non-empty, then  $Z \cap \sigma X \neq \emptyset$ .*

**Proof.** Let  $S \in \mathcal{Z}(\beta X)$  be such that  $S \cap X^* = Z$  (which exists, as  $X^*$  is closed in (the normal space)  $\beta X$ , as  $X$  is locally compact, and thus, by the Tietze-Urysohn Theorem, every continuous function from  $X^*$  to  $\mathbf{I}$  is continuously extendible over  $\beta X$ ). By Lemma 3.12 we have  $S \cap \sigma X \neq \emptyset$ . Suppose that  $S \cap (\sigma X \setminus X) = \emptyset$ . Then,  $S \cap \sigma X = X \cap S$ . Assume the notation of 2.9. Let  $J = \{i \in I : X_i \cap S \neq \emptyset\}$ . Then,  $J$  is finite. Note that since  $X_J$  is clopen in  $X$ , it has a clopen closure in  $\beta X$ . Now,

$$T = S \cap (\beta X \setminus \text{cl}_{\beta X} X_J) \in \mathcal{Z}(\beta X)$$



misses  $\sigma X$ , and therefore, by Lemma 3.12 we have  $T = \emptyset$ . But, this is a contradiction, as  $Z = S \cap (\beta X \setminus \sigma X) \subseteq T$ . This shows that

$$Z \cap (\sigma X \setminus X) = S \cap (\sigma X \setminus X) \neq \emptyset.$$

**Lemma 3.14.** *Let  $X$  be a locally compact paracompact space. For  $S, T \in \mathcal{L}(X^*)$ , if  $S \cap \sigma X \subseteq T \cap \sigma X$ , then  $S \subseteq T$ .*

**Proof.** Suppose to the contrary that  $S \setminus T \neq \emptyset$ , let  $s \in S \setminus T$ . Let  $f : \beta X \rightarrow \mathbf{I}$  be continuous and such that  $f(s) = 0$  and  $f(T) \subseteq \{1\}$ . Then,  $Z(f) \cap S$  is non-empty, and thus by Lemma 3.13 it follows that  $Z(f) \cap S \cap \sigma X \neq \emptyset$ . But, this is not possible, as

$$Z(f) \cap S \cap \sigma X \subseteq Z(f) \cap T = \emptyset.$$

The following lemma is from [9].

**Lemma 3.15.** *Let  $X$  and  $Y$  be locally compact spaces. The following are equivalent:*

- (1)  $X^*$  and  $Y^*$  are homeomorphic.
- (2)  $(\mathcal{E}^C(X), \leq)$  and  $(\mathcal{E}^C(Y), \leq)$  are order-isomorphic.

**Proof.** This follows from the fact that in a compact space the order-structure of the set of its all zero-sets (partially ordered with  $\subseteq$ ) determines its topology.

The proof of the following theorem is essentially a combination of the proofs we have given for Theorems 3.19 and 3.21 in [9] with the appropriate usage of the preceding lemmas. The reasonably detailed proof is included here for the reader's convenience.

**Theorem 3.16.** *Let  $X$  and  $Y$  be locally compact paracompact (non-compact) spaces and let  $\mathcal{P}$  be a closed hereditary topological property of compact spaces which is preserved under finite sums of subspaces and coincides with  $\sigma$ -compactness in the realm of locally compact paracompact spaces. Then, the following are equivalent:*

- (1)  $\lambda_{\mathcal{P}} X \setminus X$  and  $\lambda_{\mathcal{P}} Y \setminus Y$  are homeomorphic.
- (2)  $(\mathcal{E}_{\mathcal{P}}^C(X), \leq)$  and  $(\mathcal{E}_{\mathcal{P}}^C(Y), \leq)$  are order-isomorphic.
- (3)  $(\mathcal{E}_{local-\mathcal{P}}^*(X), \leq)$  and  $(\mathcal{E}_{local-\mathcal{P}}^*(Y), \leq)$  are order-isomorphic.

**Proof.** Let

$$X = \bigoplus_{i \in I} X_i \text{ and } Y = \bigoplus_{j \in J} Y_j,$$

for some index sets  $I$  and  $J$  with each  $X_i$  and  $Y_j$ , for  $i \in I$  and  $j \in J$  being  $\sigma$ -compact and non-compact. We will use notation of 2.9 and Remark 2.10 without mentioning. Note that by Lemma 3.9 we have  $\lambda_{\mathcal{P}}X = \sigma X$  and  $\lambda_{\mathcal{P}}Y = \sigma Y$ . Let

$$\omega\sigma X = \sigma X \cup \{\Omega\} \text{ and } \omega\sigma Y = \sigma Y \cup \{\Omega'\}$$

denote the one-point compactifications of  $\sigma X$  and  $\sigma Y$ , respectively.

(1) *implies* (2). Suppose that (1) holds. Suppose that either  $X$  or  $Y$ , say  $X$ , is  $\sigma$ -compact. Then,  $\sigma Y \setminus Y$  is compact, as it is homeomorphic to  $\sigma X \setminus X = X^*$ , and the latter is compact, as  $X$  is locally compact. Thus,

$$\sigma Y \setminus Y = Y_{H_1}^* \cup \dots \cup Y_{H_n}^* = Y_H^*$$

where  $n \in \mathbf{N}$ , the sets  $H_1, \dots, H_n \subseteq J$  are countable and

$$H = H_1 \cup \dots \cup H_n.$$

Now, if there exists some  $u \in J \setminus H$ , then since  $Y_u \cap Y_H = \emptyset$  we have

$$\text{cl}_{\beta Y} Y_u \cap \text{cl}_{\beta Y} Y_H = \emptyset.$$

Therefore,  $\text{cl}_{\beta Y} Y_u \subseteq Y$ , contradicting the fact that  $Y_u$  is non-compact. Thus,  $J = H$  and  $Y$  is  $\sigma$ -compact. Therefore,  $\sigma Y \setminus Y = Y^*$ . Note that by Lemmas 3.8 and 3.10 we have  $\mathcal{E}_{\mathcal{P}}^C(X) = \mathcal{E}^C(X)$  and  $\mathcal{E}_{\mathcal{P}}^C(Y) = \mathcal{E}^C(Y)$ . The result now follows from Lemma 3.15.

Suppose that  $X$  and  $Y$  are non- $\sigma$ -compact. Let  $f : \sigma X \setminus X \rightarrow \sigma Y \setminus Y$  denote a homeomorphism. We define an order-isomorphism

$$\phi : (\Theta_X(\mathcal{E}_{\mathcal{P}}^C(X)), \subseteq) \rightarrow (\Theta_Y(\mathcal{E}_{\mathcal{P}}^C(Y)), \subseteq).$$

Since  $\Theta_X$  and  $\Theta_Y$  are anti-order-isomorphisms, this will prove (2). Let  $D \in \Theta_X(\mathcal{E}_{\mathcal{P}}^C(X))$ . By Lemma 3.10 we have  $D \in \mathcal{Z}(X^*)$  and  $\beta X \setminus \sigma X \subseteq D$ . Since  $X^* \setminus D \subseteq \sigma X$ , being a cozero-set in  $X^*$  is  $\sigma$ -compact, there exists a countable  $G \subseteq I$  such that  $X^* \setminus D \subseteq X_G^*$ . Now, since  $D \cap X_G^* \in \mathcal{Z}(X_G^*)$ , we have

$$f(D \cap X_G^*) \in \mathcal{Z}(f(X_G^*)).$$

Since  $X_G^*$  is open in  $\sigma X \setminus X$ , its homeomorphic image  $f(X_G^*)$  is open in  $\sigma Y \setminus Y$ , and thus, is open in  $Y^*$ . But,  $f(X_G^*)$  is compact, as it is a continuous image of a compact space, and therefore,  $f(X_G^*)$  is clopen in  $Y^*$ . Thus,

$$f(D \cap X_G^*) \cup (Y^* \setminus f(X_G^*)) \in \mathcal{Z}(Y^*).$$

Let

$$\phi(D) = f(D \cap (\sigma X \setminus X)) \cup (\beta Y \setminus \sigma Y).$$

Note that since

$$\begin{aligned} f(D \cap (\sigma X \setminus X)) &= f((D \cap X_G^*) \cup ((\sigma X \setminus X) \setminus X_G^*)) \\ &= f(D \cap X_G^*) \cup ((\sigma Y \setminus Y) \setminus f(X_G^*)) \end{aligned}$$

we have

$$\begin{aligned} \phi(D) &= f(D \cap (\sigma X \setminus X)) \cup (\beta Y \setminus \sigma Y) \\ &= f(D \cap X_G^*) \cup ((\sigma Y \setminus Y) \setminus f(X_G^*)) \cup (\beta Y \setminus \sigma Y) \\ &= f(D \cap X_G^*) \cup (Y^* \setminus f(X_G^*)) \end{aligned}$$

which shows that  $\phi$  is well-defined. The function  $\phi$  is clearly an order-homomorphism. Since  $f^{-1} : \sigma Y \setminus Y \rightarrow \sigma X \setminus X$  also is a homeomorphism, as above, it induces an order-homomorphism

$$\psi : (\Theta_Y(\mathcal{E}_{\mathcal{P}}^C(Y)), \subseteq) \rightarrow (\Theta_X(\mathcal{E}_{\mathcal{P}}^C(X)), \subseteq)$$

which is defined by

$$\psi(D) = f^{-1}(D \cap (\sigma Y \setminus Y)) \cup (\beta X \setminus \sigma X),$$

for  $D \in \Theta_Y(\mathcal{E}_{\mathcal{P}}^C(Y))$ . It is now easy to see that  $\psi = \phi^{-1}$ , which shows that  $\phi$  is an order-isomorphism.

(2) *implies* (1). Suppose that (2) holds. Suppose that either  $X$  or  $Y$ , say  $X$ , is  $\sigma$ -compact (and non-compact). Then,  $\sigma X = \beta X$ , and thus, by Lemmas 3.8 and 3.10, we have  $\mathcal{E}_{\mathcal{P}}^C(X) = \mathcal{E}^C(X)$ . Suppose that  $Y$  is non- $\sigma$ -compact. Note that  $X$ , being paracompact and non-compact, is non-pseudocompact (see Theorems 3.10.21, 5.1.5 and 5.1.20 of [3]) and therefore,  $X^*$  contains at least two elements, as almost compact spaces are pseudocompact (see Problem 5U (1) of [16]; recall that a Tychonoff space  $T$  is called *almost compact* if  $\beta T \setminus T$  has at most one element). Thus, there exist two disjoint non-empty zero-sets of  $X^*$  corresponding to two elements in  $\mathcal{E}^C(X)$  with no common upper bound in  $\mathcal{E}^C(X)$ . But, this is not true, as  $\mathcal{E}^C(X)$  is order-isomorphic to  $\mathcal{E}_{\mathcal{P}}^C(Y)$ , and any two elements in the latter have a common upper bound in  $\mathcal{E}_{\mathcal{P}}^C(Y)$ . (Note that since  $Y$  is non- $\sigma$ -compact, the set  $\beta Y \setminus \sigma Y$  is non-empty, and by Lemma 3.10, the image of any element in  $\mathcal{E}_{\mathcal{P}}^C(Y)$  under  $\Theta_Y$  contains  $\beta Y \setminus \sigma Y$ .) Therefore,  $Y$  also is  $\sigma$ -compact and by Lemmas 3.8 and 3.10, we have  $\mathcal{E}_{\mathcal{P}}^C(Y) = \mathcal{E}^C(Y)$ . Now, since  $\sigma Y = \beta Y$ , the result follows from Lemma 3.15.

Next, suppose that  $X$  and  $Y$  are both non- $\sigma$ -compact. We show that the two compact spaces  $\omega\sigma X \setminus X$  and  $\omega\sigma Y \setminus Y$  are homeomorphic, by showing that their corresponding sets of zero-sets (partially ordered with  $\subseteq$ ) are order-isomorphic. Since  $\Theta_X$  and  $\Theta_Y$  are anti-order-isomorphisms, condition (2) implies the existence of an order-isomorphism

$$\phi : (\Theta_X(\mathcal{E}_P^C(X)), \subseteq) \rightarrow (\Theta_Y(\mathcal{E}_P^C(Y)), \subseteq).$$

We define an order-isomorphism

$$\psi : (\mathcal{Z}(\omega\sigma X \setminus X), \subseteq) \rightarrow (\mathcal{Z}(\omega\sigma Y \setminus Y), \subseteq)$$

as follows. Let  $Z \in \mathcal{Z}(\omega\sigma X \setminus X)$ . Suppose that  $\Omega \in Z$ . Then, since  $(\omega\sigma X \setminus X) \setminus Z$  is a cozero-set in (the compact space)  $\omega\sigma X \setminus X$ , it is  $\sigma$ -compact. Thus,  $(\omega\sigma X \setminus X) \setminus Z \subseteq X_G^*$ , for some countable  $G \subseteq I$ . Since  $X_G^*$  is clopen in  $X^*$ , we have

$$(Z \setminus \{\Omega\}) \cup (\beta X \setminus \sigma X) = (Z \cap X_G^*) \cup (X^* \setminus X_G^*) \in \mathcal{Z}(X^*).$$

In this case, we let

$$\psi(Z) = (\phi((Z \setminus \{\Omega\}) \cup (\beta X \setminus \sigma X)) \setminus (\beta Y \setminus \sigma Y)) \cup \{\Omega'\}.$$

Now, suppose that  $\Omega \notin Z$ . Then,  $Z \subseteq \sigma X \setminus X$ , and therefore  $Z \subseteq X_G^*$ , for some countable  $G \subseteq I$ , and thus, using this, one can write

$$(3.2) \quad Z = X^* \setminus \bigcup_{n=1}^{\infty} Z_n \text{ where } \beta X \setminus \sigma X \subseteq Z_n \in \mathcal{Z}(X^*) \text{ for } n \in \mathbf{N}.$$

In this case, we let

$$\psi(Z) = Y^* \setminus \bigcup_{n=1}^{\infty} \phi(Z_n).$$

We check that  $\psi$  is well-defined. Assume the representation given in (3.2). Since  $Y^* \setminus \phi(Z_n) \subseteq \sigma Y$ , for  $n \in \mathbf{N}$ , there exists a countable  $H \subseteq J$  such that  $Y^* \setminus \phi(Z_n) \subseteq Y_H^*$ , for all  $n \in \mathbf{N}$ .

**Claim.** For  $Z \in \mathcal{Z}(\omega\sigma X \setminus X)$  with  $\Omega \notin Z$  assume the representation given in (3.2). Let  $H \subseteq J$  be countable and such that  $Y^* \setminus \phi(Z_n) \subseteq Y_H^*$ , for all  $n \in \mathbf{N}$ . Let  $A$  be such that  $\phi(A) = Y^* \setminus Y_H^*$ . Then,

$$Y^* \setminus \bigcup_{n=1}^{\infty} \phi(Z_n) = \phi(A \cup Z) \setminus \phi(A).$$

*Proof of the claim.* Suppose that  $y \in Y^*$  and  $y \notin \phi(Z_n)$ , for each  $n \in \mathbf{N}$ . If  $y \notin \phi(A \cup Z) \setminus \phi(A)$ , then since  $y \notin \phi(Z_1) \supseteq \phi(A)$  we have  $y \notin \phi(A \cup Z)$ . Therefore, there exists some  $B \in \mathcal{X}(Y^*)$  containing  $y$  such that  $B \cap \phi(A \cup Z) = \emptyset$  and  $B \cap \phi(Z_n) = \emptyset$ , for  $n \in \mathbf{N}$ . Let  $C$  be such that  $\phi(C) = B \cup \phi(A \cup Z)$ , and let  $S_n$ , for  $n \in \mathbf{N}$ , be such that

$$\begin{aligned} \phi(S_n) &= \phi(C) \cap \phi(Z_n) \\ &= (B \cup \phi(A \cup Z)) \cap \phi(Z_n) \\ &= (B \cap \phi(Z_n)) \cup (\phi(A \cup Z) \cap \phi(Z_n)) = \phi(A \cup Z) \cap \phi(Z_n). \end{aligned}$$

Since  $A \subseteq Z_n$ , as  $\phi(A) \subseteq \phi(Z_n)$  and  $Z \cap Z_n = \emptyset$ , we have  $A \cap Z = \emptyset$ , which implies that

$$(A \cup Z) \cap Z_n = (A \cap Z_n) \cup (Z \cap Z_n) = A,$$

for  $n \in \mathbf{N}$ . Clearly,  $S_n \subseteq (A \cup Z) \cap Z_n$ , as by above  $\phi(S_n) \subseteq \phi(A \cup Z)$  and  $\phi(S_n) \subseteq \phi(Z_n)$ , for  $n \in \mathbf{N}$ . Thus,  $\phi(S_n) \subseteq \phi(A)$ , for  $n \in \mathbf{N}$ . But, since  $\phi(A) \subseteq \phi(Z_n)$ , we have  $\phi(A) \subseteq \phi(S_n)$ , and therefore

$$\phi(C \cap Z_n) \subseteq \phi(C) \cap \phi(Z_n) = \phi(S_n) = \phi(A),$$

for  $n \in \mathbf{N}$ . This implies that  $C \cap Z_n \subseteq A$ , for  $n \in \mathbf{N}$ . Thus,

$$C \setminus Z = C \cap \bigcup_{n=1}^{\infty} Z_n = \bigcup_{n=1}^{\infty} (C \cap Z_n) \subseteq A.$$

Therefore,  $C \subseteq A \cup Z$  and we have  $B \subseteq \phi(C) \subseteq \phi(A \cup Z)$ , which is a contradiction, as  $B \cap \phi(A \cup Z) = \emptyset$ . This shows that

$$Y^* \setminus \bigcup_{n=1}^{\infty} \phi(Z_n) \subseteq \phi(A \cup Z) \setminus \phi(A).$$

Now, suppose that  $y \in \phi(A \cup Z) \setminus \phi(A)$ . Suppose to the contrary that  $y \in \phi(Z_n)$ , for some  $n \in \mathbf{N}$ . Then,

$$y \in \phi(Z_n) \cap \phi(A \cup Z) = \phi(D),$$

for some  $D$ . Clearly,  $D \subseteq Z_n$  and  $D \subseteq A \cup Z$ , as  $\phi(D) \subseteq \phi(Z_n)$  and  $\phi(D) \subseteq \phi(A \cup Z)$ . This implies that

$$D \subseteq Z_n \cap (A \cup Z) = (Z_n \cap A) \cup (Z_n \cap Z) = Z_n \cap A \subseteq A$$

and thus  $y \in \phi(A)$ , as  $\phi(D) \subseteq \phi(A)$ , which is a contradiction. This proves the claim.

Now, suppose that

$$Z = X^* \setminus \bigcup_{n=1}^{\infty} S_n \text{ and } Z = X^* \setminus \bigcup_{n=1}^{\infty} Z_n$$

are two representations for  $Z \in \mathcal{Z}(\omega\sigma X \setminus X)$  with  $\Omega \notin Z$  such that each  $S_n, Z_n \in \mathcal{Z}(X^*)$  contains  $\beta X \setminus \sigma X$ , for  $n \in \mathbf{N}$ . Choose a countable  $H \subseteq J$  such that

$$Y^* \setminus \phi(S_n) \subseteq Y_H^* \text{ and } Y^* \setminus \phi(Z_n) \subseteq Y_H^*,$$

for  $n \in \mathbf{N}$ . Then, by the claim, we have

$$Y^* \setminus \bigcup_{n=1}^{\infty} \phi(S_n) = \phi(A \cup Z) \setminus \phi(A) = Y^* \setminus \bigcup_{n=1}^{\infty} \phi(Z_n)$$

where  $A$  is such that  $\phi(A) = Y^* \setminus Y_H^*$ . This shows that  $\psi$  is well-defined. Next, we show that  $\psi$  is an order-isomorphism. Suppose that  $S, Z \in \mathcal{Z}(\omega\sigma X \setminus X)$  and  $S \subseteq Z$ . We consider the following cases.

**Case 1:** Suppose that  $\Omega \in S$ . Then,  $\Omega \in Z$ , and clearly,

$$\begin{aligned} \psi(S) &= (\phi((S \setminus \{\Omega\}) \cup (\beta X \setminus \sigma X)) \setminus (\beta Y \setminus \sigma Y)) \cup \{\Omega'\} \\ &\subseteq (\phi((Z \setminus \{\Omega\}) \cup (\beta X \setminus \sigma X)) \setminus (\beta Y \setminus \sigma Y)) \cup \{\Omega'\} = \psi(Z). \end{aligned}$$

**Case 2:** Suppose that  $\Omega \notin S$  but  $\Omega \in Z$ . Let

$$E = \phi((Z \setminus \{\Omega\}) \cup (\beta X \setminus \sigma X))$$

and let

$$S = X^* \setminus \bigcup_{n=1}^{\infty} S_n$$

where each  $S_n \in \mathcal{Z}(X^*)$  contains  $\beta X \setminus \sigma X$ , for  $n \in \mathbf{N}$ . Clearly,  $Y^* \setminus E \subseteq \sigma Y$ . Let  $H \subseteq J$  be countable and such that  $Y^* \setminus \phi(S_n) \subseteq Y_H^*$ , for all  $n \in \mathbf{N}$  and  $Y^* \setminus E \subseteq Y_H^*$ . By the claim, we have  $\psi(S) = \phi(A \cup S) \setminus \phi(A)$ , where  $\phi(A) = Y^* \setminus Y_H^*$ . Since  $Y^* \setminus Y_H^* \subseteq E$ , we have

$$A \subseteq (Z \setminus \{\Omega\}) \cup (\beta X \setminus \sigma X).$$

Now,

$$\psi(S) = \phi(A \cup S) \setminus \phi(A) \subseteq \phi(A \cup S) \subseteq \phi((Z \setminus \{\Omega\}) \cup (\beta X \setminus \sigma X))$$

which implies that

$$\psi(S) \subseteq (\phi((Z \setminus \{\Omega\}) \cup (\beta X \setminus \sigma X)) \setminus (\beta Y \setminus \sigma Y)) \cup \{\Omega'\} = \psi(Z).$$

**Case 3:** Suppose that  $\Omega \notin Z$ . Then,  $\Omega \notin S$ . Let

$$S = X^* \setminus \bigcup_{n=1}^{\infty} S_n \text{ and } Z = X^* \setminus \bigcup_{n=1}^{\infty} Z_n$$

where each  $S_n, Z_n \in \mathcal{Z}(X^*)$  contains  $\beta X \setminus \sigma X$ , for  $n \in \mathbf{N}$ . Clearly,

$$S = S \cap Z = \left( X^* \setminus \bigcup_{n=1}^{\infty} S_n \right) \cap \left( X^* \setminus \bigcup_{n=1}^{\infty} Z_n \right) = X^* \setminus \bigcup_{n=1}^{\infty} (S_n \cup Z_n)$$

and thus, since  $\phi(Z_n) \subseteq \phi(S_n \cup Z_n)$ , for  $n \in \mathbf{N}$ , it follows that

$$\psi(S) = Y^* \setminus \bigcup_{n=1}^{\infty} \phi(S_n \cup Z_n) \subseteq Y^* \setminus \bigcup_{n=1}^{\infty} \phi(Z_n) = \psi(Z).$$

Note that since

$$\phi^{-1} : (\Theta_Y(\mathcal{E}_{\mathcal{P}}^C(Y)), \subseteq) \rightarrow (\Theta_X(\mathcal{E}_{\mathcal{P}}^C(X)), \subseteq)$$

also is an order-isomorphism, as above, it induces an order-isomorphism

$$\gamma : (\mathcal{Z}(\omega\sigma Y \setminus Y), \subseteq) \rightarrow (\mathcal{Z}(\omega\sigma X \setminus X), \subseteq)$$

which is easy to see that  $\gamma = \psi^{-1}$ . Therefore,  $\psi$  is an order-isomorphism. It then follows that there exists a homeomorphism  $f : \omega\sigma X \setminus X \rightarrow \omega\sigma Y \setminus Y$  such that  $f(Z) = \psi(Z)$ , for any  $Z \in \mathcal{Z}(\omega\sigma X \setminus X)$ . Now, since for each countable  $G \subseteq I$  we have

$$f(X_G^*) = \psi(X_G^*) \subseteq \sigma Y \setminus Y$$

it follows that  $f(\sigma X \setminus X) = \sigma Y \setminus Y$ . Thus,  $\sigma X \setminus X$  and  $\sigma Y \setminus Y$  are homeomorphic.

(1) *implies* (3). Suppose that (1) holds. Suppose that either  $X$  or  $Y$ , say  $X$ , is  $\sigma$ -compact. Then,  $\sigma X = \beta X$  and thus, arguing as in part (1) $\Rightarrow$ (2), it follows that  $Y$  also is  $\sigma$ -compact. Therefore,  $\sigma Y = \beta Y$ . Note that by Lemmas 3.7 and 3.11 we have  $\mathcal{E}_{local-\mathcal{P}}^*(X) = \mathcal{E}^*(X)$  and since  $X^* \in \mathcal{Z}(\beta X)$  (as  $X$  is  $\sigma$ -compact and locally compact, see 1B of [19]) by Lemmas 3.7 and 3.8 we have  $\mathcal{E}^*(X) = \mathcal{E}^C(X)$ . Thus,  $\mathcal{E}_{local-\mathcal{P}}^*(X) = \mathcal{E}^C(X)$  and similarly  $\mathcal{E}_{local-\mathcal{P}}^*(Y) = \mathcal{E}^C(Y)$ . The result now follows from Lemma 3.15.

Suppose that  $X$  and  $Y$  are non- $\sigma$ -compact. Let  $f : \sigma X \setminus X \rightarrow \sigma Y \setminus Y$  be a homeomorphism. We define an order-isomorphism

$$\phi : (\Theta_X(\mathcal{E}_{local-\mathcal{P}}^*(X)), \subseteq) \rightarrow (\Theta_Y(\mathcal{E}_{local-\mathcal{P}}^*(Y)), \subseteq),$$

as follows. Let  $Z \in \Theta_X(\mathcal{E}_{local-\mathcal{P}}^*(X))$ . By Lemma 3.11 we have  $Z \in \mathcal{Z}(\beta X)$  and  $Z \subseteq \sigma X \setminus X$ . Thus,  $Z \subseteq X_G^*$ , for some countable  $G \subseteq I$ . Now,  $f(Z) \in \mathcal{Z}(\sigma Y \setminus Y)$  and since  $f(Z)$  is compact, as it is a continuous image of a compact space, it follows that  $f(Z) \subseteq Y_H^*$ , for some countable  $H \subseteq J$ . Therefore,  $f(Z) \in \mathcal{Z}(Y_H^*)$  and then  $f(Z) \in \mathcal{Z}(\text{cl}_{\beta Y} Y_H)$ . Since  $\text{cl}_{\beta Y} Y_H$  is clopen in  $\beta Y$  we have  $f(Z) \in \mathcal{Z}(\beta Y)$ . Define

$$\phi(Z) = f(Z).$$

It is obvious that  $\phi$  is an order-homomorphism. If we let

$$\psi : (\Theta_Y(\mathcal{E}_{local-\mathcal{P}}^*(Y)), \subseteq) \rightarrow (\Theta_X(\mathcal{E}_{local-\mathcal{P}}^*(X)), \subseteq)$$

be defined by

$$\psi(Z) = f^{-1}(Z),$$

then  $\psi = \phi^{-1}$  which shows that  $\phi$  is an order-isomorphism.

(3) *implies* (1). Suppose that (3) holds. Suppose that either  $X$  or  $Y$ , say  $X$ , is  $\sigma$ -compact (and non-compact). Then,  $\sigma X = \beta X$ , and thus, by Lemmas 3.7 and 3.11, we have  $\mathcal{E}_{local-\mathcal{P}}^*(X) = \mathcal{E}^*(X)$ . Therefore, since  $X^* \in \mathcal{Z}(\beta X)$  the set  $\mathcal{E}_{local-\mathcal{P}}^*(X)$  has a smallest element (namely, its one-point compactification  $\omega X$ ). Thus,  $\mathcal{E}_{local-\mathcal{P}}^*(Y)$  also has a smallest element; denote this element by  $T$ . Then, for each countable  $H \subseteq J$  we have

$$Y_H^* \in \Theta_Y(\mathcal{E}_{local-\mathcal{P}}^*(Y))$$

and therefore  $\sigma Y \setminus Y \subseteq \Theta_Y(T)$ . By Lemma 3.14 (with  $\Theta_Y(T)$  and  $Y^*$  as the zero-sets in its statement) we have  $Y^* \subseteq \Theta_Y(T)$ . This implies that  $Y^* \in \mathcal{Z}(\beta Y)$  which shows that  $Y$  is  $\sigma$ -compact. Thus,  $\sigma Y = \beta Y$ , and by Lemmas 3.7 and 3.11, we have  $\mathcal{E}_{local-\mathcal{P}}^*(Y) = \mathcal{E}^*(Y)$ . Therefore, in this case (and since by Lemmas 3.7 and 3.8 we have  $\mathcal{E}^*(X) = \mathcal{E}^C(X)$  and  $\mathcal{E}^*(Y) = \mathcal{E}^C(Y)$ ) the result follows from Lemma 3.15.

Next, suppose that  $X$  and  $Y$  are both non- $\sigma$ -compact. Since  $\Theta_X$  and  $\Theta_Y$  are both anti-order-isomorphisms, there exists an order-isomorphism

$$\phi : (\Theta_X(\mathcal{E}_{local-\mathcal{P}}^*(X)), \subseteq) \rightarrow (\Theta_Y(\mathcal{E}_{local-\mathcal{P}}^*(Y)), \subseteq).$$

We extend  $\phi$  by letting  $\phi(\emptyset) = \emptyset$ . We define a function

$$\psi : (\mathcal{Z}(\omega \sigma X \setminus X), \subseteq) \rightarrow (\mathcal{Z}(\omega \sigma Y \setminus Y), \subseteq)$$

and verify that it is an order-isomorphism. Let  $Z \in \mathcal{Z}(\omega \sigma X \setminus X)$  with  $\Omega \notin Z$ . Since  $Z \subseteq X_G^*$ , for some countable  $G \subseteq I$ , we have  $Z \in \mathcal{Z}(\beta X)$ , and therefore,

$$Z \in \Theta_X(\mathcal{E}_{local-\mathcal{P}}^*(X)) \cup \{\emptyset\}.$$



In this case, let

$$\psi(Z) = \phi(Z).$$

Now, suppose that  $Z \in \mathcal{Z}(\omega\sigma X \setminus X)$  and  $\Omega \in Z$ . Then,  $(\omega\sigma X \setminus X) \setminus Z$  is a cozero-set in  $\omega\sigma X \setminus X$ , and we have

$$(3.3) \quad Z = (\omega\sigma X \setminus X) \setminus \bigcup_{n=1}^{\infty} Z_n \text{ where } Z_n \in \mathcal{Z}(\omega\sigma X \setminus X) \text{ for } n \in \mathbf{N}.$$

Thus, as above, it follows that

$$Z_n \in \Theta_X(\mathcal{E}_{local-\mathcal{P}}^*(X)) \cup \{\emptyset\},$$

for  $n \in \mathbf{N}$ . We verify that

$$(3.4) \quad \bigcup_{n=1}^{\infty} \phi(Z_n) \in \text{Coz}(\omega\sigma Y \setminus Y).$$

To show this, note that since  $\phi(Z_n) \subseteq \sigma Y \setminus Y$  there exists a countable  $H \subseteq J$  such that  $\phi(Z_n) \subseteq Y_H^*$ , for  $n \in \mathbf{N}$ .

**Claim.** For  $Z \in \mathcal{Z}(\omega\sigma X \setminus X)$  with  $\Omega \in Z$  assume the representation given in (3.3). Let  $H \subseteq J$  be countable and such that  $\phi(Z_n) \subseteq Y_H^*$ , for all  $n \in \mathbf{N}$ . Let  $A$  be such that  $\phi(A) = Y_H^*$ . Then,

$$\phi(A \cap Z) = \phi(A) \setminus \bigcup_{n=1}^{\infty} \phi(Z_n).$$

*Proof of the claim.* For each  $n \in \mathbf{N}$ , since  $A \cap Z \cap Z_n = \emptyset$ , we have  $\phi(A \cap Z) \cap \phi(Z_n) = \emptyset$ , as otherwise,  $\phi(A \cap Z)$  and  $\phi(Z_n)$  will have a common lower bound in  $\Theta_Y(\mathcal{E}_{local-\mathcal{P}}^*(Y))$ , that is,  $\phi(A \cap Z) \cap \phi(Z_n)$ , whereas  $A \cap Z$  and  $Z_n$  do not have. Also,  $\phi(A \cap Z) \subseteq \phi(A)$ . Therefore,

$$\phi(A \cap Z) \subseteq \phi(A) \setminus \bigcup_{n=1}^{\infty} \phi(Z_n).$$

To show the reverse inclusion, let  $y \in \phi(A)$  be such that  $y \notin \phi(Z_n)$ , for  $n \in \mathbf{N}$ . There exists  $B \in \mathcal{Z}(\beta Y)$  such that  $y \in B$  and  $B \cap \phi(Z_n) = \emptyset$ , for all  $n \in \mathbf{N}$ . If  $y \notin \phi(A \cap Z)$ , then there exists some  $C \in \mathcal{Z}(\beta Y)$  such that  $y \in C$  and  $C \cap \phi(A \cap Z) = \emptyset$ . Let  $D = \phi(A) \cap B \cap C$  and let  $E$  be such that  $\phi(E) = D$ . For each  $n \in \mathbf{N}$ , since  $\phi(E) \cap \phi(Z_n) = \emptyset$ , we have  $E \cap Z_n = \emptyset$ , and thus  $E \subseteq Z$ . On the other hand, since  $\phi(E) \subseteq \phi(A)$  we have  $E \subseteq A$ , and therefore  $E \subseteq A \cap Z$ . Thus,  $\phi(E) \subseteq \phi(A \cap Z)$ , which implies that  $\phi(E) = \emptyset$ , as  $\phi(E) \subseteq C$ . This contradiction shows that  $y \in \phi(A \cap Z)$ , which proves the claim.

Let  $A$  be such that  $\phi(A) = Y_H^*$ . Now,  $\phi(A \cap Z) \in \mathcal{Z}(\omega\sigma Y \setminus Y)$ , as  $\phi(A \cap Z) \subseteq \phi(A)$ . By the claim we have

$$\begin{aligned} (\omega\sigma Y \setminus Y) \setminus \bigcup_{n=1}^{\infty} \phi(Z_n) &= \left( \phi(A) \setminus \bigcup_{n=1}^{\infty} \phi(Z_n) \right) \cup ((\omega\sigma Y \setminus Y) \setminus \phi(A)) \\ &= \phi(A \cap Z) \cup ((\omega\sigma Y \setminus Y) \setminus \phi(A)) \in \mathcal{Z}(\omega\sigma Y \setminus Y) \end{aligned}$$

and (3.4) is verified. In this case, we let

$$\psi(Z) = (\omega\sigma Y \setminus Y) \setminus \bigcup_{n=1}^{\infty} \phi(Z_n).$$

Next, we show that  $\psi$  is well-defined. Assume that

$$Z = (\omega\sigma X \setminus X) \setminus \bigcup_{n=1}^{\infty} S_n$$

with  $S_n \in \mathcal{Z}(\omega\sigma X \setminus X)$ , for  $n \in \mathbf{N}$ , is another representation of  $Z$ . We need to show that

$$(3.5) \quad \bigcup_{n=1}^{\infty} \phi(Z_n) = \bigcup_{n=1}^{\infty} \phi(S_n).$$

Without any loss of generality, suppose to the contrary that there exists some  $m \in \mathbf{N}$  and  $y \in \phi(Z_m)$  such that  $y \notin \phi(S_n)$ , for all  $n \in \mathbf{N}$ . Then, there exists some  $A \in \mathcal{Z}(\beta Y)$  such that  $y \in A$  and  $A \cap \phi(S_n) = \emptyset$ , for  $n \in \mathbf{N}$ . Consider

$$A \cap \phi(Z_m) \in \Theta_Y(\mathcal{E}_{local-p}^*(Y)).$$

Let  $B$  be such that  $\phi(B) = A \cap \phi(Z_m)$ . Since  $\phi(B) \subseteq A$  we have  $\phi(B) \cap \phi(S_n) = \emptyset$  from which it follows that  $B \cap S_n = \emptyset$ , for  $n \in \mathbf{N}$ . But,  $B \subseteq Z_m$ , as  $\phi(B) \subseteq \phi(Z_m)$ , and we have

$$B \subseteq \bigcup_{n=1}^{\infty} Z_n = \bigcup_{n=1}^{\infty} S_n$$

which implies that  $B = \emptyset$ . But, this is a contradiction, as  $\phi(B) \neq \emptyset$ . Therefore, (3.5) holds, and thus  $\psi$  is well-defined. To prove that  $\psi$  is an order-isomorphism, let  $S, Z \in \mathcal{Z}(\omega\sigma X \setminus X)$  and  $S \subseteq Z$ . The case when  $S = \emptyset$  holds trivially. Assume that  $S \neq \emptyset$ . We consider the following cases.

**Case 1:** Suppose that  $\Omega \notin Z$ . Then,  $\Omega \notin S$  and we have

$$\psi(S) = \phi(S) \subseteq \phi(Z) = \psi(Z).$$

**Case 2:** Suppose that  $\Omega \notin S$  but  $\Omega \in Z$ . Let

$$Z = (\omega\sigma X \setminus X) \setminus \bigcup_{n=1}^{\infty} Z_n$$

with  $Z_n \in \mathcal{Z}(\omega\sigma X \setminus X)$ , for  $n \in \mathbf{N}$ . Then, since  $S \subseteq Z$  we have  $S \cap Z_n = \emptyset$ , and therefore  $\phi(S) \cap \phi(Z_n) = \emptyset$ , for  $n \in \mathbf{N}$ . Thus,

$$\psi(S) = \phi(S) \subseteq (\omega\sigma Y \setminus Y) \setminus \bigcup_{n=1}^{\infty} \phi(Z_n) = \psi(Z).$$

**Case 3:** Suppose that  $\Omega \in S$ . Then,  $\Omega \in Z$ . Let

$$S = (\omega\sigma X \setminus X) \setminus \bigcup_{n=1}^{\infty} S_n \text{ and } Z = (\omega\sigma X \setminus X) \setminus \bigcup_{n=1}^{\infty} Z_n$$

where  $S_n, Z_n \in \mathcal{Z}(\omega\sigma X \setminus X)$ , for  $n \in \mathbf{N}$ . Therefore,

$$\begin{aligned} S = S \cap Z &= \left( (\omega\sigma X \setminus X) \setminus \bigcup_{n=1}^{\infty} S_n \right) \cap \left( (\omega\sigma X \setminus X) \setminus \bigcup_{n=1}^{\infty} Z_n \right) \\ &= (\omega\sigma X \setminus X) \setminus \bigcup_{n=1}^{\infty} (S_n \cup Z_n). \end{aligned}$$

Thus, since  $\phi(Z_n) \subseteq \phi(S_n \cup Z_n)$ , for  $n \in \mathbf{N}$ , we have

$$\psi(S) = (\omega\sigma Y \setminus Y) \setminus \bigcup_{n=1}^{\infty} \phi(S_n \cup Z_n) \subseteq (\omega\sigma Y \setminus Y) \setminus \bigcup_{n=1}^{\infty} \phi(Z_n) = \psi(Z).$$

This shows that  $\psi$  is an order-homomorphism. To show that  $\psi$  is an order-isomorphism, we note that

$$\phi^{-1} : (\Theta_Y(\mathcal{E}_{local-\mathcal{P}}^*(Y)), \subseteq) \rightarrow (\Theta_X(\mathcal{E}_{local-\mathcal{P}}^*(X)), \subseteq)$$

is an order-isomorphism. Let

$$\gamma : (\mathcal{Z}(\omega\sigma Y \setminus Y), \subseteq) \rightarrow (\mathcal{Z}(\omega\sigma X \setminus X), \subseteq)$$

be the induced order-homomorphism which is defined as above. Then, it is straightforward to see that  $\gamma = \psi^{-1}$ , that is,  $\psi$  is an order-isomorphism. This implies the existence of a homeomorphism  $f : \omega\sigma X \setminus X \rightarrow \omega\sigma Y \setminus Y$

such that  $f(Z) = \psi(Z)$ , for every  $Z \in \mathcal{Z}(\omega\sigma X \setminus X)$ . Therefore, for any countable  $G \subseteq I$ , since  $X_G^* \in \mathcal{Z}(\omega\sigma X \setminus X)$ , we have

$$f(X_G^*) = \psi(X_G^*) = \phi(X_G^*) \subseteq \sigma Y \setminus Y.$$

Thus,  $f(\sigma X \setminus X) \subseteq \sigma Y \setminus Y$ , which shows that  $f(\Omega) = \Omega'$ . Therefore,  $\sigma X \setminus X$  and  $\sigma Y \setminus Y$  are homeomorphic.

**Example 3.17.** The Lindelöf property and the linearly Lindelöf property (besides  $\sigma$ -compactness itself) are examples of topological properties  $\mathcal{P}$  satisfying the assumption of Theorem 3.16. To see this, let  $X$  be a locally compact paracompact space. Assume a representation for  $X$  as in Notation 2.9. Recall that a Hausdorff space  $X$  is said to be *linearly Lindelöf* [6] provided that every linearly ordered (by set inclusion  $\subseteq$ ) open cover of  $X$  has a countable subcover, equivalently, if every uncountable subset of  $X$  has a complete accumulation point in  $X$ . (Recall that a point  $x \in X$  is called a *complete accumulation point* of a set  $A \subseteq X$  if for every neighborhood  $U$  of  $x$  in  $X$  we have  $|U \cap A| = |A|$ .) Note that if  $X$  is non- $\sigma$ -compact, then (using the notation of Notation 2.9) the set  $I$  is uncountable. Let  $A = \{x_i : i \in I\}$  where  $x_i \in X_i$ , for  $i \in I$ . Then,  $A$  is an uncountable subset of  $X$  without (even) accumulation points. Thus,  $X$  cannot be linearly Lindelöf as well. For the converse, note that if  $X$  is not linearly Lindelöf, then, obviously,  $X$  is not Lindelöf, and therefore, is non- $\sigma$ -compact, as it is well-known that  $\sigma$ -compactness and the Lindelöf property coincide in the realm of locally compact paracompact spaces (this fact is evident from the representation given for  $X$  in Notation 2.9).

Theorem 3.16 above might leave the impression that  $(\mathcal{E}_{\mathcal{P}}^C(X), \leq)$  and  $(\mathcal{E}_{local-\mathcal{P}}^*(X), \leq)$  are order-isomorphic. The following is to settle this, showing that in most cases this is indeed not going to be the case.

**Theorem 3.18.** *Let  $X$  be a locally compact paracompact (non-compact) space and let  $\mathcal{P}$  be a closed hereditary topological property of compact spaces which is preserved under finite sums of subspaces and coincides with  $\sigma$ -compactness in the realm of locally compact paracompact spaces. Then, the following are equivalent:*

- (1)  $X$  is  $\sigma$ -compact.
- (2)  $(\mathcal{E}_{\mathcal{P}}^C(X), \leq)$  and  $(\mathcal{E}_{local-\mathcal{P}}^*(X), \leq)$  are order-isomorphic.

**Proof.** Since  $X$  is locally compact, the set  $X^*$  is closed in (the normal space)  $\beta X$  and thus, using the Tietze-Urysohn Theorem, every zero-set of  $X^*$  is extendible to a zero-set of  $\beta X$ . Now, if  $X$  is  $\sigma$ -compact (since  $X$  is also locally compact) we have  $X^* \in \mathcal{Z}(\beta X)$  and therefore every zero-set of  $X^*$  is a zero-set of  $\beta X$ . Note that  $\lambda_{\mathcal{P}}X = \sigma X = \beta X$ . Thus, using Lemmas 3.10 and 3.11 we have

$$\Theta_X(\mathcal{E}_{\mathcal{P}}^C(X)) = \mathcal{Z}(X^*) \setminus \{\emptyset\} = \Theta_X(\mathcal{E}_{local-\mathcal{P}}^*(X))$$

from which it follows that

$$\mathcal{E}_{\mathcal{P}}^C(X) = \mathcal{E}_{local-\mathcal{P}}^*(X).$$

If  $X$  is non- $\sigma$ -compact, then any two elements of  $\mathcal{E}_{\mathcal{P}}^C(X)$  have a common upper bound while this is not the case for  $\mathcal{E}_{local-\mathcal{P}}^*(X)$ . To see this, note that by Lemma 3.10 the set  $\Theta_X(\mathcal{E}_{\mathcal{P}}^C(X))$  is closed under finite intersections (note that the finite intersections are non-empty, as they contain  $\beta X \setminus \sigma X$  and the latter is non-empty, as  $X$  is non- $\sigma$ -compact) while there exist (at least) two elements in  $\Theta_X(\mathcal{E}_{local-\mathcal{P}}^*(X))$  with empty intersection; simply consider  $X_i^*$  and  $X_j^*$ , for some distinct  $i, j \in I$  (we are assuming the representation for  $X$  given in Notation 2.9).

**Project 3.19.** Let  $X$  be a (locally compact paracompact) space and let  $\mathcal{P}$  be a (closed hereditary) topological property (of compact spaces which is preserved under finite sums of subspaces and coincides with  $\sigma$ -compactness in the realm of locally compact paracompact spaces). Explore the relationship between the order structures of  $(\mathcal{E}_{\mathcal{P}}^C(X), \leq)$  and  $(\mathcal{E}_{local-\mathcal{P}}^*(X), \leq)$ .

### Acknowledgment

This research was in part supported by a grant from IPM (No. 86540012).

### REFERENCES

- [1] G. Beer, On convergence to infinity, *Monatsh. Math.* **129** (2000) 267-280.
- [2] D. K. Burke, Covering properties, in: K. Kunen and J.E. Vaughan (Eds.), *Handbook of Set-theoretic Topology*, Elsevier, Amsterdam, 1984, pp. 347-422.
- [3] R. Engelking, *General Topology*, Second edition, Heldermann Verlag, Berlin, 1989.
- [4] Z. Frolík, A generalization of real compact spaces, *Czechoslovak Math. J.* **13** (1963) 127-138.

- [5] L. Gillman and M. Jerison, *Rings of Continuous Functions*, Springer-Verlag, New York-Heidelberg, 1976.
- [6] C. Good, The Lindelöf property, in: K. P. Hart, J. Nagata and J. E. Vaughan (Eds.), *Encyclopedia of General Topology*, Elsevier Science Publishers, B.V., Amsterdam, 2004, pp. 182-184.
- [7] M. Henriksen, L. Janos and R. G. Woods, Properties of one-point completions of a noncompact metrizable space, *Comment. Math. Univ. Carolin.* **46** (2005) 105-123.
- [8] M. R. Koushesh, On one-point metrizable extensions of locally compact metrizable spaces, *Topology Appl.* **154** (2007) 698-721.
- [9] M. R. Koushesh, On order structure of the set of one-point Tychonoff extensions of a locally compact space, *Topology Appl.* **154** (2007) 2607-2634.
- [10] M. R. Koushesh, Compactification-like extensions, *Dissertationes Math. (Rozprawy Mat.)* **476** (2011) 88 pp.
- [11] M. R. Koushesh, The partially ordered set of one-point extensions, *Topology Appl.* **158** (2011) 509-532.
- [12] J. Mack, M. Rayburn and G. Woods, Local topological properties and one-point extensions, *Canad. J. Math.* **24** (1972) 338-348.
- [13] J. Mack, M. Rayburn and G. Woods, Lattices of topological extensions, *Trans. Amer. Math. Soc.* **189** (1974) 163-174.
- [14] S. Mrówka, On local topological properties, *Bull. Acad. Polon. Sci. Cl. III* **5** (1957) 951-956.
- [15] S. Mrówka, Some comments on the author's example of a non- $\mathcal{R}$ -compact space, *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.* **18** (1970) 443-448.
- [16] J. R. Porter and G. Woods, *Extensions and Absolutes of Hausdorff Spaces*, Springer-Verlag, New York, 1988.
- [17] R. M. Stephenson, Jr., Initially  $\kappa$ -Compact and Related Spaces, in: *Handbook of set-theoretic topology*, North-Holland, Amsterdam, 1984, pp. 603-632.
- [18] J. E. Vaughan, Countably compact and sequentially compact spaces, in: *Handbook of set-theoretic topology*, North-Holland, Amsterdam, 1984, pp. 569-602.
- [19] R. C. Walker, *The Stone-Ćech Compactification*, Springer-Verlag, New York-Berlin, 1974.

**M. R. Koushesh**

Department of Mathematical Sciences, Isfahan University of Technology, Isfahan 84156-83111, Iran

and

School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box: 19395-5746, Tehran, Iran.

Email: koushesh@cc.iut.ac.ir