

## BEST PROXIMITY PAIR AND COINCIDENCE POINT THEOREMS FOR NONEXPANSIVE SET-VALUED MAPS IN HILBERT SPACES

A. AMINI-HARANDI

Communicated by Nezam Mahdavi-Amiri

ABSTRACT. This paper is concerned with the best proximity pair problem in Hilbert spaces. Given two subsets  $A$  and  $B$  of a Hilbert space  $H$  and the set-valued maps  $F : A \rightarrow 2^B$  and  $G : A_0 \rightarrow 2^{A_0}$ , where  $A_0 = \{x \in A : \|x - y\| = d(A, B) \text{ for some } y \in B\}$ , best proximity pair theorems provide sufficient conditions that ensure the existence of an  $x_0 \in A$  such that

$$d(G(x_0), F(x_0)) = d(A, B).$$

### 1. Introduction

Let  $(M, d)$  be a metric space and let  $A$  and  $B$  be nonempty subsets of  $M$ . Let  $d(A, B) = \inf\{d(a, b) : a \in A, b \in B\}$ . Let

$$B_0 := \{b \in B : d(a, b) = d(A, B) \text{ for some } a \in A\},$$

and

$$A_0 := \{a \in A : d(a, b) = d(A, B) \text{ for some } b \in B\}.$$

Let  $G : A_0 \rightarrow 2^{A_0}$  and  $F : A \rightarrow 2^B$  be set valued maps.  $(G(x_0), F(x_0))$  is called a *best proximity pair*, if  $d(G(x_0), F(x_0)) = d(A, B)$ . Best proximity pair theorems analyse the conditions on  $F$ ,  $G$ ,  $A$  and  $B$  under which the problem of minimizing the real valued function  $x \rightarrow d(G(x), F(x))$

---

MSC(2000): Primary: 47H09; Secondary: 41A65, 46B20.

Keywords: Best proximity pair, coincidence point, nonexpansive map, Hilbert space.

Received: 10 February 2010, Accepted: 27 July 2010.

© 2011 Iranian Mathematical Society.

has a solution. In the setting of normed spaces and hyperconvex metric spaces, the best proximity pair problem has been studied by many authors, see [1, 2, 3, 5, 7, 8].

Let  $H$  be a Hilbert space and  $A, B \subseteq H$ . It is well-known that if  $A$  and  $B$  are compact subsets of  $M$ , then there exist  $a_0 \in A$  and  $b_0 \in B$  such that  $d(A, B) = d(a_0, b_0)$ . Therefore, in this case

$$d(A, B) = 0 \Leftrightarrow A \cap B \neq \emptyset.$$

Let  $M$  be a metric space and let  $\mathcal{M}$  denotes the family of nonempty, closed bounded subsets of  $M$ . Let  $A, B \in \mathcal{M}$ . The Hausdorff metric  $d_H$  on  $\mathcal{M}$  defined by

$$d_H(A, B) = \inf\{\epsilon > 0 : A \subseteq N_\epsilon(B) \text{ and } B \subseteq N_\epsilon(A)\},$$

where  $N_\epsilon(A)$  denotes the closed  $\epsilon$ -neighborhood of  $A$ , that is,  $N_\epsilon(A) = \{x \in M : d(x, A) \leq \epsilon\}$ . Let  $X$  and  $Y$  be topological spaces with  $C \subseteq Y$ . Let  $G : X \rightarrow 2^Y$  be a set-valued map with nonempty values. The inverse image of  $C$  under  $G$  is

$$G^{-}(C) = \{x \in X : G(x) \cap C \neq \emptyset\}.$$

A set-valued map  $F : A \rightarrow 2^B$  is said to be *nonexpansive*, if for each  $x, y \in A$

$$d_H(F(x), F(y)) \leq \|x - y\|.$$

Given a nonempty closed convex subset  $A$  of a Hilbert space  $H$ ,  $P_A$  will always denote the nearest point projection of  $H$  onto  $A$ . We will use the well-known fact that  $P_A$  is nonexpansive and so is continuous.

**Lemma 1.1.** ([5, Lemma 3.1]) *Let  $A$  be a nonempty closed convex subset of a Hilbert space  $H$ . If  $C$  and  $D$  are nonempty closed and bounded subsets of  $H$ , then*

$$d_H(P_A(C), P_A(D)) \leq d_H(C, D).$$

Let  $(X, \|\cdot\|)$  be a reflexive Banach space and  $A \subseteq X$  be nonempty, closed, convex and bounded. It is well-known that for each  $x \in X$ ,  $P_A(x) \neq \emptyset$ . Here we give the proof for the completeness. For each  $n \in \mathbb{N}$ , let  $A_n(x) = \{y \in A : d(x, y) \leq d(x, A) + \frac{1}{n}\}$ . Notice that  $(A_n(x))$  is a decreasing sequence of nonempty closed, convex bounded subsets of the reflexive Banach space  $X$ , so by Šmulina theorem we have [4, page 433]

$$P_A(x) = \bigcap_{n=1}^{\infty} A_n(x) \neq \emptyset.$$

**Lemma 1.2.** ([5, Lemma 3.2]) *Let  $X$  be a reflexive Banach space. Let  $A$  be a nonempty bounded closed convex subset of  $X$ , and let  $B$  be a nonempty closed convex subset of  $X$ . Then,  $A_0$  and  $B_0$  are nonempty and satisfy*

$$P_B(A_0) \subseteq B_0 \text{ and } P_A(B_0) \subseteq A_0.$$

Recall that a Banach space  $X$  is *uniformly convex*, if given  $\epsilon > 0$  there is a  $\delta > 0$  such that whenever  $\|x\| = \|y\| = 1$  and  $\|x - y\| \geq \epsilon$ , then  $\|\frac{x+y}{2}\| \leq 1 - \delta$ .

**Theorem 1.3.** ([6]) *Let  $X$  be a uniformly convex Banach space, Let  $K$  be a bounded, closed and convex subset of  $X$ , and suppose  $F : K \rightarrow 2^K$  is a compact-valued, nonexpansive set-valued map. Then,  $F$  has a fixed point.*

## 2. Main results

We first present a coincidence point theorem for nonexpansive set-valued self maps.

**Theorem 2.1.** *Let  $H$  be a Hilbert space and  $K$  be a closed, bounded convex subset of  $H$ . Let  $F : K \rightarrow 2^K$  be a nonexpansive set-valued map with nonempty compact values. Let  $G : K \rightarrow 2^K$  be an onto, set-valued map for which  $G^-(C)$  is compact for each compact set  $C \subseteq K$ . Assume that for each compact subsets  $C$  and  $D$  of  $K$*

$$d_H(G^-(C), G^-(D)) \leq d_H(C, D).$$

*Then, there exists a  $x_0 \in K$  with*

$$F(x_0) \cap G(x_0) \neq \emptyset.$$

**Proof.** Since

$F(x_0) \cap G(x_0) \neq \emptyset \Leftrightarrow x_0 \in G^-(F(x_0)) = \{x \in H : G(x) \cap F(x_0) \neq \emptyset\}$ , then, the conclusion follows, if we show that the set-valued map  $J(x) = G^-(F(x)) : K \rightarrow 2^K$  has a fixed point. Since  $G$  is onto, then  $J(x) \neq \emptyset$ . For each  $x \in K$ , since  $F(x)$  is compact, then  $J(x) = G^-(F(x))$  is compact. Now, we show that  $J$  is nonexpansive. For each  $x, y \in K$  we have

$$\begin{aligned} d_H(J(x), J(y)) &= d_H(G^-(F(x)), G^-(F(y))) \leq \\ &d_H(F(x), F(y)) \leq \|x - y\|. \end{aligned}$$

Therefore,  $J$  satisfies all conditions of Theorem 1.3 and so has a fixed point.

Now, we obtain a best proximity pair theorem for nonexpansive set-valued maps in Hilbert spaces.

**Theorem 2.2.** *Let  $H$  be a Hilbert space. Let  $A$  be a nonempty bounded closed convex subset of  $H$ , and let  $B$  be a nonempty closed convex subset of  $H$ . Let  $F : A \rightarrow 2^B$  be a nonexpansive set-valued map with nonempty compact values. Let  $G : A_0 \rightarrow 2^{A_0}$  be an onto set-valued map for which  $G^-(C)$  is compact for each compact set  $C \subseteq A_0$ . Assume that for each compact subsets  $C$  and  $D$  of  $A_0$*

$$d_H(G^-(C), G^-(D)) \leq d_H(C, D).$$

*Assume that  $F(A_0) \subseteq B_0$ . Then, there exists a  $x_0 \in A_0$  such that*

$$d(G(x_0), F(x_0)) = d(A, B).$$

**Proof.** By Lemma 1.2,  $A_0$  and  $B_0$  are nonempty. Let us show that  $A_0$  is closed. To this end, let  $x_n \in A_0$  be a convergent sequence, say,  $x_n \rightarrow x_0 \in A$ . Then, for each  $n \in \mathbb{N}$ , there exists  $y_n \in B$  such that  $d(x_n, y_n) = d(A, B)$ . Thus,  $\{y_n\}$  is a bounded sequence in  $B$  (note that  $\{x_n\}$  is bounded). Since bounded subsets of a reflexive Banach space are weakly sequentially compact [4, Theorem 28, page 68], then passing to a subsequence, if necessary, we may assume that  $(y_n)$  converges weakly, say to  $y_0 \in B$ . Since  $\|\cdot\|$  is weakly lower semicontinuous, then we get

$$\|x_0 - y_0\| \leq \liminf_{n \rightarrow \infty} \|x_n - y_n\| = d(A, B).$$

Therefore,  $\|x_0 - y_0\| = d(A, B)$ , and so  $x_0 \in A_0$ . From Lemma 1.2,  $P_A(B_0) \subseteq A_0$  and by Lemma 1.1,

$$d_H(P_A(F(x)), P_A(F(y))) \leq d_H(F(x), F(y)) \leq \|x - y\|.$$

Then, the map  $P_A(F(\cdot)) : A_0 \rightarrow A_0$  is a nonxpansive set-valued map. Moreover,  $A_0$  is a nonempty closed bounded convex subsets of  $H$ , and for each  $x \in A_0$ ,  $P_A(F(x))$  is a compact subset of  $A_0$  (note  $F(x)$  is compact and  $P_A$  is continuous). Hence, by Theorem 2.1 there exists a  $x_0 \in A_0$  such that

$$P_A(F(x_0)) \cap G(x_0) \neq \emptyset.$$

Let  $z_0 \in P_A(F(x_0)) \cap G(x_0)$ , then there exists  $y_0 \in F(x_0)$  so that  $z_0 = P_{A_0}(y_0)$ . Since  $x_0 \in A_0$  and  $y_0 \in F(x_0) \subseteq B_0$ , there exists  $a_0 \in A_0$  such

that  $d(a_0, y_0) = d(A, B)$ . Therefore,

$$d(A, B) \leq d(G(x_0), F(x_0)) \leq d(z_0, F(x_0)) \leq$$

$$d(P_{A_0}(y_0), y_0) \leq d(a_0, y_0) = d(A, B)$$

Thus,

$$d(G(x_0), F(x_0)) = d(A, B).$$

**Remark 2.3.** Let  $A$  be a nonempty bounded, closed convex subset of  $H$ . Let  $G : A_0 \rightarrow A_0$  be an onto isometry. We show that  $G$  satisfies all the conditions of Theorem 2.2. Let  $C$  be a compact subset of  $A_0$ . Since  $G$  is an isometry, then  $G^-(C) = G^{-1}(C)$  is compact (note  $G^{-1}$  is isometry and so is continuous). Since  $G : A_0 \rightarrow A_0$  is isometry, then  $d_H(G^-(C), G^-(D)) = d_H(C, D)$ , for compact subsets  $C$  and  $D$  of  $A_0$ .

If we take  $G = I$ , Theorem 2.2 reduces to Theorem 3.3 of Kirk, Reich and Veeramani [5].

**Theorem 2.4.** Let  $H$  be a Hilbert space. Let  $A$  be a nonempty bounded closed convex subset of  $H$ , and let  $B$  be a nonempty closed convex subset of  $H$ . Let  $F : A \rightarrow 2^B$  be a nonexpansive set-valued map with nonempty compact values. Assume that  $F(A_0) \subseteq B_0$ . Then, there exists a  $x_0 \in A_0$  such that

$$d(x_0, F(x_0)) = d(A, B).$$

### Acknowledgments

This research was in part supported by a grant from IPM (No. 89470016). The author was also partially supported by the Center of Excellence for Mathematics, University of Shahrekord.

### REFERENCES

- [1] M. A. Al-Thagafi and N. Shahzad, Best proximity pairs and equilibrium pairs for Kakutani multimaps, *Nonlinear Anal.*, **70** (2009) 1209-1216.
- [2] A. Amini-Harandi, A. P. Farajzadeh, D. O'Regan and R. P. Agarwal, Best proximity pair theorems for continuous set-valued maps, *Fixed Point Theory Appl.*, Vol. 2008, Art ID 607926.

- [3] A. Amini-Harandi, A. P. Farajzadeh, D. O'Regan and R. P. Agarwal, Coincidence point, best approximation, and best proximity theorems for condensing set-valued maps in hyperconvex metric spaces, *Fixed Point Theory Appl.*, Vol. 2008, Article ID 543154.
- [4] N. Dunford and J. T. Schwartz, *Linear Operators*, Part I, General theory, With the assistance of William G. Bade and Robert G. Bartle, Reprint of the 1958 original, Wiley Classics Library, A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1988
- [5] W. A. Kirk, S. Reich and P. Veeramani, Proximinal retracts and best proximity pair theorems, *Numer. Funct. Anal. Optim.* **24**(7, 8) (2003) 851-862.
- [6] T. C. Lim, A fixed point theorem for multivalued nonexpansive mappings in a uniformly convex Banach space, *Bull. Amer. Math. Soc.* **80** (1974) 1123-1126.
- [7] S. Sadiq Basha and P. Veeramani, Best proximity pair theorems for multifunctions with open fibers, *J. Approx. Theory* **103** (2000) 119-129.
- [8] P. S. Srinivasan and P. Veeramani, On best proximity pair theorems and fixed-point theorems, [ MR1954244 (2004a:47069)], Proceedings of the International Conference on Fixed-Point Theory and its Applications, *Hindawi Publ. Corp., Cairo* (2003) 33-47.

**A. Amini-Harandi**

Department of Mathematics, University of Shahrekord, P.O.Box: 115, Shahrekord, Iran  
and

School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O.Box: 19395-5746, Tehran, Iran

Email: aminih\_a@yahoo.com

Archive of SID