Bulletin of the Iranian Mathematical Society Vol. 39 No. 6 (2013), pp 1273-1281.

# 2-RECOGNIZABILITY OF THE SIMPLE GROUPS $B_n(3)$ AND $C_n(3)$ BY PRIME GRAPH

#### M. FOROUDI GHASEMABADI, A. IRANMANESH\* AND N. AHANJIDEH

Communicated by Ali Reza Ashrafi

ABSTRACT. Let G be a finite group and let GK(G) be the prime graph of G. We assume that  $n \ge 5$ , is an odd number. In the paper, we show that the simple groups  $B_n(3)$  and  $C_n(3)$  are 2-recognizable by their prime graphs. As the result, the characterizability of the groups  $B_n(3)$  and  $C_n(3)$  by their spectra and by the set of orders of maximal abelian subgroups are obtained. Also, we conclude that the AAM's conjecture is true for these groups.

## 1. Introduction

For a finite group G we denote by  $\pi(G)$  the set of all prime divisors of |G| and the spectrum  $\omega(G)$  of G is the set of element orders of G, i.e., a natural number n is in  $\omega(G)$  if there is an element of order n in G. The prime graph (or Gruenberg-Kegel graph) GK(G) of G is the graph with vertex set  $\pi(G)$  where two distinct vertices p and q are adjacent by an edge (briefly, adjacent) if  $pq \in \omega(G)$ , in which case, we write  $(p,q) \in GK(G)$ . Given an arbitrary subset  $\omega$  of the set of natural numbers, denote by  $h(\omega)$  the number of pairwise nonisomorphic finite groups G such that  $\omega(G) = \omega$ . Given a natural number m, a group G is said to be *m*-recognizable by spectrum if  $h(\omega(G)) = m$ . In particular, G is said to be characterizable by spectrum if  $h(\omega(G)) = 1$ . The recognition

MSC(2010): Primary: 20D05; Secondary: 20D06, 20D60.

Keywords: Prime graph, classification of finite simple groups, recognition, spectrum.

Received: 25 May 2011, Accepted: 22 September 2011.

<sup>\*</sup>Corresponding author

 $<sup>\</sup>bigodot$  2013 Iranian Mathematical Society.

<sup>1273</sup> 

problem is solved for a finite group G if we know the value of  $h(\omega(G))$ . Along side the above definitions, we denote by k(GK(G)) the number of isomorphism classes of the finite groups H satisfying GK(G) = GK(H). Hence, if G is a finite group then  $k(GK(G)) \ge 1$ . Given a natural number m, a finite group G is called m-recognizable by prime graph if k(GK(G)) = m. Usually, a 1-recognizable group by prime graph is called a *characterizable group by prime graph*. The recognition problem by prime graph have been considered for most of finite nonabelian simple groups with disconnected prime graphs ([8, 9, 10, 15, 20]), but  $L_{16}(2)$  is the only group with connected prime graph which its recognizability by prime graph has been solved completely so far ([11, 21]). Here, we prove that the simple groups  $B_n(3)$  and  $C_n(3)$ , where n > 3 is an odd number, are 2-recognizable by prime graph and since these groups have connected prime graphs when n is non-prime, thus we obtain the first example of two infinite series of finite simple groups with connected prime graphs which their recognition problem by prime graph are solved thoroughly. In fact, we have the following main theorem:

**Main theorem.** Let n > 3 be an odd number. The simple groups  $B_n(3)$  and  $C_n(3)$  are 2-recognizable by prime graph. In fact if G is a finite group with  $GK(G) = GK(B_n(3))$ , then  $G \cong B_n(3)$  or  $G \cong C_n(3)$ . Also, if G is a finite group such that  $GK(G) = GK(B_3(3))$ , then  $G \cong B_3(3), C_3(3), D_4(3)$ , or  $G/O_2(G) = \operatorname{Aut}({}^2B_2(8))$ .

Note that for an odd prime n, the above groups have disconnected prime graphs and their 2-recognizability by prime graph have been considered in [15]. Since the method we used to prove the main theorem is based on our result in [6], we give a new proof for 2-recognizability of these groups in the disconnected case as well.

In [22, 23], it has been proved that if n > 3 is an odd prime and  $B_n(3)$ and  $C_n(3)$  have disconnected prime graphs, then these groups are characterizable by their spectra. Since the knowledge of  $\omega(G)$  determines GK(G), as the first result of the main theorem, we can conclude that these groups are characterizable by their spectra in the connected case as well.

The non-commuting graph of a nonabelian group G, denoted by  $\Gamma_G$ , is the graph with vertex set  $G \setminus Z(G)$ , where two distinct vertices x and y are connected by an edge if  $xy \neq yx$ . Problem 16.1 in the Kourovka notebook [12] is the AAM's conjecture, which asserts that simple groups are determined uniquely by the non-commuting graph. As prominent corollaries of the main theorem, the validity of the AAM's conjecture and characterizability by the set of orders of maximal abelian subgroups can be obtained for the groups  $B_n(3)$  and  $C_n(3)$ , where  $n \ge 3$  is odd.

### 2. Preliminaries

If q is a natural number, r is an odd prime and gcd(r,q) = 1, then by e(r,q) we denote the smallest natural number m such that  $q^m \equiv 1 \pmod{r}$ . If r = 2, then we put e(2,q) = 1 if  $q \equiv 1 \pmod{4}$ , and e(2,q) = 2 otherwise.

**Lemma 2.1.** [18, Corollary to Zsigmondy's theorem] Let q be a natural number greater than 1. For every natural number m there exists a prime r with e(r,q) = m, except for the cases q = 2 and m = 1, q = 3 and m = 1, and q = 2 and m = 6.

The prime r with e(r,q) = m is called a *primitive prime divisor* of  $q^m - 1$ . It is obvious that  $q^m - 1$  can have more than one primitive prime divisor. We denote by  $r_m(q)$  some primitive prime divisor of  $q^m - 1$ . Also,  $\eta(n)$  for an integer n, has been defined in [17] as follow:

$$\eta(n) = \left\{ egin{array}{c} n \ if \ n \ is \ odd; \ rac{n}{2} \ otherwise. \end{array} 
ight.$$

**Lemma 2.2.** [18, Proposition 2.4], [17, Proposition 3.1] Let G be one of the simple groups of Lie type  $B_n(q)$  or  $C_n(q)$  over a field of characteristic p and let r, s be odd primes with  $r, s \in \pi(G) \setminus \{p\}$ . Let k = e(r,q), l = e(s,q) and suppose that  $1 \leq \eta(k) \leq \eta(l)$ . The following statements hold:

- (1) r and s are nonadjacent if and only if  $\eta(k) + \eta(l) > n$  and  $\frac{l}{k}$  is not an odd natural number,
- (2) r and p are nonadjacent if and only if  $\eta(e(r,q)) > n-1$ .

**Lemma 2.3.** [17, Proposition 4.3] Let  $G = B_n(q)$  or  $G = C_n(q)$  be a finite simple group of Lie type over a field of order q with odd characteristic p. Let r be an odd prime divisor of |G|,  $r \neq p$ , and k = e(r,q). Then r and 2 are nonadjacent if and only if  $\eta(k) = n$  and one of the following holds:

- (1) *n* is odd and k = (3 e(2, q))n;
- (2) n is even and k = 2n.

**Definition 2.4.** [2] Let S be a finite simple group of Lie type in characteristic p. Let A be any abelian p-group with an S-action. An element  $s \in S$  is said to be unisingular on A if s has a (non-zero) fixed point on A. S is said to be unisingular if every element  $s \in S$  acts unisingularly on every finite abelian p-group A with an S-action.

**Lemma 2.5.** [7, Theorem 1.3] The simple groups  $B_n(q)$  and  $C_n(q)$  of characteristic p is unisingular if and only if  $q = p \neq 2$ .

**Lemma 2.6.** [14, Lemma 1] Let G be a finite group, let N be a normal subgroup of G and let G/N be a Frobenius group with Frobenius kernel F and cyclic complement C. If gcd(|F|, |N|) = 1 and F not lying in  $NC_G(N)/N$  then  $s.|C| \in \omega(G)$  for some  $s \in \pi(N)$ .

# 3. Main results

**Lemma 3.1.** If n > 3 is an odd number and q is the power of an odd prime p, then the groups  $B_n(q)$  and  $C_n(q)$  include subgroups isomorphic to Frobenius groups of the form  $U : Z_{r_n(q)}$  and  $T : Z_{r_{n-2}(q)}$ , where U and T are nontrivial p-groups.

**Proof.** By [13, Proposition 4.1.20],  $B_n(q)$  contains a subgroup isomorphic to  $U: SL_n(q)$ , where U is a nontrivial p-group. Since  $r_n(q) \in \pi(SL_n(q))$ ,  $SL_n(q)$  has a cyclic subgroup of order  $r_n(q)$ , namely Z. Now by Lemma 2.2(2), since n is odd, we can conclude that U: Z is the first desired Frobenius subgroup. Also, similar to the previous argument  $B_{n-2}(q)$  has a Frobenius subgroup of the form  $T: Z_{r_{n-2}(q)}$ , where T is a nontrivial p-group.

Let  $bd(C_1, C_2, ..., C_m)$  denote a block-diagonal matrix with square blocks  $C_1, C_2, ..., C_m$ . Put

$$W_n = \mathrm{bd}(\widetilde{W, W, ..., W}, 1), \text{ where } W = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

We have  $SO_{2n+1}(q) = \{X \in SL_{2n+1}(q) | X^t W_n X = W_n\}$  and  $B_n(q)$  is the derived subgroup of  $SO_{2n+1}(q)$ . Of course

 $M = \{ \operatorname{bd}(I_4, X) | X \in SL_{2n-3}(q), X^t W_{n-2}X = W_{n-2} \}$  is a subgroup of  $SO_{2n+1}(q)$  and it is isomorphic to  $SO_{2n-3}(q)$ . Thus the derived subgroup of M is a subgroup of  $B_n(q)$  and it is isomorphic to  $B_{n-2}(q)$ , so we can assume that  $B_{n-2}(q) \hookrightarrow B_n(q)$  and hence,  $B_n(q)$  has the second desired Frobenius subgroup as well.

2-recognizability of the simple groups  $B_n(3)$  and  $C_n(3)$  by prime graph

Now we consider  $C_n(q)$ . For  $J_n = \begin{bmatrix} \mathbf{0} & I_n \\ -I_n & \mathbf{0} \end{bmatrix}$  we have  $Sp_{2n}(q) = \{X \in GL_{2n}(q) | X^t J_n X = J_n\}$ . Let H and K be the following sets:

$$H := \left\{ \begin{bmatrix} I_n & A \\ \mathbf{0} & I_n \end{bmatrix} \mid A \in M_n(q) \text{ and } A^t = A \right\},$$
$$K := \left\{ \begin{bmatrix} B & \mathbf{0} \\ \mathbf{0} & (B^t)^{-1} \end{bmatrix} \mid B \in GL_n(q) \right\}.$$

Obviously,  $H, K \leq Sp_{2n}(q), K \cong GL_n(q)$  and  $K \leq N_{Sp_{2n}(q)}(H)$ . Also, it is easy to check that H is a nontrivial p-group and does not contain the center of  $Sp_{2n}(q)$  which consists of the matrices  $\pm I_{2n}$  and hence  $C_n(q)$  has a subgroup of the form U: K, where  $U = H\{\pm I_{2n}\}/\{\pm I_{2n}\}$ is a nontrivial p-group and  $\bar{K} = K/\{\pm I_{2n}\}$ . Since  $r_n(q) \in \pi(\bar{K}), \bar{K}$ has a cyclic subgroup of order  $r_n(q)$ , namely Z. By Lemma 2.2(2) since n is odd, we conclude that U: Z is one of the desired Frobenius subgroup of  $C_n(q)$ . Also  $N = \{ bd(I_2, X, I_2) | X \in GL_{2(n-2)}(q), X^t J_{n-2}X =$  $J_{n-2}$  is a subgroup of  $Sp_{2n}(q)$  and it is isomorphic to  $Sp_{2(n-2)}(q)$ . Thus  $N\{\pm I_{2n}\}/\{\pm I_{2n}\}$  is a subgroup of  $C_n(q)$  which is isomorphic to  $Sp_{2(n-2)}(q)$ . But as it was mentioned,  $Sp_{2(n-2)}(q)$  contains a subgroup of the form  $U_0: Z_0$ , where  $U_0$  a is nontrivial p-group and  $Z_0$  is a cyclic subgroup of order  $r_{n-2}(q)$ . We claim that  $U_0: Z_0$  is a Frobenius subgroup of  $Sp_{2(n-2)}(q)$ . If not, then  $U_0: Z_0$  contains an element of order  $pr_{n-2}(q)$ , so  $(p, r_{n-2}(q)) \in GK(Sp_{2(n-2)}(q))$ . Thus since  $p \neq 2$  and  $C_{n-2}(q) =$  $Sp_{2(n-2)}(q)/{\pm I}$ , we deduce that  $(p, r_{n-2}(q)) \in GK(C_{n-2}(q))$ , contradicting Lemma 2.2(2). This leads us to find the second desired Frobenius subgroup of  $C_n(q)$ .

**Proof of the main theorem.** In [6] we have proved that if  $n \ge 9$  and  $GK(G) = GK(B_n(3))$ , then there exists a finite nonabelian simple group S such that  $S \le G/K \le \operatorname{Aut}(S)$  and  $S \cong B_n(3)$  or  $S \cong C_n(3)$ , where K is the maximal normal solvable subgroup of G. Also, we know that  $|\operatorname{Out}(B_n(3))| = |\operatorname{Out}(C_n(3))| = 2$ , hence G/K = S or  $G/K = \operatorname{Aut}(S)$ . Thus the last step to complete the proof is to show that  $G \cong S$  and K = 1. We will do it in the following two steps:

**Step I.** In this step, we will prove that G/K = S. If not, then  $G/K = \operatorname{Aut}(S)$  and we will consider the cases  $S \cong B_n(3)$  and  $S \cong C_n(3)$  separately:

**Case 1.** If  $S \cong B_n(3)$ , then since  $[SO_{2n+1}(3) : B_n(3)] = 2$ , we have  $B_n(3)$  is a normal subgroup of  $SO_{2n+1}(3)$  and hence,

$$N_{SO_{2n+1}(3)}(B_n(3)) = SO_{2n+1}(3), \ B_n(3) \cap C_{SO_{2n+1}(3)}(B_n(3)) \triangleleft B_n(3)$$
  
and  $B_n(3)C_{SO_{2n+1}(3)}(B_n(3)) \le SO_{2n+1}(3).$ 

Thus, since  $B_n(3)$  is a nonabelian simple group, we deduce that the order of the group  $C_{SO_{2n+1}(3)}(B_n(3))$  divides 2. If  $|C_{SO_{2n+1}(3)}(B_n(3))| = 2$ , then since  $C_{SO_{2n+1}(3)}(B_n(3))$  is a normal subgroup of  $SO_{2n+1}(3)$ , we can easily conclude that  $C_{SO_{2n+1}(3)}(B_n(3)) \subseteq Z(SO_{2n+1}(3))$  which leads us to a contradiction, because  $Z(SO_{2n+1}(3)) = 1$ . Thus  $C_{SO_{2n+1}(3)}(B_n(3)) =$ 1. Now we have  $SO_{2n+1}(3) \cong N_{SO_{2n+1}(3)}(B_n(3))/C_{SO_{2n+1}(3)}(B_n(3))$ . So Aut(S) contains a subgroup isomorphic to  $SO_{2n+1}(3)$  and since  $SO_{2n+1}(3)$ has a maximal torus of order  $3^n - 1$ , we conclude that

$$(2, r_n(3)) \in GK(G) = GK(B_n(3))$$

which is impossible by Lemma 2.3.

**Case 2.** If  $S \cong C_n(3)$ , then  $(G/K)/C_n(3) = \operatorname{Out}(C_n(3))$ . As was stated in the proof of Lemma 3.1, for  $J_n = \begin{bmatrix} \mathbf{0} & I_n \\ -I_n & \mathbf{0} \end{bmatrix}$ , we have  $Sp_{2n}(3) = \{X \in GL_{2n}(3) \mid X^tJ_nX = J_n\}$  and  $C_n(3) = Sp_{2n}(3)/\{\pm I\}$ . According to [13, Relation 2.4.3 and Proposition 2.4.4],  $\operatorname{Out}(C_n(3)) = \langle \ddot{\delta} \rangle$ , where  $\delta = \operatorname{bd}(-I_n, I_n)\{\pm I\}$  and  $\ddot{\delta}$  denotes the image of  $\delta$  in  $\operatorname{Out}(C_n(3))$ . Thus  $\delta$  centralizes a subgroup  $K = \{\operatorname{bd}(A, (A^t)^{-1})\{\pm I\} | A \in GL_n(3)\}$  of  $C_n(3)$  which is isomorphic to  $GL_n(3)/\{\pm I\}$ . This shows that  $\operatorname{Aut}(C_n(3))$ has an element of order  $3^n - 1$  and we can get a contradiction similar to the previous case.

**Step II.** Here we want to show that K = 1. It suffices to show that  $GK(G) \neq GK(S)$  whenever  $K \neq 1$ . Replacing K by  $K/K_1$ , where  $K_1$  is a maximal subgroup of K which is normal in G allows us to assume that K is an elementary abelian p-group and S acts on K faithfully and irreducibly. If p = 3, then by Lemma 2.5 each element of S centralizes some nontrivial element of K, and so 3 and  $r_n(3)$  are adjacent which is impossible by Lemma 2.2(2).

If  $p \neq 3$ , then we can assume that e(p,3) = u. Of course by Lemma 3.1, there are Frobenius subgroups of the form  $U : Z_{r_n(3)}$  and  $T : Z_{r_{n-2}(3)}$  in S = G/K, where U and T are nontrivial 3-groups. According to Lemma 2.6, p is adjacent to  $r_n(3)$  and  $r_{n-2}(3)$  in GK(S).

Since  $(p, r_n(3)) \in GK(S)$ , Lemma 2.3 implies that  $p \neq 2$ . Thus by Lemma 2.2(1),  $n + \eta(u) \leq n$  or n/u is an odd number, because n is odd. This leads to  $u \mid n$ . Therefore u is an odd number, so  $\eta(u) = u$ . Also,  $(p, r_{n-2}(3)) \in GK(S)$ , so Lemma 2.2(1) implies that  $n - 2 + \eta(u) =$  $n - 2 + u \leq n$  or (n - 2)/u is an odd number. This shows that  $u \leq 2$  or  $u \mid (n, n - 2) = 1$  and hence, u = 1, because n and u are odd numbers. This implies that  $p = r_1(3)$  which is impossible by Lemma 2.1.

Thus, according to Steps I and II, the main theorem is obtained for  $n \ge 9$ . Also, if  $n \le 7$ , then the Main theorem in [15] completes the proof.

As consequences of our main result, we have the following corollaries:

**Corollary 3.2.** Let n be an odd number. The simple groups  $B_n(3)$  and  $C_n(3)$  are characterizable by their spectra except the case  $B_3(3)$  which is 2-recognizable.

*Proof.* Since  $\omega(B_n(3)) \neq \omega(C_n(3))$  [16, Proposition] and  $\omega(B_3(3)) = \omega(D_4(3))$  [4], thus the proof is straightforward by the main theorem.  $\Box$ 

If G is a finite group, by M(G) we denote the set of orders of maximal abelian subgroups of G and a group G is said to be characterizable by the set of orders of its maximal abelian subgroups, if G is uniquely determined by M(G).

**Lemma 3.3.** [3, Lemma 2] Let G and H be finite groups. If M(G) = M(H), then GK(G) = GK(H).

**Corollary 3.4.** If n is an odd number, then the simple groups  $B_n(3)$  and  $C_n(3)$  are characterizable by their sets of orders of maximal abelian subgroups.

Proof. Table 2 in [19] shows that  $M(C_n(3)) \neq M(B_n(3))$  and  $M(B_3(3)) \neq M(D_4(3)) \neq M(C_3(3))$ . Also, it implies that  $\max(M(B_3(3))) = 3^5$ ,  $\max(M(C_3(3))) = 3^6$  and  $\max(M(^2B_2(8))) = 2^4$ .

Also, since  $[\operatorname{Aut}({}^{2}B_{2}(8)): {}^{2}B_{2}(8)] = 3$ ,  $M(G/O_{2}(G)) \neq M(\operatorname{Aut}({}^{2}B_{2}(8)))$ , for every finite group G with  $M(G) = M(B_{3}(3))$  or  $M(G) = M(C_{3}(3))$ and hence, Lemma 3.3 and the main theorem complete the proof.  $\Box$ 

The Non-commuting graph of a nonabelian group G, denoted by  $\Gamma_G$  is the graph with vertex set  $G \setminus Z(G)$ , where two distinct vertices x and y are adjacent by an edge if  $xy \neq yx$ . Problem 16.1 in the Kourovka notebook [12] is AAM's conjecture, which says simple groups

П

are determined uniquely by the non-commuting graph. This conjecture is valid for all non-abelian finite simple groups with disconnected prime graph (for examples see paper by Darafsheh [5] and references quoted in that paper).

**Lemma 3.5.** [1, Theorem 2] Let P be a finite nonabelian simple group and G is a group such that  $\Gamma_G \cong \Gamma_P$ , then GK(G) = GK(P) and M(G) = M(P).

**Corollary 3.6.** The AAM's Conjecture is true for the groups under study.

Proof. By Lemma 3.5 and Corollary 3.4, the proof is straightforward.

### Acknowledgment

The authors wish to thank the referees for their invaluable comments. This work was partially supported by Center of Excellence of Algebraic Hyperstructures and its Applications of Tarbiat Modares University (CEAHA). This research is partially supported by Iran National Science Foundation (INSF) (Grant No. 91058621).

## References

- N. Ahanjideh and A. Iranmanesh, On the relation between the non-commuting graph and the prime graph, Int. J. Group Theory, 1 (2012), No. 1, 25–28.
- [2] L. Babai and A. Shalev, Recognizing simplicity of black-box groups and the frequency of *p*-singular elements in affine groups, 39–62, Groups and Computation, III (Columbus, OH, 1999), Ohio State Univ. Math. Res. Inst. Publ., 8, de Gruyter, Berlin, 2001.
- [3] G. Chen, A characterization of alternating groups by the set of orders of their maximal abelian subgroups, Sibrian Math. J. 47 (2006), no. 3, 594–596.
- [4] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, Atlas of Finite Groups, Clarendon Press, Oxford University Press, Eynsham, 1985.
- [5] M. R. Darafsheh, Groups with the same non-commuting graph, Discrete Appl. Math. 157 (2009), no. 4, 833–837.
- [6] M. Foroudi Ghasemabadi and A. Iranmanesh, 2-quasirecognizability of the simple groups  $B_n(p)$  and  $C_n(p)$  by prime graph, Bull. Iranian Math. Soc. **38** (2012), no. 3, 647–668.
- [7] R. M. Guralnick and P. H. Tiep, Finite simple unisingular groups of Lie type, J. Group Theory 6 (2003), no. 3, 271–310.
- [8] M. Hagie, The prime graph of a sporadic simple group, Comm. Algebra 31 (2003), no. 9, 4405–4424.

2-recognizability of the simple groups  $B_n(3)$  and  $C_n(3)$  by prime graph

- [9] A. Khosravi and B. Khosravi, 2-recognizability of  $PSL(2, p^2)$  by a prime graph, Sibrian Math. J. **49** (2008), no. 4, 749–757.
- [10] B. Khosravi, n-Recognition by prime graph of the simple group PSL(2,q), J. Algebra Appl. 7 (2008), no. 6, 735–748.
- [11] B. Khosravi, B. Khosravi and B. Khosravi, A characterization of the finite simple group L<sub>16</sub>(2) by its prime graph, *Manuscripta Math.* **126** (2008), no. 1, 49–58.
- [12] E. I. Khukhro and V. D. Mazurov, Unsolved problems in group theory: The Kourovka Notebook, Sobolev Institute of Mathematics, Novosibirsk, 17th edition, 2010.
- [13] P. B. Kleidman and M. Liebeck, The Subgroup Structure of Finite Classical Groups, Cambridge Univ. Press, Cambridge, 2005.
- [14] V. D. Mazurov, Characterizations of finite groups by sets of the orders of their elements, Algebra Logic 36 (1997), no. 1, 23–32.
- [15] Z. Momen and B. Khosravi, On *r*-recognition by prime graph of  $B_p(3)$  where *p* is an odd prime, *Monatsh. Math.* **166** (2012), no. 2, 239–253.
- [16] W. J. Shi, Pure quantitative characterization of finite simple groups, Front. Math. China 2 (2007), no. 1, 123–125.
- [17] A. V. Vasil'ev and E. P. Vdovin, An adjacency criterion in the prime graph of a finite simple group, *Algebra Logic* 44 (2005), no. 6, 381–406.
- [18] A. V. Vasil'ev and E. P. Vdovin, Cocliques of maximal size in the prime graph of a finite simple group, *Algebra Logic* 50 (2011), no. 4, 291–322.
- [19] E. P. Vdovin, Maximal orders of abelian subgroups in finite Chevalley groups, Mat. Zametki 69 (2001), no. 4, 524–549.
- [20] A. V. Zavarnitsine, Recognition of finite groups by the prime graph, Algebra Logic 45 (2006), no. 4, 220–231.
- [21] A. V. Zavarnitsine, Uniqueness of the prime graph of  $L_{16}(2)$ , Sib. Èlektron. Mat. Izv. 7 (2010), 119-121.
- [22] M. R. Zinov'eva, Recognition by spectrum of simple groups  $C_p(3)$ , Tr. Inst. Mat. Mekh. 16 (2010), no. 3, 88–95.
- [23] M. R. Zinov'eva, R. Shen and W. J. Shi, Recognition of simple groups  $B_p(3)$  by the set of element orders, *Sibrian Math. J.* **51** (2010), no. 2, 244–254.

#### Mahnaz Foroudi Ghasemabadi

Department of pure Mathematics, Faculty of Mathematical Sciences Tarbiat Modares University, P.O. Box 14115-137, Tehran, Iran E-mail: foroudi@modares.ac.ir

#### Ali Iranmanesh

Department of pure Mathematics, Faculty of Mathematical Sciences Tarbiat Modares University, P.O. Box 14115-137, Tehran, Iran E-mail: iranmanesh@modares.ac.ir

#### Neda Ahanjideh

Department of Mathematics, Faculty of basic Sciences University of Shahre-kord, P.O. Box 115, Shahre-kord, Iran E-mail: ahanjideh.neda@sci.sku.ac.ir