

## DETERMINANTS AND PERMANENTS OF HESSENBERG MATRICES AND GENERALIZED LUCAS POLYNOMIALS

K. KAYGISIZ\* AND A. ŞAHİN

Communicated by Amir Daneshgar

**ABSTRACT.** In this paper, we give some determinantal and permanent representations of generalized Lucas polynomials, which are a general form of generalized bivariate Lucas  $p$ -polynomials, ordinary Lucas and Perrin sequences etc., by using various Hessenberg matrices. In addition, we show that determinant and permanent of these Hessenberg matrices can be obtained by using combinations. Then we show, the conditions under which the determinants of the Hessenberg matrix become its permanents.

### 1. Introduction

Fibonacci numbers  $f_n$ , Lucas numbers  $l_n$  and Perrin numbers  $r_n$  are defined by

$$\begin{aligned}f_n &= f_{n-1} + f_{n-2} \text{ for } n > 2 \text{ and } f_1 = f_2 = 1, \\l_n &= l_{n-1} + l_{n-2} \text{ for } n > 1 \text{ and } l_0 = 2, l_1 = 1, \\r_n &= r_{n-2} + r_{n-3} \text{ for } n > 3 \text{ and } r_0 = 3, r_1 = 0, r_2 = 2,\end{aligned}$$

respectively.

---

MSC(2010): Primary: 11B37; Secondary: 15A15, 15A51.

Keywords: Generalized Lucas polynomials, generalized Perrin polynomials, Hessenberg matrix, determinant and permanent.

Received: 11 May 2012, Accepted: 12 August 2012.

\*Corresponding author

© 2013 Iranian Mathematical Society.

Generalizations of these sequences have been studied by a number of researchers. For instance; Miles [16] defined generalized order- $k$  Fibonacci numbers, Er [2] defined  $k$  sequences of generalized order- $k$  Fibonacci numbers and Kaygısız and Bozkurt [4] defined  $k$ -generalized order- $k$  Perrin numbers.

MacHenry [12] defined generalized Fibonacci polynomials  $F_{k,n}(t)$  and Lucas polynomials  $G_{k,n}(t)$  as follows;

$$\begin{aligned} F_{k,n}(t) &= 0, \quad n < 0, & G_{k,n}(t) &= 0, \quad n < 0, \\ F_{k,0}(t) &= 1, & G_{k,0}(t) &= k, \\ F_{k,n}(t) &= \sum_{j=1}^k t_j F_{k,n-j}(t), & G_{k,1}(t) &= t_1, \\ & & G_{k,n}(t) &= G_{k-1,n}(t), \quad 1 \leq n < k, \\ & & G_{k,n}(t) &= \sum_{j=1}^k t_j G_{k,n-j}(t), \quad n \geq k \end{aligned}$$

where  $t_i$  ( $1 \leq i \leq k$ ) are constant coefficients of the core polynomial

$$P(x; t_1, t_2, \dots, t_k) = x^k - t_1 x^{k-1} - \dots - t_k.$$

In [13, 14], authors obtained some properties of this polynomials. In addition, in [15], authors obtained ( $n, k \in \mathbb{N}, n \geq 1$ )

$$(1.1) \quad G_{k,n}(t) = \sum_{a \vdash n} \frac{n}{|a|} \binom{|a|}{a_1, \dots, a_k} t_1^{a_1} \dots t_k^{a_k}$$

where  $a_i$  are nonnegative integers for all  $i$  ( $1 \leq i \leq k$ ), with initial conditions given by

$$G_{k,0}(t) = k, \quad G_{k,-1}(t) = 0, \quad \dots, \quad G_{k,-k+1}(t) = 0.$$

Throughout this paper, the notations  $a \vdash n$  and  $|a|$  are used instead of  $\sum_{j=1}^k j a_j = n$  and  $\sum_{j=1}^k a_j$ , respectively.

Kaygısız and Şahin [5] defined generalized Perrin polynomials  $R_{k,n}(t)$  by using generalized Lucas polynomials.

The generalized bivariate Lucas  $p$ -polynomials [20] are defined by

$$L_{p,n}(x, y) = x L_{p,n-1}(x, y) + y L_{p,n-p-1}(x, y)$$

for  $n > p$ , with boundary conditions  $L_{p,0}(x, y) = (p+1)$ ,  $L_{p,n}(x, y) = x^n$ ,  $n = 1, 2, \dots, p$ .

TABLE 1. Cognate polynomial sequences 1.

$\mathbf{k}$	$\mathbf{t}_1$	$\mathbf{t}_i(2 \leq i \leq (k - 1))$	$\mathbf{t}_k$	$\mathbf{G}_{\mathbf{k},\mathbf{n}}(\mathbf{t})$
$k$	0	$t_i$	$t_k$	$R_{k,n}(t)$
$k$	$x$	0	$y$	$L_{p,n}(x, y)$
3	0	1	1	$r_n$

TABLE 2. [20]Cognate polynomial sequences 2.

$\mathbf{x}$	$\mathbf{y}$	$\mathbf{p}$	$\mathbf{L}_{\mathbf{p},\mathbf{n}}(\mathbf{x}, \mathbf{y})$
$x$	$y$	1	bivariate Lucas polynomials $L_n(x, y)$
$x$	1	$p$	Lucas $p$ -polynomials $L_{p,n}(x)$
$x$	1	1	Lucas polynomials $l_n(x)$
1	1	$p$	Lucas $p$ -numbers $L_p(n)$
1	1	1	Lucas numbers $L_n$
$2x$	$y$	$p$	bivariate Pell-Lucas $p$ -polynomials $L_{p,n}(2x, y)$
$2x$	$y$	1	bivariate Pell-Lucas polynomials $L_n(2x, y)$
$2x$	1	$p$	Pell-Lucas $p$ -polynomials $Q_{p,n}(x)$
$2x$	1	1	Pell-Lucas polynomials $Q_n(x)$
2	1	1	Pell-Lucas numbers $Q_n$
$2x$	-1	1	Chebysev polynomials of the first kind $T_n(x)$
$x$	$2y$	$p$	bivariate Jacobsthal-Lucas $p$ -polynomials $L_{p,n}(x, 2y)$
$x$	$2y$	1	bivariate Jacobsthal-Lucas polynomials $L_n(x, 2y)$
1	$2y$	1	Jacobsthal-Lucas polynomials $j_n(y)$
1	2	1	Jacobsthal-Lucas numbers $j_n$

Tables 1 and 2 show that  $G_{k,n}(t)$  are general form of many sequences and polynomials. Therefore, any result obtained from the polynomials  $G_{k,n}(t)$  is valid for all sequences and polynomials mentioned in these tables.

On the other hand, many researchers studied determinantal and permanental representations of  $k$  sequences of generalized order- $k$  Fibonacci and Lucas numbers. For example, Minc [17] defined an  $n \times n$  (0,1)-matrix  $F(n, k)$  and showed that the permanents of  $F(n, k)$  is equal to the generalized order- $k$  Fibonacci numbers.

In [10, 11], authors defined two (0,1)-matrices and showed that the permanents of these matrices are the generalized Fibonacci and Lucas numbers. Öcal et al. [18] gave some determinantal and permanental representations of  $k$ -generalized Fibonacci and Lucas numbers and obtained Binet's formulas for these sequences. Kılıç and Stakhov [8] gave

permanent representation of Fibonacci and Lucas  $p$ -numbers. Kılıç and Taşçı [9] studied permanents and determinants of Hessenberg matrices. Yılmaz and Bozkurt [22] derived some relationships between Pell and Perrin sequences, as well as permanents and determinants of a type of Hessenberg matrices. Kaygısız and Şahin [7] gave some determinantal and permanental representations of Fibonacci type numbers. Kaygısız and Şahin [6] gave some determinantal and permanental representations of generalized bivariate Lucas  $p$ -polynomials. In [3, 19, 21], authors gave some relations between determinants and permanents.

The main purpose of this paper is to give some determinantal and permanental representations of generalized Lucas polynomials by using various Hessenberg matrices. Then we provide some conditions under which the determinants of the Hessenberg matrix become its permanents.

## 2. The determinantal representations

An  $n \times n$  matrix  ${}_+A_n = (a_{ij})$  is called lower Hessenberg matrix if  $a_{ij} = 0$  when  $j - i > 1$  i.e.,

$$(2.1) \quad {}_+A_n = \begin{bmatrix} a_{11} & a_{12} & 0 & \cdots & 0 \\ a_{21} & a_{22} & a_{23} & \cdots & 0 \\ a_{31} & a_{32} & a_{33} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \cdots & a_{n-1,n} \\ a_{n,1} & a_{n,2} & a_{n,3} & \cdots & a_{n,n} \end{bmatrix}$$

**Lemma 2.1.** [1] Let  ${}_+A_n$  be the  $n \times n$  lower Hessenberg matrix for all  $n \geq 1$  and define  $\det({}_+A_0) = 1$ . Then,  $\det({}_+A_1) = a_{11}$  and for  $n \geq 2$

$$(2.2) \quad \det({}_+A_n) = a_{n,n} \det({}_+A_{n-1}) + \sum_{r=1}^{n-1} [(-1)^{n-r} a_{n,r} \left( \prod_{j=r}^{n-1} a_{j,j+1} \right) \det({}_+A_{r-1})].$$

Now, we define two Hessenberg matrices  ${}_+C_{k,m}$  and  ${}_+C_{k,m}$  whose determinants give the generalized Lucas polynomials.

**Theorem 2.2.** Let  $k \geq 2$  and  $n \geq 1$  be integers, and let  $G_{k,n}(t)$  be the generalized Lucas polynomials and  ${}_+C_{k,n} = (c_{rs})$  an  $n \times n$  Hessenberg

matrix, given by

$$c_{rs} = \begin{cases} i^{|r-s|} \cdot \frac{t_{r-s+1}}{t_2^{(r-s)}}, & \text{if } s \neq 1 \text{ and } -1 \leq r-s < k, \\ i^{|r-s|} \cdot \frac{t_{r-s+1}}{t_2^{(r-s)}} \cdot (r-s+1), & \text{if } s = 1 \text{ and } -1 \leq r-s < k, \\ 0, & \text{otherwise} \end{cases}$$

i.e.,

$$+C_{k,n} = \begin{bmatrix} t_1 & it_2 & 0 & 0 & \cdots & 0 \\ 2i & t_1 & it_2 & 0 & \cdots & 0 \\ 3i^2 \frac{t_3}{t_2} & i & t_1 & it_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ ki^{k-1} \frac{t_k}{t_2^{k-1}} & i^{k-2} \frac{t_{k-1}}{t_2^{k-2}} & i^{k-3} \frac{t_{k-2}}{t_2^{k-3}} & i^{k-4} \frac{t_{k-3}}{t_2^{k-4}} & \cdots & 0 \\ 0 & i^{k-1} \frac{t_k}{t_2^{k-1}} & i^{k-2} \frac{t_{k-1}}{t_2^{k-2}} & i^{k-3} \frac{t_{k-2}}{t_2^{k-3}} & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & it_2 \\ 0 & 0 & 0 & \cdots & i & t_1 \end{bmatrix}$$

where  $t_0 = 1$  and  $i = \sqrt{-1}$ . Then,

$$(2.3) \quad \det(+C_{k,n}) = G_{k,n}(t).$$

*Proof.* To prove (2.3), we use the mathematical induction on  $m$ . The result is true for  $m = 1$  by hypothesis.

Assume that it is true for all positive integers less than or equal to  $m$ , namely,  $\det(+C_{k,m}) = G_{k,m}(t)$ . Then, by using Lemma 2.1, we have

$$\begin{aligned} \det(+C_{k,m+1}) &= c_{m+1,m+1} \det(+C_{k,m}) \\ &\quad + \sum_{r=1}^m \left[ (-1)^{m+1-r} c_{m+1,r} \prod_{j=r}^m c_{j,j+1} \det(+C_{k,r-1}) \right] \\ &= t_1 \det(+C_{k,m}) + \sum_{r=1}^{m-k+1} \left[ (-1)^{m+1-r} c_{m+1,r} \prod_{j=r}^m c_{j,j+1} \det(+C_{k,r-1}) \right] \\ &\quad + \sum_{r=m-k+2}^m \left[ (-1)^{m+1-r} c_{m+1,r} \prod_{j=r}^m c_{j,j+1} \det(+C_{k,r-1}) \right] \end{aligned}$$

$$\begin{aligned}
&= t_1 \det(+C_{k,m}) \\
&\quad + \sum_{r=m-k+2}^m \left[ (-1)^{m+1-r} c_{m+1,r} \prod_{j=r}^m c_{j,j+1} \det(+C_{k,r-1}) \right] \\
&= t_1 \det(+C_{k,m}) \\
&\quad + \sum_{r=m-k+2}^m \left[ (-1)^{m+1-r} .i^{m+1-r} \frac{t_{m-r+2}}{t_2^{(m-r+1)}} \prod_{j=r}^m it_2 \det(+C_{k,r-1}) \right] \\
&= t_1 \det(+C_{k,m}) \\
&\quad + \sum_{r=m-k+2}^m \left[ (-i)^{m+1-r} \frac{t_{m-r+2}}{t_2^{(m-r+1)}} .i^{m+1-r} .t_2^{(m-r+1)} \det(+C_{k,r-1}) \right] \\
&= t_1 \det(+C_{k,m}) + \sum_{r=m-k+2}^m t_{m-r+2} \det(+C_{k,r-1}) \\
&= t_1 \det(+C_{k,m}) + t_2 \det(+C_{k,m-1}) + \cdots + t_k \det(+C_{k,m-(k-1)}).
\end{aligned}$$

From the hypothesis and definition of generalized Lucas polynomials we obtain

$$\det(+C_{k,m+1}) = t_1 G_{k,m}(t) + \cdots + t_k G_{k,m-(k-1)}(t) = G_{k,m+1}(t).$$

Therefore, (2.3) holds for all positive integers.  $\square$

**Example 2.3.** We obtain 6-th generalized Lucas polynomial for  $k = 5$ , by using (2.3).

$$\begin{aligned}
\det(+C_{5,6}) &= \det \begin{bmatrix} t_1 & it_2 & 0 & 0 & 0 & 0 \\ 2i & t_1 & it_2 & 0 & 0 & 0 \\ 3 \frac{-t_3}{t_2^2} & i & t_1 & it_2 & 0 & 0 \\ 4 \frac{-it_4}{t_2^3} & \frac{-t_3}{t_2^2} & i & t_1 & it_2 & 0 \\ 5 \frac{t_5}{t_2^4} & \frac{-it_4}{t_2^3} & \frac{-t_3}{t_2^2} & i & t_1 & it_2 \\ 0 & \frac{t_5}{t_2^4} & \frac{-it_4}{t_2^3} & \frac{-t_3}{t_2^2} & i & t_1 \end{bmatrix} \\
&= t_1^6 + 6t_1^4 t_2 + 9t_1^2 t_2^2 + 2t_2^3 + 6t_1^3 t_3 + 3t_3^2 + 12t_1 t_2 t_3 \\
&\quad + 6t_1^2 t_4 + 6t_2 t_4 + 6t_1 t_5 \\
&= G_{5,6}(t).
\end{aligned}$$

**Theorem 2.4.** Let  $k \geq 2$  and  $n \geq 1$  be integers,  $G_{k,n}$  the generalized Lucas polynomial and  $-C_{k,n} = (b_{ij})$  an  $n \times n$  lower Hessenberg matrix,

given by

$$b_{ij} = \begin{cases} -t_2, & \text{if } j = i + 1, \\ \frac{t_{i-j+1}}{t_2^{(i-j)}}, & \text{if } j \neq 1 \text{ and } 0 \leq i - j < k, \\ \frac{t_{i-j+1}}{t_2^{(i-j)}} \cdot (i - j + 1), & \text{if } j = 1 \text{ and } 0 \leq i - j < k, \\ 0, & \text{otherwise} \end{cases}$$

i.e.,

$$-C_{k,n} = \begin{bmatrix} t_1 & -t_2 & 0 & 0 & \cdots & 0 \\ 2 & t_1 & -t_2 & 0 & \cdots & 0 \\ 3 \frac{t_3}{t_2^2} & 1 & t_1 & -t_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ k \frac{t_k}{t_2^{k-1}} & \frac{t_{k-1}}{t_2^{k-2}} & \frac{t_{k-2}}{t_2^{k-3}} & \cdots & \cdots & 0 \\ 0 & \frac{t_k}{t_2^{k-1}} & \frac{t_{k-1}}{t_2^{k-2}} & \frac{t_{k-2}}{t_2^{k-3}} & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & -t_2 \\ 0 & 0 & 0 & \cdots & \cdots & t_1 \end{bmatrix}$$

where  $t_0 = 1$ . Then,

$$(2.4) \quad \det(-C_{k,n}) = G_{k,n}(t).$$

*Proof.* The proof is similar to the proof of Theorem 2.2, by using Lemma 2.1.  $\square$

**Example 2.5.** We obtain 5-th generalized Lucas polynomial for  $k = 4$ , by using (2.4).

$$\begin{aligned} \det(-C_{4,5}) &= \det \begin{bmatrix} t_1 & -t_2 & 0 & 0 & 0 \\ 2 & t_1 & -t_2 & 0 & 0 \\ 3 \frac{t_3}{t_2^2} & 1 & t_1 & -t_2 & 0 \\ 4 \frac{t_4}{t_2^3} & \frac{t_3}{t_2^2} & 1 & t_1 & -t_2 \\ 0 & \frac{t_4}{t_2^3} & \frac{t_3}{t_2^2} & 1 & t_1 \end{bmatrix} \\ &= t_1^5 + 5t_1^3t_2 + 5t_1t_2^2 + 5t_1^2t_3 + 5t_2t_3 + 5t_1t_4 \\ &= G_{4,5}(t). \end{aligned}$$

**Corollary 2.6.** If we rewrite equalities (2.3) and (2.4) for  $t_i = 1$  and  $k = 2$ , then we obtain

$$\det(+C_{k,n}) = \det(-C_{k,n}) = l_n$$

where  $l_n$  are the ordinary Lucas numbers.

**Corollary 2.7.** *If we rewrite equalities (2.3) and (2.4) for  $t_1 = 0$ , then we obtain*

$$\det({}_+C_{k,n}) = \det({}_-C_{k,n}) = R_{k,n}(t)$$

where  $R_{k,n}(t)$  are the generalized Perrin polynomials.

**Corollary 2.8.** *If we rewrite the right hand side of equalities (2.3) and (2.4) for  $t_1 = x$ ,  $t_k = y$ ,  $t_i = 0$  ( $2 \leq i \leq k-1$ ) and  $k = (p+1)$ , then we obtain*

$$\det({}_+C_{k,n}) = \det({}_-C_{k,n}) = L_{p,n}(x, y)$$

where  $L_{p,n}(x, y)$  are the generalized bivariate Lucas  $p$ -polynomials.

**Corollary 2.9.** *If we rewrite equalities (2.3) and (2.4) for  $t_1 = 0$  and  $t_i = 1$  ( $2 \leq i \leq k$ ) and  $k = 3$ , then we obtain*

$$\det({}_+C_{k,n}) = \det({}_-C_{k,n}) = r_n$$

where  $r_n$  are the ordinary Perrin numbers.

Now we show that determinants of Hessenberg matrices  ${}_+C_{k,n}$  and  ${}_+C_{k,n}$  can be obtained by using combinations.

**Corollary 2.10.**

$$\det({}_+C_{k,n}) = \det({}_-C_{k,n}) = \sum_{a \mid n} \frac{n}{|a|} \binom{|a|}{a_1, \dots, a_k} t_1^{a_1} \dots t_k^{a_k}$$

*Proof.* It is obvious from Theorem 2.2, Theorem 2.4 and (1.1).  $\square$

### 3. The permanent representations

Let  $A = (a_{i,j})$  be an  $n \times n$  square matrix over a ring  $R$ . Then, it is well known that, the permanent of  $A$  is defined by

$$\text{per}(A) = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i,\sigma(i)}$$

where  $S_n$  denotes the symmetric group on  $n$  letters.

**Lemma 3.1.** [18] *Let  $A_n$  be an  $n \times n$  lower Hessenberg matrix for all  $n \geq 1$  and define  $\text{per}(A_0) = 1$ . Then,  $\text{per}(A_1) = a_{11}$  and for  $n \geq 2$*

$$(3.1) \quad \text{per}(A_n) = a_{n,n} \text{per}(A_{n-1}) + \sum_{r=1}^{n-1} (a_{n,r} \prod_{j=r}^{n-1} a_{j,j+1} \text{per}(A_{r-1})).$$



We define two Hessenberg matrices  $-H_{k,n}$  and  $+H_{k,n}$  whose permanents give the generalized Lucas polynomials.

**Theorem 3.2.** Let  $k \geq 2$  and  $n \geq 1$  be integers,  $G_{k,n}(t)$  the generalized Lucas polynomials and  $-H_{k,n} = (h_{rs})$  an  $n \times n$  lower Hessenberg matrix, given by

$$h_{rs} = \begin{cases} i^{(r-s)} \cdot \frac{t_{r-s+1}}{t_2^{(r-s)}}, & \text{if } s \neq 1 \text{ and } -1 \leq r-s < k, \\ i^{(r-s)} \cdot \frac{t_{r-s+1}}{t_2^{(r-s)}} \cdot (r-s+1), & \text{if } s = 1 \text{ and } -1 \leq r-s < k, \\ 0, & \text{otherwise} \end{cases}$$

i.e.,

$$-H_{k,n} = \begin{bmatrix} t_1 & -it_2 & 0 & 0 & \dots & 0 \\ 2i & t_1 & -it_2 & 0 & \dots & 0 \\ 3i^2 \frac{t_3}{t_2} & i & t_1 & -it_2 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ ki^{k-1} \frac{t_k}{t_2^{k-1}} & i^{k-2} \frac{t_{k-1}}{t_2^{k-2}} & i^{k-3} \frac{t_{k-2}}{t_2^{k-3}} & i^{k-4} \frac{t_{k-3}}{t_2^{k-4}} & \dots & 0 \\ 0 & i^{k-1} \frac{t_k}{t_2^{k-1}} & i^{k-2} \frac{t_{k-1}}{t_2^{k-2}} & i^{k-3} \frac{t_{k-2}}{t_2^{k-3}} & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & \dots & t_1 \end{bmatrix}$$

where  $t_0 = 1$  and  $i = \sqrt{-1}$ . Then,

$$(3.2) \quad \text{per}(-H_{k,n}) = G_{k,n}(t).$$

*Proof.* The proof is similar to the proof of Theorem 2.2, by using Lemma 3.1. □

**Example 3.3.** We obtain the 3-rd generalized Lucas polynomial for  $k = 4$ , by using (3.2)

$$\begin{aligned} \text{per}(-H_{4,3}) &= \text{per} \begin{bmatrix} t_1 & -it_2 & 0 \\ 2i & t_1 & -it_2 \\ 3 \frac{t_3}{t_2} & i & t_1 \end{bmatrix} \\ &= t_1^3 + 3t_1t_2 + 3t_3. \end{aligned}$$

**Theorem 3.4.** Let  $k \geq 2$  and  $n \geq 1$  be integers,  $G_{k,n}(t)$  the generalized Lucas polynomials and  $+H_{k,n} = (p_{ij})$  an  $n \times n$  lower Hessenberg matrix

given by

$$p_{ij} = \begin{cases} \frac{t_{i-j+1}}{t_2^{(i-j)}}, & \text{if } j \neq 1 \text{ and } -1 \leq i-j < k, \\ \frac{t_{i-j+1}}{t_2^{(i-j)}} \cdot (i-j+1), & \text{if } j = 1 \text{ and } 0 \leq i-j < k, \\ 0, & \text{otherwise} \end{cases}$$

i.e.,

$${}_+H_{k,n} = \begin{bmatrix} t_1 & t_2 & 0 & 0 & \cdots & 0 \\ 2 & t_1 & t_2 & 0 & \cdots & 0 \\ 3 \frac{t_3}{t_2^2} & 1 & t_1 & t_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ k \frac{t_k}{t_2^{k-1}} & \frac{t_{k-1}}{t_2^{k-2}} & \frac{t_{k-2}}{t_2^{k-3}} & \frac{t_{k-3}}{t_2^{k-4}} & \cdots & 0 \\ 0 & \frac{t_k}{t_2^{k-1}} & \frac{t_{k-1}}{t_2^{k-2}} & \frac{t_{k-2}}{t_2^{k-3}} & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \cdots & t_1 \end{bmatrix}$$

where  $t_0 = 1$ . Then,

$$(3.3) \quad \text{per}({}_+H_{k,n}) = G_{k,n}(t).$$

*Proof.* The proof is similar to the proof of Theorem 2.2, by using Lemma 3.1.  $\square$

**Corollary 3.5.** *If we rewrite equalities (3.2) and (3.3) for  $t_i = 1$  ( $1 \leq i \leq k$ ) and  $k = 2$ , then we obtain*

$$\text{per}(-H_{k,n}) = \text{per}({}_+H_{k,n}) = l_n$$

where  $l_n$  are the ordinary Lucas numbers.

**Corollary 3.6.** *If we rewrite equalities (3.2) and (3.3) for  $t_1 = 0$  and  $t_i = 1$  ( $2 \leq i \leq k$ ), then we obtain*

$$\text{per}(-H_{k,n}) = \text{per}({}_+H_{k,n}) = R_{k,n}(t)$$

where  $R_{k,n}(t)$  are the generalized Perrin polynomials.

**Corollary 3.7.** *If we rewrite the right hand side of equalities (3.2) and (3.3) for  $t_1 = x$ ,  $t_k = y$ ,  $t_i = 0$  ( $2 \leq i \leq k-1$ ) and  $k = (p+1)$ , then we obtain*

$$\text{per}(-H_{k,n}) = \text{per}({}_+H_{k,n}) = L_{p,n}(x, y)$$

where  $L_{p,n}(x, y)$  are the generalized bivariate Lucas  $p$ -polynomials.

**Corollary 3.8.** *If we rewrite equalities (3.2) and (3.3) for  $t_1 = 0, t_i = 1$  ( $2 \leq i \leq k$ ) and  $k = 3$ , then we obtain*

$$\text{per}(-H_{k,n}) = \text{per}(+H_{k,n}) = r_n$$

where  $r_n$  are the ordinary Perrin numbers.

Now we show that permanent of Hessenberg matrices  $-H_{k,n}$  and  $+H_{k,n}$  can be obtained by using combinations.

**Corollary 3.9.**

$$\text{per}(-H_{k,n}) = \text{per}(+H_{k,n}) = \sum_{a^+} \frac{n}{|a|} \binom{|a|}{a_1, \dots, a_k} t_1^{a_1} \dots t_k^{a_k}$$

*Proof.* It is obvious from Theorem 3.2, Theorem 3.4 and (1.1). □

#### 4. Determinant and permanent of a Hessenberg matrix

Gibson [3] gave an identity between permanent and determinant of a semitriangular matrix. We give a different proof of this identity for Hessenberg matrices.

**Theorem 4.1.** *Let  $+A_n$  be the Hessenberg matrix in (2.1) and*

$-A_n = (b_{ij})$  *an  $n \times n$  Hessenberg matrix, given by*

$$b_{ij} = \begin{cases} 0, & \text{if } j - i > 1, \\ -a_{ij}, & \text{if } j - i = 1, \\ a_{ij}, & \text{otherwise} \end{cases}$$

*i. e.,*

$$-A_n = \begin{bmatrix} a_{11} & -a_{12} & 0 & \cdots & 0 \\ a_{21} & a_{22} & -a_{23} & \cdots & 0 \\ a_{31} & a_{32} & a_{33} & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \cdots & -a_{n-1,n} \\ a_{n,1} & a_{n,2} & a_{n,3} & \cdots & a_{n,n} \end{bmatrix}.$$

*Then,*

$$(4.1) \quad \det(-A_n) = \text{per}(+A_n) \text{ and } \det(+A_n) = \text{per}(-A_n).$$

*Proof.* To prove (4.1), we use the mathematical induction on  $m$ . The result is true for  $m = 1$  by hypothesis.

Assume that it is true for all positive integers less than or equal to  $m$ , namely  $\det(-A_m) = \text{per}(+A_m)$ . Then, by using (2.2) and (3.1), we have

$$\begin{aligned}
 \det(-A_{m+1}) &= a_{m+1,m+1} \det(-A_m) \\
 &+ \sum_{r=1}^m [(-1)^{m+1-r} a_{m+1,r} \prod_{j=r}^m b_{j,j+1} \det(-A_{r-1})] \\
 &= a_{m+1,m+1} \text{per}(+A_m) \\
 &+ \sum_{r=1}^m [(-1)^{m+1-r} a_{m+1,r} \prod_{j=r}^m (-a_{j,j+1}) \text{per}(+A_{r-1})] \\
 &= a_{m+1,m+1} \text{per}(+A_m) \\
 &+ \sum_{r=1}^m [(-1)^{m+1-r} a_{m+1,r} (-1)^{m+1-r} \prod_{j=r}^m a_{j,j+1} \text{per}(+A_{r-1})] \\
 &= a_{m+1,m+1} \text{per}(+A_m) + \sum_{r=1}^m [a_{m+1,r} \prod_{j=r}^{m+1} a_{j,j+1} \text{per}(+A_{r-1})] \\
 &= \text{per}(+A_{m+1}).
 \end{aligned}$$

Therefore, the result is true for all positive integers.  $\square$

### Conclusion

Generalized Lucas polynomials are a general form of several polynomials and number sequences. Therefore any result obtained from these polynomials is applicable to the others. In addition, the relation between determinant and permanent of a Hessenberg matrix make it possible to transfer any result between them.

### Acknowledgments

The authors would like to express their appreciation to the anonymous reviewers for their careful reading and making some useful comments, improving the presentation of the paper.

## REFERENCES

- [1] N. D. Cahill, J.R. D'Errico, D. A. Narayan and J. Y. Narayan, Fibonacci determinants, *College Math. J.* **33** (2002), no. 3, 221–225.
- [2] M. C. Er, Sums of Fibonacci numbers by matrix method, *Fibonacci Quart.* **22** (1984), no. 3, 204–207.
- [3] P. M. Gibson, An identity between permanents and determinants, *Amer. Math. Monthly* **76** (1969) 270–271.
- [4] K. Kaygısız and D. Bozkurt,  $k$ -generalized order- $k$  Perrin number presentation by matrix method, *Ars Combin.* **105** (2012) 95–101.
- [5] K. Kaygısız and A. Şahin, Generalized Van der Laan and Perrin polynomials, and generalizations of Van der Laan and Perrin numbers, *Selçuk J. Appl. Math.* **14** (2013), no. 1, 89–103.
- [6] K. Kaygısız and A. Şahin, Generalized bivariate Lucas  $p$ -polynomials and Hessenberg Matrices, *J. Integer Seq.* **15** (2012), no. 3, 8 pages.
- [7] K. Kaygısız and A. Şahin, Determinant and permanent of Hessenberg matrices and Fibonacci type numbers, *Gen. Math. Notes* **9** (2012), no. 2, 32–41.
- [8] E. Kılıç and A. P. Stakhov, On the Fibonacci and Lucas  $p$ -numbers, their sums, families of bipartite graphs and permanents of certain matrices, *Chaos Solitons Fractals* **40** (2009), no. 5, 2210–2221.
- [9] E. Kılıç and D. Taşçı, On the generalized Fibonacci and Pell sequences by Hessenberg matrices, *Ars Combin.* **94** (2010) 161–174.
- [10] G. Y. Lee and S. G. Lee, A note on generalized Fibonacci numbers, *Fibonacci Quart.* **33** (1995), no. 3, 273–278.
- [11] G.-Y. Lee,  $k$ -Lucas numbers and associated bipartite graphs, *Linear Algebra Appl.* **320** (2000), no. 1-3, 51–61.
- [12] T. MacHenry, A subgroup of the group of units in the ring of arithmetic functions, *Rocky Mountain J. Math.* **39** (1999), no. 3, 1055–1065.
- [13] T. MacHenry, Generalized Fibonacci and Lucas polynomials and multiplicative arithmetic functions, *Fibonacci Quart.* **38** (2000), no. 2, 167–173.
- [14] T. MacHenry and K. Wong, Degree  $k$  linear recursions mod  $(p)$  and number fields, *Rocky Mountain J. Math.* **41** (2011), no. 4, 1303–1327.
- [15] T. MacHenry and K. Wong, A correspondence between the isobaric ring and multiplicative arithmetic functions, *Rocky Mountain J. Math.* **42** (2012), no. 4, 1247–1290.
- [16] E. P. Miles, Generalized Fibonacci numbers and associated matrices, *Amer. Math. Monthly.* **67** (1960) 745–752.
- [17] H. Minc, *Encyclopedia of Mathematics and its Applications*, 6, Addison-Wesley Publishing Co., London, 1978.
- [18] A. A. Öcal, N. Tuglu, E. Altinisik, On the representation of  $k$ -generalized Fibonacci and Lucas numbers, *Appl. Math. Comput.* **170** (2005), no. 1, 584–596.
- [19] V. E. Tarakanov and R. A. Zatorskii, A relationship between determinants and permanents, *Math. Notes* **85** (2009) 267–273.
- [20] N. Tuglu, E. G. Kocer and A. Stakhov, Bivariate Fibonacci like  $p$ -polynomials, *Appl. Math. Comput.* **217** (2011), no. 24, 10239–10246.
- [21] C. Wenchang, Determinant, permanent, and MacMahon's master theorem, *Linear Algebra Appl.* **255** (1997) 171–183.

- [22] F. Yılmaz and D. Bozkurt, Hessenberg matrices and the Pell and Perrin numbers, *J. Number Theory* **131** (2011), no. 8, 1390–1396.

**K. Kaygısız**

Department of Mathematics, Faculty of Arts and Sciences, Gaziosmanpaşa University, 60250 Tokat, Turkey

Email: [kenan.kaygisiz@gop.edu.tr](mailto:kenan.kaygisiz@gop.edu.tr)

**A. Şahin**

Department of Mathematics, Faculty of Education, Gaziosmanpaşa University, 60250 Tokat, Turkey

Email: [adem.sahin@gop.edu.tr](mailto:adem.sahin@gop.edu.tr)

Archive of SID