Bulletin of the Iranian Mathematical Society Vol. 39 No. 6 (2013), pp 1213-1222.

THE *nc*-SUPPLEMENTED SUBGROUPS OF FINITE GROUPS[†]

S. GUO, S. LIU^{*} AND W. SHI

Communicated by Jamshid Moori

ABSTRACT. A subgroup H is said to be *nc*-supplemented in a group G if there exists a subgroup $K \leq G$ such that $HK \triangleleft G$ and $H \cap K$ is contained in H_G , the core of H in G. We characterize the supersolubility of finite groups G with that every maximal subgroup of the Sylow subgroups is *nc*-supplemented in G.

1. Introduction

In this paper the word group always means finite group.

A subgroup H is said to be complemented in G if there exists a subgroup K such that G = HK and $H \cap K = 1$. Hall proved that a group is soluble if and only if every Sylow subgroup is complemented [7]. Ramadan in [13] proved that if G/H is supersoluble and all maximal subgroups of the Sylow subgroups of H are normal in G, then G is supersoluble. A subgroup H is c-normal in G if there exists a normal subgroup N of G such that HN = G and $H \cap N$ is contained in H_G , the core of H in G (see [17]). Obviously c-normality is weaker than normality. A subgroup H is said to be c-supplemented in a group G if

MSC(2010): Primary: 20D10; Secondary: 20D20, 20D40.

Keywords: Soluble group, *nc*-supplemented subgroup, normal subgroup, supersoluble group. Received: 10 September 2011, Accepted: 24 November 2012.

^{*} Corresponding author

[†] The editorial board of the Bulletin of the Iranian Mathematical Society (BIMS) has decided to retract this paper (for more details see Vol. 40 (2014), No. 1, page 293.)

^{© 2013} Iranian Mathematical Society.

¹²¹³

there exists a subgroup K such that HK = G and $H \cap K$ is contained in H_G , the core of H in G (see [3]). The notion of c-supplementation is a generalization of the notions of complement and c-normality. Li et al. in [12] defined the following concept: A subgroup H is said to be *nc*-supplemented in a group G if there exists a subgroup $K \leq G$ such that $HK \triangleleft G$ and $H \cap K$ is contained in H_G , the core of H in G.

In this note, we give some generalization of supersolubility based on the concept of nc-supplementation.

We will prove the following theorem:

Theorem 1.1. Suppose that G is a group with a normal subgroup H such that G/H is supersoluble. If every maximal subgroup of every Sylow subgroup of H is nc-supplemented in G, then G is supersoluble.

A class of finite group \mathfrak{F} is said to be a formation if every epimorphic image of an \mathfrak{F} -group is an \mathfrak{F} -group and if $G/N_1 \cap N_2$ belongs to \mathfrak{F} whenever G/N_1 and G/N_2 belong to \mathfrak{F} . A formation \mathfrak{F} is said to be saturated if a finite group $G \in \mathfrak{F}$ whenever $G/\Phi(G) \in \mathfrak{F}$ (see [14, p. 277]). The class of supersoluble group is a saturated formation (see [14, 9.4.5]). Let \mathfrak{U} denote the class of all supersoluble groups.

Also we prove:

Theorem 1.2. Let \mathfrak{F} be a saturated formation containing \mathfrak{U} . Suppose that G is a group with a normal subgroup H such that $G/H \in \mathfrak{F}$. If every maximal subgroup of all Sylow subgroups of H is nc-supplemented in G, then $G \in \mathfrak{F}$.

Further definitions and notations are standard, please refer to [11] and [9].

2. Preliminaries

In this section, we give some concepts and some lemmas.

Definit on 2.1. ([3]) A subgroup H is said to be c-supplemented in group G if there exists a subgroup K such that HK = G and $H \cap K$ is contained in $Core_G(H)$. Then we say that K is a c-supplement of H in G.

Definition 2.2. ([12]) Let G be a group and H a subgroup of G. Then H is said to be nc-supplemented in G if there is a subgroup K of G such that $HK \leq G$ and $H \cap K \leq H_G$. We say that K is a nc-supplement of H in G. nc-supplemented subgroups

Remark 2.3. If H is a maximal subgroup of G, then an nc-supplement of H in G is a c-supplement of H in G.

Proof. If H is *nc*-supplemented in G, then there exists a subgroup K such that $HK \triangleleft G$ and $H \cap K \leq H_G$. The maximality of H implies that HK = G or HK = H. In the former case, H is *c*-supplemented in G. In the latter case, $H \triangleleft G$ and so H is also *c*-supplemented in G. \Box

Remark 2.4. Being nc-supplement is weaker than c-supplementation and normality.

nc-supplemented is a generalized *c*-supplemented. In general, *nc*-supplementation does not imply *c*-supplementation. For example (see [12, Example 3]), let $G = A_4$ and $B = \{(1), (12)(34), (13)(24), (14)(23)\}$. Let $C = \{(1), (12)(34)\}$ and $H = \{(1), (13)(24)\}$. Then $B = CH \leq G$ and *C* is *nc*-supplemented in *G* but not *c*-supplemented in *G* since $C_G = 1$ and *G* has no subgroup of order 6.

Lemma 2.5. ([12, Lemma 4]) If H is nc-supplemented in G, then there exists a subgroup C of G such that $H \cap C = H_G$ and $HC \leq G$.

Lemma 2.6. ([12, Lemma 5]) Let G be a group. Then

(1) If $H \leq M \leq G$ and H is nc-supplemented in G, then H is nc-supplemented in M.

(2) If $N \leq G$ and $N \leq H$, then H is nc-supplemented in G if and only if H/N is nc-supplemented in G/N.

(3) If $N \leq G$ and (|N|, |H|) = 1. If H is nc-supplemented in G, then HN/N is nc-supplemented in G/N.

Lemma 2.7. ([16, 2.16]) Let \mathfrak{F} be a formation containing \mathfrak{U} and let G be a group with a normal subgroup H such that $G/H \in \mathfrak{F}$. If H is cyclic, then $G \in \mathfrak{F}$.

3. Main results and their applications

In this section, we give the proofs of the main theorems. The proof of Theorem 1.1

Proof. Suppose that G is a counter-example of minimal order. We have: Step 1. Every proper subgroup M of G containing H is supersoluble and G is soluble.

Since $H \leq M$, it follows that M/H is a proper subgroup of G/H. Since G/H is supersoluble, it follows that M/H is supersoluble. Thus M satisfies the hypotheses of the theorem, and by the minimality of G, M is supersoluble. In particular, H is supersoluble and so G is soluble by [4].

Step 2. $\Phi(G) < H$ and $\Phi(G) = 1$.

Since the class of supersoluble group is a saturated formation by [14, 9.4.5], it is easy to get the result.

In the following, let L be a minimal normal subgroup of G contained in H. Then, by Step 1 and [10, Lemma 8. 6, p. 102] L is an elementary abelian p-group for some prime divisor p of |G|.

Step 3. G/L is supersoluble and L is the unique minimal normal subgroup of G which is contained in H.

First, we check that (G/L, H/L) satisfies the hypothesis as (G, H). Let $\overline{Q} = QL/L$ be a Sylow q-subgroup of $H/L = \overline{H}$. Then $\overline{G} = G/L$. Hence we assume that Q is a Sylow q-subgroup of H.

Case a. If p = q, we assume that L < P, then P = Q > L. Let P_1 be a maximal subgroup of P. By hypothesis P_1 is *nc*-supplemented in G, and by Lemma 2.6, $\overline{P_1}$ is *nc*-supplemented in \overline{G} . The minimality of G implies that \overline{G} is supersoluble.

Case b. Assume that $p \neq q$. Let $\overline{Q_1}$ be a maximal subgroup of a Sylow q-subgroup \overline{Q} of \overline{H} . Without loss of generality, we assume that $\overline{Q_1} = Q_1 L/L$. Since Q_1 is *nc*-supplemented in G, it follows, by Lemma 2.6, that $\overline{Q_1}$ is *nc*-supplemented in \overline{G} . The minimality of G implies that \overline{G} is supersoluble.

Now, let R be another minimal normal subgroup of G contained in H. Then G/R is supersoluble by Step 3. Since $G/R \cap L \leq G/R \times G/L$, it follows, from [1, Theorem 3] that, $G/R \cap L$ is supersoluble. On the other hand, $R \cap L \leq L$ and so $R \cap L = 1$ or $R \cap L = L$ by the minimality of L. In the former case, $G/1 \cong G$ is supersoluble, a contradiction. In the latter, L is unique.

Step 4. $L = F(H) = C_H(L)$.

Since L is an elementary abelian normal subgroup of $G, L \leq H$. So by [11, 6.5.4], F(H), the Fitting subgroup of H contains every minimal normal subgroup of H. By [6, Theorem 1.9.17] and Step 2, F(H) is the direct product of minimal normal subgroups of G contained in H. Then L = F(H) by Step 3. Since G is soluble by Step 1, $F(H) \leq C_H(L) =$ $C_H(F(H)) \leq F(H)$ by [19, Lemma 2.3].

Step 5. L is a Sylow subgroup of H.

nc-supplemented subgroups

Let q be the largest prime divisor of |H| and let Q be a Sylow qsubgroup of H. Since H/L is supersoluble, it follows, by [9, VI-9.1(c)], that LQ/L is characteristic in G/L and so $LQ \leq G$. Thus we have:

Case a. If p = q, then $L \leq P = Q \triangleleft G$. Therefore, by Step 1 and [4, Hilfssatz C], L = Q is a Sylow subgroup of H.

Case b. If p < q, then $L \leq P$ and PQ = PLQ is a subgroup of G. Since every maximal subgroup of all Sylow subgroups of PQ is *nc*-supplemented in PQ by Lemma 2.2(1), PQ satisfies the hypothesis of the theorem. Then we have:

Subcase a. If PQ < G, then, by Step 1, PQ is supersoluble and so $Q \triangleleft PQ$ by [9, VI-9.1]. Hence $LQ = L \times Q$ and so $Q \leq C_G(L) \leq L$ by [19, Lemma 2.3], a contradiction.

Subcase b. Assume that PQ = H = G and L < P in the case $Q \not \leq G$. Since $L \cap N_G(Q) = 1$ and LQ is characteristic in H = PQ = G, it follows that $G = [L]N_G(Q)$. Let P_2 be a Sylow *p*-subgroup of $N_G(Q)$. Then LP_2 is a Sylow *p*-subgroup of *G*. Choose a maximal subgroup P_1 of LP_2 with $P_2 \leq P_1$. Obviously, $L \leq P_1$ and $P_{1G} = 1$. Otherwise, $L = P_{1G}$, which contradicts that $L \cap N_G(Q) = 1$. By hypotheses, P_1 is *nc*-supplemented in G, then there exists a subgroup K such that $P_1K \triangleleft G$ and so $P_1 \cap K \leq P_{1G} = 1$. Hence if K is a q-subgroup of a Sylow q-subgroup Q of G, then P_1K is supersoluble by Step 1 and K is characteristic in P_1K which is normal in G. Then $LK = L \times K$ and so, by [19, Lemma 2.3], $K \leq C_G(L) \leq L$, a contradiction. Thus we assume that K is not a q-group. Since $|K|_p = |G : P_1|_p = p$, it follows that K has a normal p-complement Q^* . Obviously, P_1Q^* is a subgroup of G. By Step 1, P_1Q^* is supersoluble. And so, by [9, VI-9.1], $Q^* \triangleleft P_1 Q^*$. Thus $LQ^* = L \times Q^*$ and $Q^* \leq C_{P_1 Q^*}(L) \leq L$ by [19, Lemma 2.3], a contradiction. So we have $P_1K = G$. Now $|K|_p = |G: P_1|_p = p$ implies that K has a normal p-complement Q_1 which is also a Sylow q-subgroup of G. By [8, Theorem 4.2.2], there exists a $g \in LP_2 = P$ such that $Q_1^g = Q$. Since $P_1 \triangleleft P$, we have G = $P_1K = (P_1K)^g = P_1K^g$ and $P_1 \cap K^g = 1$. Since $K^g \cong K$ has a normal *p*-complement and $Q_1^g = Q \leq K^g$, it follows that $K^g \leq N_G(Q)$. Since
$$\begin{split} P &= LP = P LP = P LP \cap G = P (LP \cap K^g), \text{ if } P_1(LP_2 \cap K^g) \leq P_2, \\ \text{then } LP_2^2 \leq P_1^1 P_2^2 \leq P_2^1, \text{ a contradiction.} ^2\text{So } P_1(LP_2 \cap K^g) \nleq P_2 \text{ and} \end{split}$$
 P_2 must be a proper subgroup of $P_3 = \langle P_2, LP_2 \cap K^g \rangle$, where P_3 is a subgroup of a Sylow p-subgroup P. Thus P_2 and K^g are contained in $N_G(Q)$ and so P_3 is a p-subgroup of G containing a proper Sylow *p*-subgroup P_2 of $N_G(Q)$, a contradiction.

Thus L is a Sylow subgroup of H.

Step 6. |L| = p.

Let L_1 be a maximal subgroup of L. Then, by hypothesis, L_1 is *nc*supplemented in G and so, by Lemma 2.5, there exists a subgroup Kof G such that $L_1K \trianglelefteq G$ and $L_1 \cap K \le L_{1G}$. By Step 3, $L_1K \ge L$, and so $L = L \cap (L_1K) = L_1(L \cap K)$. It follows that $L \cap K = L$ or $L \cap K < L$. In the first case, it is easy to get $L \cap K \lhd G$. In the second case, $L_1 \cap K < L_1 < L$, and so $L_1 \cap K = L_1 \cap K \cap K < L \cap K < L$. Since $L_1 \cap K \lhd G$ and $L \lhd G$, it follows that $L(L_1 \cap K) \lhd G$. As $L(L_1 \cap K) =$ $(LL_1) \cap K = L \cap K$, we have $L \cap K \lhd G$ and so $L \cap K \ge L$ by the minimality and uniqueness of L. Then $L \cap K = L$ and so $L \le K$. Hence $L_1 \cap K \le L \cap K = L$ and so $L_1 \cap K = 1$. Thus $L_1 = 1$ and |L| = p. **Step 7.** The final contradiction.

By Step 3, G/L is supersoluble. By Step 6, L is a cyclic subgroup of prime order. Then by Lemma 2.7, G is supersoluble, a contradiction.

The final contradiction completes the proof.

Remark 3.1. The condition of Theorem 1.1 " G/H is supersoluble " cannot be replaced by " G/H is soluble ". Let $G = A_4 \times C_5$, where A_4 is the alternating group of degree 4 and C_5 is a cyclic group of order 5. Then $G/C_5 \cong A_4$ is soluble. Obviously, C_5 satisfies the hypotheses, but G is not supersoluble.

Corollary 3.2. ([3, Theorem 3.3]) Let G be a finite group and let N be a normal subgroup of G such that G/N is supersoluble. If every maximal subgroup of every Sylow subgroup of N is c-supplemented in G, then G is supersoluble.

Corollary 3.3. ([17, Theorem 1.1]) Let G be a finite group. Suppose P_1 is c-normal in G for every Sylow subgroup P of G and every maximal subgroup P_1 of P. Then G is supersoluble.

Corollary 3.4. ([2, Theorem 3.2]) Let G be a finite solvable group. Then G is supersoluble if and only if G/H is supersoluble and all maximal subgroups of every Sylow subgroup of F(H) are normal in G.

Corollary 3.5. ([15, Theorem 1]) Let G be a finite group such that all maximal subgroups of Sylow subgroups are normal in G. Then G is supersoluble.

Corollary 3.6. ([13, Theorem 3.5]) Assume that G/H is supersolvable and all maximal subgroups of the Sylow subgroups of H are normal in G. Then G is supersolvable.

The proof of the theorem 1.2

Proof. Assume that the theorem is false. And suppose that G is a counter-example of minimal order. By Lemma 2.6, we have that every maximal subgroup of the Sylow subgroups of H is *nc*-supplemented in H and so G is soluble. Then by [12, Theorem 11], H is soluble. We consider the following two cases:

Case 1. H is a p-group for some prime number p.

Step 1. Let N be the \mathfrak{F} -residual subgroup of G. Then $N = C_H(N) = F(H)$.

Let M be a nontrivial normal subgroup of G and let B be a maximal subgroup of MH with $M \leq B$. Then $B = M(H \cap B)$. Since $p = |MH | B| = |MH : M(H \cap B)| = |H : H \cap B|$, it follows that $H \cap B$ is a maximal subgroup of H. By hypothesis, $H \cap B$ is *nc*-supplemented in G and so is B. Thus B/M is *nc*-supplemented in G/M by Lemma 2.6(2). The minimal choice of G implies that $G/M \in \mathfrak{F}$. Since N is the \mathfrak{F} -residual subgroup of G, it follows that $\Phi(G) = 1$ and N is an elementary abelian subgroup of G since \mathfrak{F} is a saturated formation. Obviously $N \leq H$. Let F(H) be the Fitting subgroup of H. Then N = F(H) since \mathfrak{F} is a saturated formation. Then $F(H) \leq C_H(N) \leq N$ since H is solvable. Thus $N = C_H(N) = F(H)$ is a minimal normal nontrivial p-subgroup of G.

Step 2. H is a Sylow *p*-subgroup of G.

Suppose that H is not a Sylow p-subgroup of G and G is soluble. It follows, from [5, Theorem 3.5, p. 229], that there exists a Hall $\{p, q\}$ -subgroup of G, where q is a prime which is not equal to p, and that HQ is a subgroup of G since H is normal in the Sylow p-subgroup of G and $H \triangleleft G$. Since G/H is supersoluble, HQ/H is supersoluble . If HQ < G, then HQ is supersoluble and so is NQ. Then $N \cap Q = 1$, and $NQ = N \times Q$ since $N \triangleleft NQ$ and NQ is supersoluble. By [5, Theorem 1.3, p. 218], $Q \leq C_G(N) \leq N$, a contradiction. So H is a Sylow p-subgroup of G.

Step 3. |N| = p.

Let H_1 be a maximal subgroup of H. Then $N < H_1$. Otherwise, $N = H_1 \triangleleft G$, it follows, from [17, Theorem 1.1], that $G \in \mathfrak{F}$. H_1 is *nc*supplemented in G by hypothesis and so there exists a subgroup K of Gsuch that $H_1K \triangleleft G$ and $H_1 \cap K \leq H_{1G}$. Thus we have that $H_1 \cap K = 1$ or $H_1 \cap K = N$. If the former, $H_1K \geq H$ or $H_1K = H_1$ and so $K \geq H$ or $H_1 \geq K$, which contradicts $H_1 \cap K = 1$. Hence $N \leq K$ and N is a Sylow p-subgroup of K. If N is not a Sylow p-subgroup of K, then there is a Sylow p-subgroup P_K of G with $N < P_K$, and so $H_1P_K = H$ or $H_1P_K = H_1$. In the former case, $P_K = H$ and so $H_1 \cap K = H_1 \cap H = H_1 \triangleleft G$. It follows, from [13, Theorem 3.5], that G is supersoluble, a contradiction. In the latter, $N < P_K \leq H_1$ and so $N = H_1 \cap K = H_1 \cap P_K = P_K > N$, another contradiction. Thus N is a normal Sylow p-subgroup of K. By Step 2, K < G and so HK < G. Since HK/H is supersoluble and every maximal subgroup of H is nc-supplemented in HK, it follows, from the minimal choice of G that, HK is supersoluble and so K is supersoluble. Let Q be a Sylow q-subgroup of K, where q is the largest prime of |K|. Thus Q is normal in K, and $NQ = N \times Q$. This means $Q \leq C_K(N) \leq N$, a contradiction. Hence there does not exist non-trivial maximal subgroup of H, that is, H is a Sylow p-subgroup of G of order p. Namely, |H| = |N| = p.

Step 4. The final contradiction.

By Step 3, H is a cyclic subgroup. By Lemma 2.7, $\overline{G} \in \mathfrak{F}$, a contradiction.

Case 2. *H* is not of prime power order.

Let P be a Sylow p-subgroup of H. Then by hypothesis and Lemma 2.6(1), the maximal subgroups of every Sylow subgroup of H are nc-supplemented in H. Then by Theorem 1.1, H is supersoluble, and so by [4, Hillssatz C] H has a normal Sylow subgroup P

Since P is characteristic in H and $H \triangleleft G$, it follows that $P \triangleleft G$. Clearly, $(G/P)/(H/P) \cong G/H \in \mathfrak{F}$. By the minimality of $G, G/P \in \mathfrak{F}$. But now $G \in \mathfrak{F}$ by Case 1, a contradiction.

So the minimal counter-example does not exist.

This completes the proof.

Remark 3.7. The condition of Theorem 1.2, " \mathfrak{U} " cannot be replaced by " \mathfrak{N} ", where \mathfrak{N} is the class of all nilpotent groups. Let $G = S_3$ the symmetric group of degree 3. Then G is supersoluble, but G not nilpotent.

Corollary 3.8. ([18, Theorem 1]) Let \mathfrak{F} be a saturated formation containing \mathfrak{U} . Suppose that G is a group with a soluble normal subgroup H such that $G/H \in \mathfrak{F}$. If all maximal subgroups of all Sylow subgroups of F(H) are c-normal in G, then $G \in \mathfrak{F}$.

Corollary 3.9. ([19, Theorem 3.1]) Let \mathfrak{F} be a saturated formation containing \mathfrak{U} . Suppose that G is a group with a normal subgroup H such

nc-supplemented subgroups

that $G/H \in \mathfrak{F}$. If all maximal subgroups of all Sylow subgroups of $F^*(H)$ are c-normal in G, then $G \in \mathfrak{F}$.

Corollary 3.10. ([20, Theorem 1.2]) Let \mathfrak{F} be a saturated formation containing \mathfrak{U} . Suppose that G is a group G with a normal subgroup H such that $G/H \in \mathfrak{F}$. If all maximal subgroups of all Sylow subgroups of $F^*(H)$ are c-supplemented in G, then $G \in \mathfrak{F}$.

Acknowledgments

The authors would like to thank the referee with deep gratitude for pointing out some mistakes in a previous version of the paper, especially his/her valuable suggestions on revising the paper, which made the proof of theorems read smoothly and with technical support. The third author is supported by NSF of China and the subject is partially supported by NSF of SUSE (Grant Number: 2010XJKYL017) and Scientific Research Fund of School of Science of SUSE (Grant Number: 09LXYB02).

References

- M. Asaad, On the supersolvability of finite groups I, Acta Math. Acad. Sci. Hungar. 38 (1981), no. 1-4, 57–59.
- [2] M. Asaad, M. Ramadan and A. Shaalan, Influence of π-quasinormality on maximal subgroups of Sylow subgroups of Fitting subgroup of a finite group, Arch. Math. 56 (1991), no. 6, 521–527.
- [3] A. Ballester-Bolinches, Y. Wang and X. Guo, c-supplemented subgroups of finite groups, Glasg. Math. J. 42(2000), no. 3, 383–389.
- [4] K. Doerk, Minimal nicht überauflösbare, endliche gruppen, Math. Z. 91 (1966) 198–205.
- [5] D. Gorenstein, Finite Groups, Second Edition, Chelsea Publishing Co., New York, 1980.
- [6] W. Guo, The Theory of Classes of Groups, Kluwer Academic Publishers Group, Dordrecht, Science Press, Beijing, 2000.
- [7] P. Hall, A characteristic property of soluble groups, J. London Math. Soc. 12 (1937), no. 2, 198–200
- [8] M. Hall, The Theory of Groups, The Macmillan Co., New York, 1959.
- [9] B. Huppert, Endliche Gruppen I, Springer-Verlag, Berlin-Heidelberg-New York, 1967.
- [10] I. M. Isaacs, Algebra: A Graduate Course, Brooks/Cole Publishing Co, Pacific Grove, 1994.

- [11] H. Kurzweil and B. Stellmacher, The Theory of Finite Groups: An Introduction, Springer-Verlag, New York, 2004.
- [12] S. Li, D. Liang and W. Shi, A generalization of c-supplementation, Southeast Asian Bull. Math. 30 (2006), no. 5, 889–895.
- [13] M. Ramadan, Influence of normality on maximal subgroups of Sylow subgroups of a finite group, Acta Math. Hungar. 59 (1992), no. 1-2, 107–110.
- [14] D. J. Robinson, A Course in the Theory of Groups, 2nd Edition, Springer-Verlag, New York, 1995.
- S. Srinivasan, Two sufficient conditions for supersolvability of finite groups, *Israel J. Math.* 35 (1980), no. 3, 210–214.
- [16] A. N. Skiba, On weakly s-permutable subgroups of finite groups, J. Algebra 315 (2007), no. 1, 192–209.
- [17] Y. Wang, c-normality of groups and its properties, J. Algebra 180 (1996), no. 3, 954–965.
- [18] H. Wei, On c-normal maximal and minimal subgroups of Sylow subgroups of finite groups, Comm. Algebra 29 (2001), no. 5, 2193–2200.
- [19] H. Wei, Y. Wang and Y. Li, On *c*-normal maximal and minimal subgroups of Sylow subgroups of finite groups II, *Comm. Algebra* **31** (2003), no. 10, 4807–4816.
- [20] H. Wei, Y. Wang and Y. Li, On c-supplemented maximal and minimal subgroups of Sylow subgroups of finite groups, Proc. Amer. Math. Soc. 132 (2004), no. 8, 2197–2204.

Shiguang Guo

School of Science, Sichuan University of Science & Engineering, 643000, Zigong, P. R. China

Email: 710442986@qq.com

Shitian Liu

School of Science, Sichuan University of Science & Engineering, 643000, Zigong, P. R. China

Email: liust@suse.edu.cn

Wujie Shi

School of Mathematics and Statistics, Chongqing University of Arts and Sciences, 402160, Chongqing, P. R. China Email: wjshi@suda.edu.cn