

IMPLICIT ITERATION APPROXIMATION FOR A FINITE FAMILY OF ASYMPTOTICALLY QUASI-PSEUDOCONTRACTIVE TYPE MAPPINGS

S. ZHANG

(Communicated by Behzad Djafari-Rouhani)

ABSTRACT. In this paper, strong convergence theorems of Ishikawa type implicit iteration process with errors for a finite family of asymptotically nonexpansive in the intermediate sense and asymptotically quasi-pseudocontractive type mappings in normed linear spaces are established by using a new analytical method, which essentially improve and extend some recent results obtained by Yang [Convergence theorems of implicit iteration process for asymptotically pseudocontractive mappings, Bulletin of the Iranian Mathematical Society, Available Online from 12 April 2011] and others.

Keywords: Normed linear spaces, implicit iteration process, asymptotically quasi-pseudocontractive type mappings, nonexpansive mappings.

MSC(2010): Primary: 47H05; Secondary: 47H10, 47J25.

1. Introduction

Let E be an arbitrary real normed linear space with norm $\|\cdot\|$ and E^* be the duality space of E . Let $\langle \cdot, \cdot \rangle$ denote the duality pairing between E and E^* . For $1 < p < \infty$, the mapping $J_p : E \rightarrow 2^{E^*}$ defined by

$$J_p(x) = \{f \in E^* : \langle x, f \rangle = \|x\| \cdot \|f\|, \|f\| = \|x\|^{p-1}\}$$

is called the duality mapping with gauge function $\varphi(t) = t^{p-1}$. In particular, for $p = 2$, the duality mapping J_2 with gauge function $\varphi(t) = t$ is called the normalized duality mapping. It is well known that the duality

Article electronically published on February 25, 2014.

Received: 6 December 2011, Accepted: 28 January 2013.

©2014 Iranian Mathematical Society

mapping J_p has the following properties:

- (i) $J_p(x) = \|x\|^{p-2} J_2(x)$ for all $x \in E (x \neq 0)$,
- (ii) $J_p(\alpha x) = \alpha^{p-1} J_p(x)$ for all $\alpha \geq 0$,
- (iii) J_p can be equivalently defined as the subdifferential of the functional $\psi(x) = p^{-1} \|x\|^p$, i.e., $J_p(x) = \partial\psi(x) = \{f \in E^* : \psi(y) - \psi(x) \geq \langle y - x, f \rangle, \forall y \in E\}$ (Asplund [1]).

Definition 1.1. Let K be a nonempty subset of E . A mapping $T : K \rightarrow K$ is said to be

- (i) asymptotically nonexpansive, if there exists a sequence $\{k_n\} \subset [1, \infty)$ with $\lim_{n \rightarrow \infty} k_n = 1$, such that

$$\|T^n x - T^n y\| \leq k_n \|x - y\|, \quad \forall x, y \in K, n \geq 1,$$

- (ii) asymptotically pseudo-contractive, if for all $x, y \in K$, there exist $j(x - y) \in J(x - y)$ and a sequence $\{k_n\} \subset [1, \infty)$ with $\lim_{n \rightarrow \infty} k_n = 1$, such that

$$\langle T^n x - T^n y, j(x - y) \rangle \leq k_n \|x - y\|^2, \quad n \geq 1,$$

- (iii) asymptotically quasi-pseudocontractive type if $F(T) \neq \emptyset$ and there exists a sequence $\{k_n\} \subset [1, \infty)$ with $\lim_{n \rightarrow \infty} k_n = 1$, such that

$$\limsup_{n \rightarrow \infty} \left\{ \sup_{x \in K} \inf_{j_p(x - x^*) \in J_p(x - x^*)} \langle T^n x - x^*, j_p(x - x^*) \rangle - k_n \|x - x^*\|^p \right\} \leq 0,$$

- (iv) asymptotically nonexpansive in the intermediate sense if

$$\limsup_{n \rightarrow \infty} \left\{ \sup_{x, y \in K} (\|T^n x - T^n y\| - \|x - y\|) \right\} \leq 0.$$

It is easy to see that an asymptotically nonexpansive mapping is asymptotically nonexpansive in the intermediate sense if the domain of T is bounded. Every asymptotically nonexpansive mapping is asymptotically pseudocontractive, and every asymptotically pseudocontractive mapping is asymptotically quasi-pseudocontractive type mapping. But the inverse is not true, in general.

The concept of asymptotically nonexpansive mappings was introduced by Goebel and Kirk [5], while the concept of asymptotically pseudocontractive mapping was introduced by Schu [12] in 1991. The iterative approximation problems for asymptotically nonexpansive mappings and asymptotically pseudo-contractive mappings were studied extensively by

Schu [12], Chang [3], Khan et al. [7], Ofoedu [8], Plubtieng et al [10], Xu and Ori [14], Zhou [19], Sun [13], Yang and Hu [15] and Yang [16] in the setting of Hilbert spaces or Banach spaces.

Let K be a nonempty closed convex subset of E and $\{T_i\}_{i=1}^m$ be a finite family of nonexpansive mappings from K into itself (i.e., $\|T_i x - T_i y\| \leq \|x - y\|$ for $x, y \in K$ and $i = 1, 2, \dots, m$). In 2001, Xu and Ori [14] introduced the following implicit iteration process. For an arbitrary $x_0 \in K$ and $\alpha_n \in [0, 1]$, the sequence $\{x_n\}$ is generated as follows:

$$\begin{cases} x_1 = (1 - \alpha_1)x_0 + \alpha_1 T_1 x_1, \\ x_2 = (1 - \alpha_2)x_1 + \alpha_2 T_2 x_2, \\ \vdots \\ x_N = (1 - \alpha_N)x_{N-1} + \alpha_N T_N x_N, \\ x_{N+1} = (1 - \alpha_{N+1})x_N + \alpha_{N+1} T_{N+1} x_{N+1}, \\ \vdots \end{cases}$$

The scheme is expressed in its compact form by

$$x_n = (1 - \alpha_n)x_n + \alpha_n T_{n(\text{mod } N)} x_n, n \geq 1.$$

Using this iteration, they proved that the sequence $\{x_n\}$ converges weakly to a common fixed point of a finite family of nonexpansive mappings $\{T_i\}_{i=1}^m$ in a Hilbert space under certain conditions.

In 2006, Chang et al.[3] introduced another implicit iteration process with error. In the sense of [3], the implicit iteration process with errors for a finite family of asymptotically nonexpansive mappings $\{T_i\}_{i=1}^m$ is generated from an arbitrary $x_0 \in K$ by

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_{i(n)}^{k(n)} x_n + u_n, n \geq 1,$$

where $n = (k - 1)m + i, i = i(n) \in \{1, 2, \dots, m\}, k = k(n) \geq 1$ is a positive integer and $k(n) \rightarrow \infty$, as $n \rightarrow \infty$. $\{\alpha_n\}$ is a suitable sequence in $[0, 1]$ and $\{u_n\} \subset K$ is such that $\sum_{n=0}^{\infty} \|u_n\| < \infty$. They extended the results of [14] from Hilbert spaces to more general uniformly convex Banach spaces and from nonexpansive mappings to asymptotically nonexpansive mappings.

Yang and Hu [15] proposed another implicit iteration process which appears to be more satisfactory as follows:

$$(1.1) \quad x_n = \alpha_n x_{n-1} + \beta_n T_{i(n)}^{k(n)} x_n + \gamma_n u_n, n \geq 1,$$

where $\{\alpha_n\}, \{\beta_n\}$ and $\{\gamma_n\} \subset [0, 1]$ with $\alpha_n + \beta_n + \gamma_n = 1$, and $\{u_n\}$ is bounded in K .

Since for each $n \geq 1$, it can be written as $n = (k - 1)m + i$, where $i = i(n) \in \{1, 2, \dots, m\}$, $k = k(n) - 1$ is a positive integer and $k(n) \rightarrow \infty$ as $n \rightarrow \infty$. Hence, (1.1) can be expressed in the following form:

$$(1.2) \quad x_n = (1 - \alpha_n - \gamma_n)x_{n-1} + \alpha_n T_{i(n)}^n x_n + \gamma_n u_n, n \geq 1,$$

where $\alpha_n + \gamma_n \leq 1$, and $\{u_n\}$ is bounded in K .

Very recently, Yang [16] proved the following result.

Theorem 1.2. ([16]). *Let E be a real normed linear space, K be a nonempty convex subset of E , $T_i : K \rightarrow K, i = 1, 2, \dots, m$ be a finite family of asymptotically nonexpansive in the intermediate sense and asymptotically pseudo-contractive mappings with $\{k_{in}\} \subset [1, \infty)$ such that $\lim_{n \rightarrow \infty} k_n = 1$, where $k_n = \max_{1 \leq i \leq m} \{k_{in}\}$. Assume that $F = \bigcap_{i=1}^m F(T_i) \neq \emptyset$ denotes the set of common fixed points of $\{T_i\}_{i=1}^m$. Let $\{x_n\}$ be the sequence defined by (1.2). Suppose that $\{u_n\}$ is bounded in K and that $\{\alpha_n\}, \{\gamma_n\}$ are sequences in $[0, 1]$ satisfying the following conditions:*

- (i) $\sum_{n=0}^{\infty} \gamma_n < \infty, \sum_{n=0}^{\infty} \alpha_n(k_n - 1) < \infty$,
- (ii) $\sum_{n=0}^{\infty} \alpha_n = \infty, \lim_{n \rightarrow \infty} \alpha_n = 0$.

Assume that there exists a strict increasing function $\varphi : [0, \infty) \rightarrow [0, \infty)$ with $\varphi(0) = 0$ such that

$$\limsup_{n \rightarrow \infty} \left\{ \langle T_i^n x_n - x^*, j(x_n - x^*) \rangle - k_n \|x_n - x^*\|^2 + \varphi(\|x_n - x^*\|) \right\} \leq 0$$

for $x^ \in F$ and $i = 1, 2, \dots, m$. Then $\{x_n\}$ converges strongly to a common fixed point p of $\{T_i\}_{i=1}^m$.*

Remark 1.3. *We point out here that the conditions (i) is not always true.*

Example 1.4. *Let $\alpha_n = \frac{1}{\sqrt{n+1}}$ and $k_n = 1 + \frac{1}{\sqrt{n+1}}$, then $\sum_{n=0}^{\infty} \alpha_n(k_n - 1) = \sum_{n=0}^{\infty} \frac{1}{n+1} = \infty$, which show that conditions (i) in Theorem 1.2 is not satisfied. Hence Theorem 1.2 need to be improved.*

The purpose of this paper is, under the condition of removing the restriction $\sum_{n=0}^{\infty} \alpha_n(k_n - 1) < \infty$, to prove strong convergence theorems of Ishikawa type implicit iteration process with errors for a finite family of

asymptotically nonexpansive in the intermediate sense and asymptotically quasi-pseudocontractive type mappings in normed linear spaces by using a new analytical method. Our results essentially extend and improve some recent results obtained by Yang [16] and others.

Now we consider Ishikawa type implicit iteration process with errors for a finite family of asymptotically quasi-pseudocontractive type mappings as follows:

$$(1.3) \quad \begin{cases} x_n = (1 - \alpha_n - \gamma_n)x_{n-1} + \alpha_n T_{i(n)}^n y_n + \gamma_n u_n \\ y_n = (1 - \beta_n - \mu_n)x_n + \beta_n T_{i(n)}^n x_n + \mu_n v_n, \quad (n \geq 0), \end{cases}$$

where $n = (k-1)m + i, i = i(n) \in \{1, 2, \dots, m\}, k = k(n) \geq 1$ is a positive integer and $k(n) \rightarrow \infty$, as $n \rightarrow \infty$. $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\mu_n\}$ are four suitable sequences in $[0, 1]$ with $\alpha_n + \gamma_n \leq 1, \beta_n + \mu_n \leq 1$ and $\{u_n\}, \{v_n\}$ are bounded sequences in K .

The following lemmas plays an important role in this paper.

Lemma 1.5 ([17]). / Let E be a real normed linear space and $J_p : E \rightarrow 2^{E^*}$ a duality mapping. Then

$$\|x + y\|^p \leq \|x\|^p + p\langle y, j_p(x + y) \rangle$$

for all $x, y \in E, 1 < p < \infty$ and $j_p(x + y) \in J_p(x + y)$.

Lemma 1.6. Let $\varphi_i (i = 1, 2, \dots, m) : [0, \infty) \rightarrow [0, \infty)$ be strictly increasing functions with $\varphi_i(0) = 0$ and let $\{a_n\}, \{b_n\}, \{c_n\}, \{\delta_n\}$ be non-negative real sequences such that $\sum_{n=1}^{\infty} \delta_n = \infty, \lim_{n \rightarrow \infty} \frac{b_n}{\delta_n} = 0, \sum_{n=1}^{\infty} c_n < \infty$. Suppose that

$$(1.4) \quad a_{n+1}^p \leq a_n^p - \delta_n \varphi_i(a_{n+1}) + b_n + c_n, \quad n \geq n_0,$$

where n_0 is some nonnegative integer and $p \in (1, \infty)$, then $\lim_{n \rightarrow \infty} a_n = 0$.

Proof. Setting $\liminf_{n \rightarrow \infty} a_n = \tau$, then $\tau \geq 0$. Now we prove $\tau = 0$. If $\tau > 0$, then there exists a positive integer $N_1 > 0$ such that $a_n \geq \frac{\tau}{2}$ for all $n \geq N_1$. By the strictly increasing property of φ_i , we have $\varphi_i(a_{n+1}) > \varphi_i(\frac{\tau}{2}) \geq \min_{1 \leq i \leq m} \varphi_i(\frac{\tau}{2}) =: \sigma$. Since $\lim_{n \rightarrow \infty} \frac{b_n}{\delta_n} = 0$, there exists a positive integer $N_2 > N_1$ such that $\frac{b_n}{\delta_n} \leq \frac{1}{2}\sigma$ for all $n \geq N_2$. Taking $N_3 = \max\{N_2, n_0\}$, then from (1.4), we have

$$a_{n+1}^p \leq a_n^p - \delta_n \sigma + \delta_n \frac{\sigma}{2} + c_n = a_n^p - \delta_n \frac{\sigma}{2} + c_n$$

for all $n \geq N_3$, which means that $\delta_n \frac{\sigma}{2} \leq a_n^p - a_{n+1}^p + c_n$. Hence for any positive integer $h \geq N_3$, we obtain

$$\frac{\sigma}{2} \sum_{n=N_3}^h \delta_n \leq a_{N_3}^p - a_{h+1}^p + \sum_{n=N_3}^h c_n \leq a_{N_3}^p + \sum_{n=N_3}^h c_n,$$

and so

$$\infty = \frac{\sigma}{2} \sum_{n=N_3}^{\infty} \delta_n \leq a_{N_3}^p + \sum_{n=N_3}^{\infty} c_n,$$

a contradiction. This implies that $\tau > 0$ is impossible. Therefore $\tau = 0$, which there exists a subsequence $\{a_{n_j}\} \subset \{a_n\}$ such that $a_{n_j} \rightarrow 0$ ($j \rightarrow \infty$). Since $\lim_{n \rightarrow \infty} \frac{b_n}{\delta_n} = 0$, $\sum_{n=1}^{\infty} c_n < \infty$, for any given $\varepsilon > 0$, there exist two positive integers $j_0 > 0$ and $N_4 > 0$, such that for all $n \geq N_4$, $\sum_{n=N_4}^{\infty} c_n < \varepsilon^p$, $\frac{b_n}{\delta_n} < \frac{1}{2}\omega$ and $a_{n_j} < \varepsilon$ for all $j \geq j_0$, where $\omega = \min\{\varphi_1(\varepsilon), \varphi_2(\varepsilon), \dots, \varphi_m(\varepsilon)\}$. Let $N_5 = \max\{j_0, N_4\}$. For fixed $j_* > N_5$ and all $k \geq 0$, we now want to show that $a_{n_{j_*}+k} < 2\varepsilon$. To see this consider two possible cases.

Case I: $a_{n_{j_*}+1} < \varepsilon$.

In this case, $a_{n_{j_*}+1}^p < \varepsilon^p + c_{n_{j_*}} + c_{n_{j_*}+1}$ and so we have the desired result.

Case II: $a_{n_{j_*}+1} \geq \varepsilon$.

In this case, $\varphi_i(a_{n_{j_*}+1}) \geq \varphi_i(\varepsilon) \geq \omega > 0$ since for each $i = 1, 2, \dots, m$, φ_i is a strictly increasing function. From (1.4), we also have

$$\begin{aligned} a_{n_{j_*}+1}^p &\leq a_{n_{j_*}}^p - \delta_{n_{j_*}} \varphi_i(a_{n_{j_*}+1}) + b_{n_{j_*}} + c_{n_{j_*}} \\ &\leq a_{n_{j_*}}^p - \delta_{n_{j_*}} \left(\omega - \frac{\omega}{2} \right) + c_{n_{j_*}} \\ &< \varepsilon^p + c_{n_{j_*}} + c_{n_{j_*}+1}. \end{aligned}$$

By using induction, we have

$$a_{n_{j_*}+k}^p < \varepsilon^p + \sum_{i=n_{j_*}}^{n_{j_*}+k} c_i < \varepsilon^p + \varepsilon^p = 2\varepsilon^p < (2\varepsilon)^p$$

for all $k \geq 0$. This shows $a_n \rightarrow 0$ as $n \rightarrow \infty$. The proof of Lemma 1.6 is completed.

2. Main results

Theorem 2.1. Let E be a real normed linear space, K a nonempty convex subset of E and $T_i : K \rightarrow K$ ($i = 1, 2, \dots, m$) a finite family of asymptotically nonexpansive in the intermediate sense and asymptotically quasi-pseudocontractive type mappings with $\{k_n^{(i)}\} \subset [1, \infty)$ such that $\lim_{n \rightarrow \infty} k_n = 1$, where $k_n = \max_{1 \leq i \leq m} \{k_n^{(i)}\}$. Assume that $F = \bigcap_{i=1}^m F(T_i) \neq \emptyset$ denotes the set of common fixed points of $\{T_i\}_{i=1}^m$. Let $\{x_n\}$ be the sequence defined by (1.3). Suppose that $\{u_n\}$ and $\{v_n\}$ are bounded sequences in K and that $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$ and $\{\mu_n\}$ are sequences in $[0, 1]$ satisfying the following conditions:

- (i) $\sum_{n=0}^{\infty} \alpha_n = \infty$,
- (ii) $\alpha_n \rightarrow 0, \beta_n \rightarrow 0$ ($n \rightarrow \infty$),
- (iii) $\sum_{n=0}^{\infty} \gamma_n < \infty, \mu_n \rightarrow 0$ ($n \rightarrow \infty$).

Assume that there exist strict increasing functions $\varphi_i : [0, \infty) \rightarrow [0, \infty)$ with $\varphi_i(0) = 0$ such that

$$(2.1) \quad \limsup_{n \rightarrow \infty} \left\{ \inf_{j_p(x_n - x^*) \in J_p(x_n - x^*)} \langle T_i^n x_n - x^*, j_p(x_n - x^*) \rangle - k_n \|x_n - x^*\|^p + \varphi_i(\|x_n - x^*\|) \right\} \leq 0$$

for $x^* \in F$ and $i = 1, 2, \dots, m$. Then $\{x_n\}$ converges strongly to a common fixed point x^* of $\{T_i\}_{i=1}^m$.

Proof. For $i = 1, 2, \dots, m$, let

$$\sigma_n^{(i)} = \inf_{j_p(x_n - x^*) \in J_p(x_n - x^*)} \langle T_i^n x_n - x^*, j_p(x_n - x^*) \rangle - k_n \|x_n - x^*\|^p + \varphi_i(\|x_n - x^*\|),$$

then there exist $j_p^{(i)}(x_n - x^*) \in J_p(x_n - x^*)$, such that

$$(2.2) \quad \begin{aligned} \langle T_i^n x_n - x^*, j_p^{(i)}(x_n - x^*) \rangle &= k_n \|x_n - x^*\|^p + \varphi_i(\|x_n - x^*\|) \\ &< \sigma_n^{(i)} + \varepsilon_n^{(i)} \leq \xi_n, \end{aligned}$$

where $\varepsilon_n^{(i)} \in (0, 1)$ with $\varepsilon_n^{(i)} \rightarrow 0$ ($n \rightarrow \infty$), and $\xi_n = \max_{1 \leq i \leq m} \{\sigma_n^{(i)}, 0\} +$

$\max_{1 \leq i \leq m} \{\varepsilon_n^{(i)}\}$. It is easy to see (using (2.1)) that $\lim_{n \rightarrow \infty} \xi_n = 0$. Since $\{u_n\}$ and $\{v_n\}$ are bounded sequences in K , $M = \sup_{n \geq 0} \{\|u_n - x^*\| + \|v_n - x^*\|\} < \infty$. Also, since for each $i = 1, 2, \dots, m$, $T_i : K \rightarrow K$ is an asymptotically

nonexpansive in the intermediate sense, there exists $n_0 \geq 1$ such that $\sup_{x,y \in K} (\|T_i^n x - T_i^n y\| - \|x - y\|) \leq 1$ for all $n \geq n_0, i = 1, 2, \dots, m$.

It follows from (1.3) that

$$\begin{aligned}
 & \frac{\|x_n - x^*\|}{1 + \|x_{n-1} - x^*\|} \\
 = & \frac{\|(1 - \alpha_n - \gamma_n)(x_{n-1} - x^*) + \alpha_n(T_i^n y_n - x^*) + \gamma_n(u_n - x^*)\|}{1 + \|x_{n-1} - x^*\|} \\
 \leq & \frac{\|x_{n-1} - x^*\| + \alpha_n\|T_i^n y_n - x^*\| + \|u_n - x^*\|}{1 + \|x_{n-1} - x^*\|} \\
 \leq & \frac{\|x_{n-1} - x^*\| + \sup_{x,y \in K} (\|T_i^n x - T_i^n y\| - \|x - y\|)}{1 + \|x_{n-1} - x^*\|} \\
 + & \frac{\alpha_n\|y_n - x^*\| + \|u_n - x^*\|}{1 + \|x_{n-1} - x^*\|} \\
 \leq & \frac{\|x_{n-1} - x^*\| + 1 + \alpha_n[\|x_n - x^*\| + \|T_i^n x_n - x^*\|]}{1 + \|x_{n-1} - x^*\|} \\
 + & \frac{\|v_n - x^*\| + \|u_n - x^*\|}{1 + \|x_{n-1} - x^*\|} \\
 \leq & \frac{\|x_{n-1} - x^*\| + 1 + 2\alpha_n\|x_n - x^*\|}{1 + \|x_{n-1} - x^*\|} \\
 + & \frac{\sup_{x,y \in K} (\|T_i^n x - T_i^n y\| - \|x - y\|) + 2M}{1 + \|x_{n-1} - x^*\|} \\
 (2.3) \leq & 3 + 2M + \frac{2\alpha_n\|x_n - x^*\|}{1 + \|x_{n-1} - x^*\|}
 \end{aligned}$$

for all $n \geq n_0$.

Since $1 - 2\alpha_n \rightarrow 1$ ($n \rightarrow \infty$), there exists $n_1 \geq n_0$ such that $1 - 2\alpha_n > \frac{1}{2} > 0$ for all $n \geq n_1$, which together with (2.3) gives that

$$(2.4) \quad \frac{\|x_n - x^*\|}{1 + \|x_{n-1} - x^*\|} \leq \frac{3 + 2M}{1 - 2\alpha_n} \leq 6 + 4M.$$

Let $c_n^{(i)} = \sup_{x,y \in K} (\|T_i^n x - T_i^n y\| - \|x - y\|)$, $d_n = \max \left\{ 0, \max_{1 \leq i \leq m} c_n^{(i)} \right\}$, then $\lim_{n \rightarrow \infty} d_n = 0$.

By (1.3), we have

$$\begin{aligned}
 \|x_n - y_n\| &\leq \beta_n \|T_i^n x_n - x_n\| + \mu_n \|v_n - x_n\| \\
 &\leq \beta_n (\|T_i^n x_n - x^*\| - \|x_n - x^*\|) \\
 &\quad + (2\beta_n + \mu_n) \|x_n - x^*\| + \mu_n \|v_n - x^*\| \\
 &\leq \beta_n \sup_{x, y \in K} (\|T_i^n x - T_i^n y\| - \|x - y\|) \\
 &\quad + (2\beta_n + \mu_n) \|x_n - x^*\| + \mu_n M \\
 (2.5) \quad &\leq (2\beta_n + \mu_n) \|x_n - x^*\| + \beta_n d_n + \mu_n M
 \end{aligned}$$

for all $n \geq n_1$.

For $j_p^{(i)}(x_n - x^*) \in J_p(x_n - x^*)$, $\forall n \geq 0$, we have from (1.3) and Lemma 1.5 that

$$\begin{aligned}
 &\left(\frac{\|x_n - x^*\|}{1 + \|x_{n-1} - x^*\|} \right)^p \\
 = &\frac{\|(1 - \alpha_n - \gamma_n)(x_{n-1} - x^*) + \alpha_n(T_i^n y_n - x^*) + \gamma_n(u_n - x^*)\|^p}{(1 + \|x_{n-1} - x^*\|)^p} \\
 \leq &\frac{(1 - \alpha_n)^p \|x_{n-1} - x^*\|^p + p\alpha_n \langle T_i^n x_n - x^*, j_p^{(i)}(x_n - x^*) \rangle}{(1 + \|x_{n-1} - x^*\|)^p} \\
 &+ \frac{p\alpha_n \langle T_i^n y_n - T_i^n x_n, j_p^{(i)}(x_n - x^*) \rangle}{(1 + \|x_{n-1} - x^*\|)^p} \\
 (2.6) + &\frac{p\gamma_n \langle u_n - x^*, j_p^{(i)}(x_n - x^*) \rangle}{(1 + \|x_{n-1} - x^*\|)^p}
 \end{aligned}$$

for all $n \geq n_1, i = 1, 2, \dots, m$.

Next we consider the second and third term on the right side of (2.6). From (2.4) and (2.5), we obtain that

$$\begin{aligned}
 &\frac{p\alpha_n \langle T_i^n y_n - T_i^n x_n, j_p^{(i)}(x_n - x^*) \rangle}{(1 + \|x_{n-1} - x^*\|)^p} \\
 \leq &p\alpha_n \frac{\|T_i^n y_n - T_i^n x_n\| \|x_n - x^*\|^{p-1}}{(1 + \|x_{n-1} - x^*\|)^p} \\
 = &p\alpha_n \frac{(\|T_i^n x_n - T_i^n y_n\| - \|x_n - y_n\|) + \|x_n - y_n\|}{1 + \|x_{n-1} - x^*\|} \\
 &\cdot \left(\frac{\|x_n - x^*\|}{1 + \|x_{n-1} - x^*\|} \right)^{p-1}
 \end{aligned}$$

$$\begin{aligned}
&\leq p\alpha_n \left(\frac{d_n}{1 + \|x_{n-1} - x^*\|} + \frac{\|x_n - y_n\|}{1 + \|x_{n-1} - x^*\|} \right) \\
&\quad \cdot \left(\frac{\|x_n - x^*\|}{1 + \|x_{n-1} - x^*\|} \right)^{p-1} \\
&\leq p\alpha_n \left(d_n + \frac{(2\beta_n + \mu_n)\|x_n - x^*\| + \beta_n d_n + \mu_n M}{1 + \|x_{n-1} - x^*\|} \right) \\
&\quad \cdot \left(\frac{\|x_n - x^*\|}{1 + \|x_{n-1} - x^*\|} \right)^{p-1} \\
(2.7) \quad &\leq p\alpha_n (6 + 4M)^{p-1} [d_n + (2\beta_n + \mu_n)(6 + 4M) + \beta_n d_n + \mu_n M]
\end{aligned}$$

for all $n \geq n_1$.

In view of (2.4), we deduce that

$$\begin{aligned}
\frac{p\gamma_n \langle u_n - x^*, j_p^{(i)}(x_n - x^*) \rangle}{(1 + \|x_{n-1} - x^*\|)^p} &\leq \frac{p\gamma_n \|u_n - x^*\| \|x_n - x^*\|^{p-1}}{(1 + \|x_{n-1} - x^*\|)^p} \\
(2.8) \quad &\leq p\gamma_n M (6 + 4M)^{p-1}.
\end{aligned}$$

Substituting (2.2), (2.7) and (2.8) into (2.6) yields that

$$\begin{aligned}
\left(\frac{\|x_n - x^*\|}{1 + \|x_{n-1} - x^*\|} \right)^p &\leq \frac{(1 - \alpha_n)^p \|x_{n-1} - x^*\|^p + p\alpha_n \xi_n}{(1 + \|x_{n-1} - x^*\|)^p} \\
&\quad + \frac{p\alpha_n (k_n \|x_n - x^*\|^p - \varphi_i(\|x_n - x^*\|))}{(1 + \|x_{n-1} - x^*\|)^p} \\
&\quad + p\alpha_n (6 + 4M)^{p-1} [d_n + (2\beta_n + \mu_n)(6 + 4M) \\
(2.9) \quad &\quad + \beta_n d_n + \mu_n M] + p\gamma_n M (6 + 4M)^{p-1}
\end{aligned}$$

for all $n \geq n_1, i = 1, 2, \dots, m$.

Since $1 - p\alpha_n k_n \rightarrow 1$ ($n \rightarrow \infty$), there exists $n_2 \geq n_1$ such that $0 < \frac{1}{2} < 1 - p\alpha_n k_n < 1$ for all $n \geq n_2$. It follows from (2.9) that

$$\begin{aligned}
\|x_n - x^*\|^p &\leq \frac{(1 - \alpha_n)^p \|x_{n-1} - x^*\|^p + p\alpha_n \xi_n - p\alpha_n \varphi_i(\|x_n - x^*\|)}{1 - p\alpha_n k_n} \\
(2.10) \quad &+ \frac{(p\alpha_n A_n + p\gamma_n M (6 + 4M)^{p-1}) (1 + \|x_{n-1} - x^*\|)^p}{1 - p\alpha_n k_n}
\end{aligned}$$

for all $n \geq n_2, i = 1, 2, \dots, m$,

where $A_n = (6 + 4M)^{p-1} [d_n + (2\beta_n + \mu_n)(6 + 4M) + \beta_n d_n + \mu_n M] \rightarrow$

$0(n \rightarrow \infty)$. Note that $(1 + \|x_{n-1} - x^*\|)^p \leq 2^{p-1}(1 + \|x_{n-1} - x^*\|^p)$,

$$\begin{aligned}(1 - \alpha_n)^p &= 1 - p\alpha_n + \frac{p(p-1)\alpha_n^2}{2!} - \frac{p(p-1)(p-2)\alpha_n^3}{3!} + \cdots + (-\alpha_n)^p \\ &= 1 - p\alpha_n + \alpha_n B_n,\end{aligned}$$

where

$$B_n = \frac{p(p-1)\alpha_n}{2!} - \frac{p(p-1)(p-2)\alpha_n^2}{3!} + \cdots + (-\alpha_n)^{p-1} \rightarrow 0(n \rightarrow \infty).$$

In virtue of (2.10), we conclude that

$$\begin{aligned}\|x_n - x^*\|^p &\leq \left[1 + \frac{p\alpha_n(k_n - 1) + \alpha_n B_n + p2^{p-1}\alpha_n A_n}{1 - p\alpha_n k_n} \right. \\ &\quad \left. + \frac{pM2^{p-1}(6 + 4M)^{p-1}\gamma_n}{1 - p\alpha_n k_n} \right] \|x_{n-1} - x^*\|^p \\ &\quad + \frac{p\alpha_n(\xi_n + 2^{p-1}A_n) + pM2^{p-1}(6 + 4M)^{p-1}\gamma_n}{1 - p\alpha_n k_n} \\ &\quad - \frac{p\alpha_n \varphi_i(\|x_n - x^*\|)}{1 - p\alpha_n k_n} \\ &\leq [1 + 2p\alpha_n(k_n - 1) + 2\alpha_n B_n + p2^p\alpha_n A_n \\ &\quad + pM2^p(6 + 4M)^{p-1}\gamma_n] \|x_{n-1} - x^*\|^p \\ &\quad + 2p\alpha_n(\xi_n + 2^{p-1}A_n) + pM2^p(6 + 4M)^{p-1}\gamma_n \\ &\quad - p\alpha_n \varphi_i(\|x_n - x^*\|)\end{aligned}\tag{2.11}$$

for all $n \geq n_2, i = 1, 2, \dots, m$.

Now we take a nonnegative integer $n_3 \geq n_2$ such that $x_{n_3} \neq x^*$ (if not, $x_n = x^*$ for all $n \geq n_2$, then $x_n \rightarrow x^*(n \rightarrow \infty)$, and so we have done). Since $k_n \rightarrow 1, \xi_n \rightarrow 0, B_n \rightarrow 0, A_n \rightarrow 0 (n \rightarrow \infty)$, $\sum_{n=0}^{\infty} \gamma_n < \infty$, there exists a positive integer $N > n_3$ such that, for all $n \geq N$, $(k_n - 1 + \frac{1}{p}B_n)(2G)^p + (\xi_n + 2^{p-1}A_n(1 + (2G)^p)) < \frac{\min_{1 \leq i \leq m} \{\varphi_i(G)\}}{4}$ and $\sum_{n=N}^{\infty} \gamma_n < \frac{G^p}{pM2^p(6+4M)^{p-1}(1+(2G)^p)}$, where $G = \max \{\|x_{n_3} - x^*\|, \|x_{n_3+1} - x^*\|, \dots, \|x_{N-1} - x^*\|, \|x_N - x^*\|\}$, and obviously $0 < G < \infty$.

Next we proceed by induction to show $\|x_{N+k} - x^*\| \leq 2G$ for all $k \geq 1$.

To see this consider two possible cases.

Case III: $\|x_{N+1} - x^*\| \leq G$.

In this case, $\|x_{N+1} - x^*\|^p \leq G^2 + pM2^p(6 + 4M)^{p-1}(1 + (2G)^p)\gamma_{N+1}$ and so we have the desired result.

Case IV: $\|x_{N+1} - x^*\| > G$.

In this case, $\varphi_i(\|x_{N+1} - x^*\|) > \varphi_i(G) \geq \min_{1 \leq i \leq m} \{\varphi_i(G)\} > 0$ since for each $i = 1, 2, \dots, m$, φ_i is a strictly increasing function. From (2.11), we also have

$$\begin{aligned}
 & \|x_{N+1} - x^*\|^2 \leq \|x_N - x^*\|^p \\
 & + [2p\alpha_{N+1}(k_{N+1} - 1) + 2\alpha_{N+1}B_{N+1} + pM2^p(6 + 4M)^{p-1}\gamma_{N+1} \\
 & + p2^p\alpha_{N+1}A_{N+1}](2G)^p + 2p\alpha_{N+1}(\xi_{N+1} + 2^{p-1}A_{N+1}) \\
 & + pM2^p(6 + 4M)^{p-1}\gamma_{N+1} - p\alpha_{N+1} \min_{1 \leq i \leq m} \{\varphi_i(G)\} \\
 & = \|x_N - x^*\|^p \\
 & - p\alpha_{N+1} \left[\min_{1 \leq i \leq m} \{\varphi_i(G)\} - 2(\xi_{N+1} + 2^{p-1}A_{N+1})(1 + (2G)^p) \right. \\
 & \left. - 2\left(k_{N+1} - 1 + \frac{1}{p}B_{N+1}\right)(2G)^p \right] + pM2^p(6 + 4M)^{p-1}(1 + (2G)^p)\gamma_{N+1} \\
 & \leq \|x_N - x^*\|^p - p\alpha_{N+1} \left(\min_{1 \leq i \leq m} \{\varphi_i(G)\} - \frac{\min_{1 \leq i \leq m} \{\varphi_i(G)\}}{4} \right) \\
 & + pM2^p(6 + 4M)^{p-1}(1 + (2G)^p)\gamma_{N+1} \\
 & \leq \|x_N - x^*\|^p + pM2^p(6 + 4M)(1 + (2G)^p)\gamma_{N+1} \\
 & \leq G^p + pM2^p(6 + 4M)^{p-1}(1 + (2G)^p)\gamma_{N+1}.
 \end{aligned}$$

By using induction, we get that

$$\begin{aligned}
 \|x_{N+k} - x^*\|^2 & \leq G^p + pM2^p(6 + 4M)^{p-1}(1 + (2G)^p) \sum_{i=N+1}^{N+k} \gamma_i \\
 & \leq G^p + G^p = 2G^p \leq (2G)^p
 \end{aligned}$$

for all $k \geq 1$.

This shows $\|x_n - x^*\| \leq 2G$ for all $n \geq N$. Therefore, it follows from (2.11) that

$$\|x_n - x^*\|^p \leq \|x_{n-1} - x^*\|^p + 2p\alpha_n \left[\left(k_n - 1 + \frac{B_n}{p} + 2^{p-1}A_n \right) (2G)^p \right]$$

$$\begin{aligned}
 & + \left[\xi_n + 2^{p-1}A_n \right] + pM2^p(6+4M)^{p-1}(1+(2G)^p)\gamma_n \\
 (2.12) \quad & - p\alpha_n\varphi_i(\|x_n - x^*\|)
 \end{aligned}$$

for all $n \geq N, i = 1, 2, \dots, m$.

Taking $\delta_n = p\alpha_n$, $c_n = pM2^p(6+4M)^{p-1}(1+(2G)^p)\gamma_n$, $a_n = \|x_n - x^*\|$ and $b_n = 2p\alpha_n \left[\left(k_n - 1 + \frac{B_n}{p} + 2^{p-1}A_n \right) (2G)^p + \xi_n + 2^{p-1}A_n \right]$ for all $n \geq N$. By (2.12) and Lemma 1.6 ensures that $\|x_n - x^*\| \rightarrow 0$ as $n \rightarrow \infty$, that is, $x_n \rightarrow x^*$ as $n \rightarrow \infty$. This completes the proof.

Remark 2.2. Theorem 2.1 improves and extends Theorem 1.2 (i.e., Theorem 2.1 of Yang [16]) in the following aspects:

(1) Extend asymptotically pseudocontractive mapping to asymptotically quasi-pseudocontractive type mappings.

(2) It abolishes the condition that $\sum_{n=0}^{\infty} \alpha_n(k_n - 1) < \infty$.

(3) The proof of sequence $\{x_n\}$ boundedness is entirely different from what it was before.

(4) Extend implicit iterative scheme (1.2) to Ishikawa type implicit iteration process (1.3).

(5) Condition

$$\limsup_{n \rightarrow \infty} \left\{ \langle T_i^n x_n - x^*, j(x_n - x^*) \rangle - k_n \|x_n - x^*\|^2 + \varphi(\|x_n - x^*\|) \right\} \leq 0$$

is replaced by the condition

$$\begin{aligned}
 \limsup_{n \rightarrow \infty} \left\{ \inf_{j_p(x_n - x^*) \in J_p(x_n - x^*)} \langle T_i^n x_n - x^*, j_p(x_n - x^*) \rangle \right. \\
 \left. - k_n \|x_n - x^*\|^p + \varphi_i(\|x_n - x^*\|) \right\} \leq 0.
 \end{aligned}$$

From Theorem 2.1, we obtain the following result immediately.

Theorem 2.3. Let E be a real normed linear space, K a nonempty bounded convex subset of E and $T_i : K \rightarrow K (i = 1, 2, \dots, m)$ a finite family of asymptotically nonexpansive mappings with $\{k_n^{(i)}\} \subset [1, \infty)$ such that $\lim_{n \rightarrow \infty} k_n = 1$, where $k_n = \max_{1 \leq i \leq m} \{k_n^{(i)}\}$. Assume that $F =$

$\bigcap_{i=1}^m F(T_i) \neq \emptyset$ denotes the set of common fixed points of $\{T_i\}_{i=1}^m$. Let $\{x_n\}$ be the sequence defined by (1.3). Suppose that $\{u_n\}$ and $\{v_n\}$ are bounded sequences in K and that $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$ and $\{\mu_n\}$ are sequences in $[0, 1]$ satisfying the following conditions:

- (i) $\sum_{n=0}^{\infty} \alpha_n = \infty$,
- (ii) $\alpha_n \rightarrow 0, \beta_n \rightarrow 0 (n \rightarrow \infty)$,
- (iii) $\sum_{n=0}^{\infty} \gamma_n < \infty, \mu_n \rightarrow 0 (n \rightarrow \infty)$.

Assume that there exist strict increasing functions $\varphi_i : [0, \infty) \rightarrow [0, \infty)$ with $\varphi_i(0) = 0$ such that

$$\limsup_{n \rightarrow \infty} \left\{ \inf_{j_p(x_n - x^*) \in J_p(x_n - x^*)} \langle T_i^n x_n - x^*, j_p(x_n - x^*) \rangle - k_n \|x_n - x^*\|^p + \varphi_i(\|x_n - x^*\|) \right\} \leq 0$$

for $x^* \in F$ and $i = 1, 2, \dots, m$. Then $\{x_n\}$ converges strongly to a common fixed point x^* of $\{T_i\}_{i=1}^m$.

Proof. Since T_i is an asymptotically nonexpansive mapping with $\{k_n\} \subset [1, \infty)$ such that $\lim_{n \rightarrow \infty} k_n = 1$, we have

$$\begin{aligned} \limsup_{n \rightarrow \infty} \{ \sup_{x, y \in K} (\|T_i^n x - T_i^n y\| - \|x - y\|) \} \\ \leq \limsup_{n \rightarrow \infty} [(k_n - 1) \text{diam}(K)] = 0, \end{aligned}$$

where $\text{diam}(K) = \sup_{x, y \in K} \|x - y\|$. This implies that every asymptotically nonexpansive mapping is asymptotically nonexpansive in the intermediate sense. Also since every asymptotically nonexpansive mapping is asymptotically pseudo-contractive mapping. The conclusion now follows easily from Theorem 2.1.

If $\gamma_n = \mu_n = 0 (\forall n \geq 1)$ in Theorem 2.1 and Theorem 2.3, then we have the following results.

Theorem 2.4. Let E be a real normed linear space, K a nonempty convex subset of E and $T_i : K \rightarrow K (i = 1, 2, \dots, m)$ a finite family of asymptotically nonexpansive in the intermediate sense and asymptotically quasi-pseudocontractive type mappings with $\{k_n^{(i)}\} \subset [1, \infty)$ such that $\lim_{n \rightarrow \infty} k_n = 1$, where $k_n = \max_{1 \leq i \leq m} \{k_n^{(i)}\}$. Assume that $F =$

$\bigcap_{i=1}^m F(T_i) \neq \emptyset$ denotes the set of common fixed points of $\{T_i\}_{i=1}^m$. Let $\{x_n\}$ be the sequence defined by

$$\begin{cases} x_n = (1 - \alpha_n)x_{n-1} + \alpha_n T_{i(n)}^n y_n \\ y_n = (1 - \beta)x_n + \beta_n T_{i(n)}^n x_n, \quad (n \geq 0). \end{cases}$$

Suppose that $\{\alpha_n\}, \{\beta_n\}$ are sequences in $[0, 1]$ satisfying the following conditions:

(i) $\sum_{n=0}^{\infty} \alpha_n = \infty$,

(ii) $\alpha_n \rightarrow 0, \beta_n \rightarrow 0 (n \rightarrow \infty)$.

Assume that there exist strict increasing functions $\varphi_i : [0, \infty) \rightarrow [0, \infty)$ with $\varphi_i(0) = 0$ such that

$$\limsup_{n \rightarrow \infty} \left\{ \inf_{j_p(x_n - x^*) \in J_p(x_n - x^*)} \langle T_i^n x_n - x^*, j_p(x_n - x^*) \rangle - k_n \|x_n - x^*\|^p + \varphi_i(\|x_n - x^*\|) \right\} \leq 0$$

for $x^* \in F$ and $i = 1, 2, \dots, m$. Then $\{x_n\}$ converges strongly to a common fixed point x^* of $\{T_i\}_{i=1}^m$.

Theorem 2.5. Let E be a real normed linear space, K a nonempty bounded convex subset of E and $T_i : K \rightarrow K (i = 1, 2, \dots, m)$ a finite family of asymptotically nonexpansive mappings with $\{k_n^{(i)}\} \subset [1, \infty)$ such that $\lim_{n \rightarrow \infty} k_n = 1$, where $k_n = \max_{1 \leq i \leq m} \{k_n^{(i)}\}$. Assume that $F = \bigcap_{i=1}^m F(T_i) \neq \emptyset$ denotes the set of common fixed points of $\{T_i\}_{i=1}^m$. Let $\{x_n\}$ be the sequence defined by

$$\begin{cases} x_n = (1 - \alpha_n)x_{n-1} + \alpha_n T_{i(n)}^n y_n \\ y_n = (1 - \beta)x_n + \beta_n T_{i(n)}^n x_n, \quad (n \geq 0). \end{cases}$$

Suppose that $\{\alpha_n\}, \{\beta_n\}$ are sequences in $[0, 1]$ satisfying the following conditions:

(i) $\sum_{n=0}^{\infty} \alpha_n = \infty$,

(ii) $\alpha_n \rightarrow 0, \beta_n \rightarrow 0 (n \rightarrow \infty)$.

Assume that there exist strict increasing functions $\varphi_i : [0, \infty) \rightarrow [0, \infty)$ with $\varphi_i(0) = 0$ such that

$$\limsup_{n \rightarrow \infty} \left\{ \inf_{j_p(x_n - x^*) \in J_p(x_n - x^*)} \langle T_i^n x_n - x^*, j_p(x_n - x^*) \rangle - k_n \|x_n - x^*\|^p + \varphi_i(\|x_n - x^*\|) \right\} \leq 0$$

for $x^* \in F$ and $i = 1, 2, \dots, m$. Then $\{x_n\}$ converges strongly to a common fixed point x^* of $\{T_i\}_{i=1}^m$.

REFERENCES

- [1] E. Aspiund, Positivity of duality mappings, *Bull. Amer. Math. Soc.* **73** (1967) 200-203.
- [2] R. E. Bruck, T. Kuczumow and S. Reich, Convergence of iterates of asymptotically nonexpansive mappings in Banach spaces with the uniform Opial property, *Colloq. Math.* **65** (1993), no. 2, 169-179.
- [3] S. S. Chang, K. K. Tanc, H. W. J. Lee and C. K. Chan, On the convergence of implicit iteration process with errors for a finite family of asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* **313** (2006), no. 1, 273-283.
- [4] C. E. Chidume, H. Zegeye, Strong convergence theorems for common fixed points of uniformly L -Lipschitzian pseudocontractive semi-groups, *Appl. Anal.* **86** (2007), no. 3, 353-366.
- [5] K. Goebel, W. A. Kirk, A fixed point theorem for asymptotically nonexpansive mappings, *Proc. Amer. Math. Soc.* **35** (1972) 171-174.
- [6] F. Gu, The new composite implicit iterative process with errors for common fixed points of a finite family of strictly pseudocontractive mappings, *J. Math. Anal. Appl.* **329** (2007), no. 2, 766-776.
- [7] S. H. Khan and W. Takahashi, Approximating common fixed points of two asymptotically nonexpansive mappings, *Sci. Math. Jpn.* **53** (2001), no. 1, 143-148.
- [8] E. U. Ofoedu, Strong convergence theorem for uniformly L -Lipschitzian asymptotically pseudocontractive mapping in a real Banach space, *J. Math. Anal. Appl.* **321** (2006), no. 2, 722-728.
- [9] M. O. Osilike, Implicit iteration process for common fixed points of a finite family of strictly pseudocontractive maps, *J. Math. Anal. Appl.* **294** (2004), no. 1, 73-81.
- [10] S. Plubtieng, K. Ungchittrakool and R. Wangkeeree, Implicit iterations of two finite families for nonexpansive mappings in Banach spaces, *Numer. Funct. Anal. Optim.* **28** (2007), no. 5-6, 737-749.
- [11] B. E. Rhoades, A comparison of various definition of contractive mappings, *Trans. Amer. Math. Soc.* **226** (1977) 257-290.
- [12] J. Schu, Iterative construction of fixed point of asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* **158** (1991), no. 2, 407-413.
- [13] Z. H. Sun, Strong convergence of an implicit iteration process for a finite family of asymptotically quasi-nonexpansive mappings, *J. Math. Anal. Appl.* **286** (2003), no. 1, 351-358.
- [14] H. K. Xu and R. G. Ori, An implicit iteration process for nonexpansive mappings, *Numer. Funct. Anal. Optim.* **22** (2001), no. 5, 767-773.
- [15] L. P. Yang and G. Hu, Convergence of implicit iteration process with random errors, *Acta Math. Sinica (Chin. Ser.)* **51** (2008), no. 1, 11-22.

- [16] L. P. Yang, Convergence theorems of an implicit iteration process for asymptotically pseudo-contractive mappings, *Bull. Iranian Math. Soc.* **38** (2012), no. 3, 699–713.
- [17] S. Y. Zhang, Modified generalized Ishikawa iterative approximations of fixed point for asymptotically quasi pseudocontractive type mappings in normed linear spaces, *Acta Math. Appl. Sin. (Chinese Series)* **34** (2011), no. 5, 886–894.
- [18] S. Y. Zhang, Iterative solution of strongly accretive operator equation in normed linear spaces, *Acta Anal. Funct. Appl.* **12** (2010), no. 12, 75–78.
- [19] H. Y. Zhou, S. S. Chang, Convergence of implicit iterative process for a finite family of asymptotically nonexpansive mappings in Banach spaces, *Numer. Funct. Anal. Optim.* **23** (2002), no. 7-8, 911–921.
- [20] H. Y. Zhou, Y. J. Cho, Ishikawa and Mann iterative processes with errors for nonlinear Φ -strongly quasi-accretive mappings in normed linear spaces, *J. Korean Math. Soc.* **36** (1999), no. 6, 1061–1073

(Shuyi Zhang) DEPARTMENT OF MATHEMATICS, BOHAI UNIVERSITY, JINZHOU
121013 CHINA

E-mail address: jzzhangshuyi@126.com

Archive of SID