

The Effect of Nutrition and Supplements on Ocular Health

HR Jahadi Hosseini*¹, M Mosallaei¹, Z Asadi Kalameh²

¹Department of Ophthalmology, ²Medical Student, Shiraz University of Medical Sciences, Shiraz, Iran

Abstract

Nutrition is a subject of interest in many fields of medicine. So ophthalmologists have also attempted to find possible ways to preserve vision through diet and supplements. Ocular disorders such as cataracts, age-related macular degeneration and glaucoma are the leading causes of visual impairment and blindness in the world, so most of the studies have focused on these major disorders and nutritions containing antioxidant such as vitamin C and E. Zexanthin/luteins and omega 3 have been the main substances studied in this relation. Although benefits of the regimens with high amounts of antioxidants were observed in reducing progression of cataract, age-related macular degeneration and so on, as many of these studies have been observational, the cause and effect relationship cannot be definitely concluded and multiple cohort prospective studies will be desired to evaluate the exact role of nutrition. Somehow, a healthy diet which means the diet which increases our health can be achieved in regimens with low saturated fatty acids and rich in fresh fruits, vegetables and fish. On the whole, even though they may not affect disease progression, they are generally good for overall health.

Keywords: Nutrition; Supplement; Ocular health

Introduction

Ocular disorders such as cataracts, age-related macular degeneration (ARMD) and glaucoma are the leading causes of visual blindness in the world. Currently, cataract extraction is the most common surgical procedure performed. It has been estimated that if the progression of cataracts could be delayed by at least 10 years, the number of cataract surgeries would be approximately reduced by 45 percent per year.¹ On the other hand, the severity and irreversibility of ARMD and glaucoma have caused interest in finding a modality to prevent or retard cataract formation. In this respect, nutrition modification is one promising subject which seems to be effective to fulfill our desire to some extent; however, the evaluation of the exact effects of nutrition on such long lasting ocular disorders, which are manifested as people age, is not practical anyway.² As a result, multiple observational cohort and misanalysis studies were performed to investigate any effect of nutrition on modula-

tion of those ocular disorders. Considering the eye as a certain area of the body with heavy exposure to light, the oxidative damage mediated by light exposure remains as the main cause of ocular problems.³⁻⁵ Oxidation is a process that allows free radicals to alter the genetic structure of a cell leading to disastrous chronic diseases in humans. However, any substance consumption which neutralizes this process of oxidation will protect human cells from damage. The question remaining to be answered is what substances and with what amount to be included in daily nutrition to gain optimal effect. Other issues concern the cost effectiveness of such regimens.⁶

As studies on this issue have come up with conflicting results, we attempted to have an overview of the nutritional substances pertaining to ocular health, the main sources of obtaining them, and the possible effect of these regimens in each ocular disorder.

Materials and Methods

The Pubmed databases were searched from 1993 to 2008, using the search term (nutrition), (nutrition and ocular health), (nutrition and eye), (nutrition and eye disease/ ocular disorder/cataract/age related macular

*Correspondence: Hamid Reza Jahadi Hosseini, MD, Department of Ophthalmology, Poostchi Eye Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. Tel: +98-711-2302830, Fax: +98-711-2302830, e-mail: jahadih@sums.ac.ir
Received: July 10, 2008 Accepted: November 2, 2008

disease or glaucoma) and (sources and caretenoid/zeaxanthin/lutein/vitamin C or vitamin E). In addition, English abstracts of articles which are non-English and of significant interest were included.

Nutritional substances and their main sources

Multiple kinds of carotenoids (plant pigments) give each fruit and vegetable its special colors. Such carotenoids as zeaxanthin and lutein, which give color from yellow to orange, provide a bright red hue to foods such as tomatoes, watermelon and pink grapefruit. Spinach, peas and many yellow/orange fruits and vegetables are rich in lutein. Corn, orange, peppers and certain leafy greens provide zeaxanthin. Lutein and zeaxanthin, however, can be found in many of the same foods and are often considered together as one value in the literature.

See the Tables below for food sources of lutein and zeaxanthin

FOOD SERVING SIZE LUTEIN (MG)	
Spinach, cooked 1 cup	13.3
Savoy cabbage, shredded 1 cup	10.2
Greens, cooked 1 cup	8.4
Broccoli, cooked 1 cup	3.4
Peas, cooked 1 cup	3.2
Spinach, raw 1 cup	1.8
Butter head lettuce 1 cup	0.9
Green pepper, sliced 1 cup	0.9

* Provided by American Optometric Association

Vitamin C

Vitamin C is an essential water-soluble vitamin considered as an important antioxidant which works to protect cells from damaging in the face of free radicals. Many fruits and vegetables are excellent sources of vitamin C.

See the Table below for a list of good food sources of vitamin C

FOOD SERVING SIZE VITAMIN C (MG)	
Papaya 1 medium	188
Orange juice 6 ounces	93
Strawberries 1 cup (8 whole)	84
Grapefruit juice 6 ounces	70
Cantaloupe, cubed 1 cup	68
Mango 1 medium	57
Green peppers, raw 1/2 cup	45
Broccoli, raw 1/2 cup	41
Cauliflower, raw 1/2 cup	23
Tomato 1 medium	23

*Provided by American Optometric Association

Vitamin E

Vitamin E is a fat-soluble vitamin. Alphatocopherol is the most active form of vitamin E that may protect the body against free radicals. Vegetable oils and nuts are also the good dietary sources of vitamin E.

See the Table below for a list of good food sources of vitamin E.

FOOD SERVING SIZE VITAMIN E MG/ (IU)	
Almonds 1 ounce	5.0 (7.5)
Safflower Oil 1 tablespoon	3.1 (4.7)
Corn Oil 1 tablespoon	1.9 (2.9)
Turnip Greens 1/2 cup	1.6 (2.4)
Mango 1 fruit	1.5 (2.3)
Peanuts 1 ounce	1.3 (2.1)
Broccoli 1/2 cup	1.0 (1.5)
Pistachio Nuts 1 ounce	0.8 (1.2)

*Provided by American Optometric Association

Omega-3 Poly Unsaturated Fatty Acid (PUFA)

The primary food source of omega-3 is sea food, especially cold water fish such as salmon, sardines and tuna.⁷⁻¹⁰

Age-related macular degeneration (ARMD) and nutrition

Age-related macular degeneration (ARMD) is a common cause of decreased vision and blindness in patients with the mean age over 40. The cause of ARMD remains unknown and the pathophysiologic mechanism of the damage of the retinal cells is beyond the scope of this text; however, numerous epidemiological studies have suggested a relationship between nutritional factors and ARMD.¹¹⁻¹⁶ As an instance, the relationship of lipid and ARMD is expected since certain kinds of lipids present in the retina have properties capable of reducing cell damage that may be encountered in patients with advanced ARMD.¹⁷ A daily diet high in n-3 polyunsaturated fat, especially from fish, may have protection against both forms of early and late ARMD. Also, people with the highest versus lowest quantities of 3-polyunsaturated fat as well as omega-3 fatty acids have a lower risk of developing neovascular ARMD and such substances may even hinder the progression of this event.^{18,19} The blue mountain Eye Study of more than 2000 Australian adults found that those whose diet was enriched in omega 3 fatty acid were less prone to developing both early and late ARMD.¹⁷ A recent case control study on 4000 individuals also

reported the benefits of regimens of omega -3 on abolishing the process of progression to neovascular type.²⁰ Moreover, as found in many studies, docosahexaenoic acid (an omega-3 polyunsaturated fatty acid) found more in the retina than pigmented epithelium, has especial functional properties in photoreceptors.^{16-18,21} The other compound is oleic acid or olive oil that was reported to be used as a suitable fat source to increase the absorption of dietary lutein in the management of age-related macular degeneration.²² An analysis by NHANES (National Health and Nutrition Examination Survey) showed the relationship between blood vitamin level and reduction in the incidence of early ARMD. The mechanism by which this finding was interpreted was related to the anti-inflammatory effect of vitamin D, leading to the establishment and reduction of the growth of new blood vessels.²³ On the other hand, the eye is particularly sensitive to oxidative stress because of direct exposure to light. Antioxidants, such as vitamin E, C or zinc, clearly have a protective and beneficial effect on ARMD.²¹ Individuals randomly assigned to receive antioxidants and zinc reduced their risk of aggravation of ARMD and loss of visual acuity.^{24,25} A previous study implied that the use of high doses of a combination of antioxidants (vitamin C, vitamin E, and beta carotene) and zinc reduced the risk of progression of advanced ARMD by about 25% in participants who had at least a moderate risk of developing ARMD.¹⁷ Also, levels of serum vitamin D were inversely associated with early ARMD but not advanced ARMD. This provides evidence that vitamin D may protect against ARMD.²⁶ Those with intermediate risk of age-related macular degeneration or advanced age-related macular degeneration in one eye are recommended to take the supplements. In Age-Related Eye Disease Study (AREDS), it has been proved to be promising in the prevention of advanced age-related macular degeneration by 25%. The formulation used by them was made of vitamins C, E, beta-carotene and zinc.²⁷ Subsequently a cocktail of antioxidants (supplements of vitamins C and E, beta carotene, and zinc) has been shown to reduce the risk of developing advanced (wet) ARMD by about 25%. In individuals with unilateral advanced ARMD, the combination of antioxidants studied in AREDS -- vitamins C and E, beta carotene, and zinc-- reduced the risk of developing advanced ARMD in the other eye.²⁸ The AREDS ARMD study also indicated that, compared with placebo, both the combination of antioxidants plus zinc and zinc alone considerably

reduced the risk of advanced ARMD in persons at risk of progression. They claimed that the combination of antioxidants and zinc had the most benefit, reducing the risk of development of moderate visual acuity loss (OR, 0.73; 99% CI, 0.54-0.99) and also advanced ARMD.^{29,30} However, the formulation of zinc and antioxidants is only recommended for persons at high risk, determined by a dilated eye examination by an ophthalmologist. This therapy is not a cure that resolves ARMD completely, but these are important in demonstrating a benefit for patients with early ARMD, before the development of advanced disease, as well as those who have already developed advanced disease in one eye.³¹ Development of ARMD appears to have several carbohydrate-related mechanisms as it was shown that a reduction in the dietary glycemic index (dGI), a modifiable risk factor, may provide a way in reducing the risk of ARMD.³²

In other words, higher dGI is associated with a greater risk of age-related macular degeneration (ARMD). Those at risk of ARMD progression, especially those at high risk of advanced ARMD may benefit from consuming a smaller amount of carbohydrates especially of the refined type.³³ It has been argued that the xanthophylls carotenoids lutein and zeaxanthin may protect the eyes against the age-related macular degeneration (ARMD). Macular pigment optical density (MPOD) is directly related to dietary consumption of lutein and zeaxanthin and also to their serum concentrations.³⁴⁻³⁷ Many studies suggest that people with ARMD have lower concentrations of lutein and zeaxanthin in the macula than those without ARMD.^{38,39} Suzan et al. (2006) studied the association between intermediate ARMD and lutein and zeaxanthin in the carotenoids in Age-Related Eye Disease Study (CAREDS). They selected women aged 50 to 79 years with intake of lutein plus zeaxanthin above the 78th (high) and below the 28th (low) percentiles, in whom the presence of AMD was determined by fundus. They concluded that diets rich in lutein plus zeaxanthin may protect against intermediate AMD in healthy women younger than 75 years.⁴⁰ In this regard, three mechanisms were interdicted by which lutein and zeaxanthin would protect against ARMD by absorbing the blue light, by quenching free radicals (reviewed by Landrum and Bone), or by increasing the membrane stability. Also, It has been hypothesized that lutein and zeaxanthin protect the macula against photo oxidative damage by their function as antioxidants and/or optical filters.^{40,41} Observational and clinical trials support the safety of higher

intakes of the lutein and zeaxanthin and their roles in improving clinical features of ARMD in patients.⁴² Also, it was reported that dietary lutein/zeaxanthin intake was inversely associated with neovascular ARMD.¹⁹ Moreover, the increase in the serum levels of lutein/zeaxanthin correlates with increases in the serum levels of their metabolites that have previously been detected in the ocular tissues. Elderly human subjects with and without ARMD can safely take supplements of lutein up to 10 mg/d for 6 months with no apparent toxicity or side effects.⁴³ However, in cigarette smokers, a high rate of lung cancer was documented in those receiving beta carotene supplements.^{44,45} So, except in cigarette smokers, carotenoids have been advocated as potential therapeutic agents in treating age-related macular degeneration (ARMD).⁴⁴⁻⁴⁷

Cataract and nutrition

Cataracts are opacities of the lens in the eye that result in decreased visual acuity in the elderly. According to literature regarding many experimental and observational researches, antioxidants, such as vitamin E, C or zinc, probably have a protective effect on cataract by their antioxidant properties.^{21,30} Moreover, the lens contains carotenoids, lutein and zeaxanthin, which have been shown to have antioxidant properties. It has been hypothesized that increasing the intake of lutein and zeaxanthin may prolong the onset of age-related cataracts.⁴⁸ In an experimental study on animals with sodium selenite induced cataract, L-cysteine and vitamin C were highly effective in preventing and retarding the process of cataractogenicity of this substance by maintaining the soluble protein concentrations of the lens.⁴⁹ In previous studies, it was reported that women in the group with high dietary levels of lutein and zeaxanthin had a 23% lower prevalence of nuclear cataract compared with those with low levels but the final results were in favor of only moderate association of those regimens and prevalence of nuclear cataract in older patients.⁵⁰⁻⁵³ Consequently, it was said that supplementation with vitamin C, lutein, zeaxanthin, or a multivitamin may help only certain populations in prevention of cataract, but it will probably affect the progression of cataracts in most patients.⁵⁴ Christen et al. (2008) in a study on women reported that higher dietary intakes of vitamin E from food and supplements were associated with significantly decreased risk of cataract.⁵³ Although in observational studies high levels of antioxidants especially vitamin E was found to be associated

with lower rates of nuclear cataract, in other prospective randomized placebo controlled clinical trials, the Age Related Eye Disease Study (AREDS) and recent vitamin E, cataract and Age Related Maculopathy Trial, it was found that neither the combination of vitamin E, C and B carotene nor vitamin E alone was effective in this relation.⁵⁵ Furthermore, osmotic stress caused by accumulation of polyols within the lens has been shown to be associated with glucose-induced cataractogenesis. Taurine has an antioxidant capacity and its level in diabetic cataractous lens is markedly decreased. Higgines et al. (2007) claimed that pretreatment of the lense with 30 miliMole taurine significantly reversed the level of protein carbonylation and reduced glutathione (GSH) compared to the control group. Therefore, taurine might spare GSH and protect ional effect on the lens in the face of oxidative stress induced by a high concentration of glucose.⁵⁷

According to the nutritional lipid profile, higher dietary intake of omega-3 polyunsaturated fatty acids (n-3 PUFA) was associated with a reduced incidence of nuclear cataract as dietary fat may affect the lens cell membrane composition and function, and these are related to age-related cataract.⁵⁸ In addition, higher alpha-linolenic acid (ALA) intake was associated with a greater age-related change in the lens nuclear density.⁵⁹ Suzan et al. studied the association between age-related nuclear cataract and lutein and zeaxanthin in the diet and serum in the carotenoids in the Age-Related Eye Disease Study (CAREDS). They reported that diets rich in lutein and zeaxanthin are moderately associated with decreased prevalence of nuclear cataract in older patients as patients in the group with high dietary levels of lutein and zeaxanthin had a 23% lower prevalence of nuclear cataract compared with those with low levels.⁶⁰ Conversely, an Italian trial which randomized more than 1000 older adults to a daily multivitamin supplement or placebo reported that while supplements reduced the risk of nuclear cataract by 34 percent, it increased the risk of posterior subcapsular cataract.⁶¹

Glaucoma and Nutrition

Glaucoma is one of the common causes of blindness in the elderly population. Increasing dietary omega-3 reduces IOP with age because of increased outflow facility, probably as the result of an increase in docosanoids. So, dietary formulations may provide a modifiable factor for IOP regulation. This can reduce the risk for glaucoma and lead to a role in treatment of the disease.³⁵ A higher intake of certain fruits

and vegetables enriched in vitamins may result in decreased risk of glaucoma.³⁶

Dry Eye and Nutrition

The women health study (WHS) found that women with a higher dietary intake of omega-3 fatty acids were less prone to dry eye.⁶² The mechanism by which this result could be interpreted may be related to the anti-inflammatory effect of Omega-3 found in meibomian gland. The same result was observed in patients with sjögrens by abolishing lacrimal gland inflammation. So, it was recommended to use fish oil as a main source of this substance especially in regions with low intake of fish. However, caution is recommended as these supplements may not be useful for health by increasing bleeding time and interfering with COX-2 inhibitors.^{63,64}

Summary

Nutrition is a subject of interest in all fields of medicine. Accordingly, in the field of ophthalmology attempts have been made to find possible ways to preserve vision through diet and supplements. In spite of several studies in this field,⁶⁵⁻⁷⁰ the most famous was the Age Related Eye Disease Study (AREDS), a randomized trial that reported the benefit of supplementation with antioxidants and zinc in patients with intermediate ARMD or loss of vision in one eye as a result of advanced disease. Age Related Eye Disease Study (AREDS) is important since it revealed a certain effect of supplementation on abolishing specific disease progression in specific groups of patients. The effect of nutrition on ocular health has not been neglected any way though most of the studies were observational and this made it difficult to make a definite conclusion about what to eat or what supplements to use as pills. Moreover, the results of some of the recent prospective cohort studies, while not confirming the previous results, may conversely give different conclusions. In other words, as stressed by Dr Chew, "most of those studies tell us that there may be something there, but still need to do well-conducted randomized controlled trials."^{65,71,74}

As previously mentioned, since the eyes are more vulnerable to light and oxidative damage by light,

trends are towards nutrition with high antioxidative properties such as vegetables containing beta carotene zexanthine /lutein, vitamin C, E and also those with high amounts of unsaturated fatty acid and omega 3 such as fish oil. Researchers hope that antioxidants might at least slow disease progression if not effective on prevention. Although the results of the first AREDS are currently used as a guideline in treatment of patients with ARMD, it is impossible to know if the relation of diet and disease progression is a real cause and effect one until cohort studies confirm it.^{72,73} "Fortunately a randomized trial called ARDES-2 is investigating the effect of these nutrients on the development of ARMD. Enrolment has begun and the results will be available in about five years. Therefore, "The results from ARDES-2 which is looking for cataract will be available in 2012. They are going to study the effect of lutein/Zexanthin and omega 3 fatty acid on cataract following 4000 participants in five years."⁷⁴ Somehow, a health diet which means the one that increases the life health can be achieved in regimens with low saturated fatty acids and enriched in fresh fruits, vegetables and fish. We can state briefly that even though they may not affect disease progression, they are generally good for overall health. It would be better to obtain these substances through natural foods and include them in the daily regimens rather than through supplements, since many fat soluble supplements such as vitamin E are risky for health. In addition, other supplements containing Beta carotene increase the risk of lung cancer in asbestoses workers and cigarette smokers. Also, combination of selenium, Beta carotene and vitamin E has been found to increase the risk of esophageal cancer.^{75,76} Despite these arguments, researchers still work on making pills proper for ocular health.

Acknowledgement

The authors would like to thank Dr. Nasrin Shokrpour and Shiva Mostafavi for editing the manuscript.

Conflict of interest: None declared.

References

- 1 Herm R.J. Age-related macular degeneration. *N Engl J Med* 2008; **359**:1735-6. [18927955] [doi:10.1056/NEJM081470]
- 2 Jacques PF. The potential preventive effects of vitamins for cataract and age-related macular degeneration. *Int J Vitam Nutr Res* 1999; **69**:198-205. [10389028] [doi:10.1024/0300-9831.69.3.198]
- 3 Seddon JM, Ajani UA, Sperduto RD, Hiller R, Blair N, Burton TC, Farber

- MD, Gragoudas ES, Haller J, Miller DT, et al. Dietary carotenoids, vitamins A, C, and E, and advanced age-related macular degeneration. Eye Disease Case-Control Study Group. *JAMA* 1994;**272**:1455-6. [7933422] [doi:10.1001/jama.272.18.1413]
- 4 Teichmann KD. Treatment of macular degeneration, according to Bangerter. *Eur J Med Res* 1997;**2**: 445-54. [9348273]
 - 5 Ciulla TA, Danis RP, Harris A. Age-related macular degeneration: a review of experimental treatments. *Surv Ophthalmol* 1998;**43**:134-46. [9763138] [doi:10.1016/S0039-6257(98)00014-9]
 - 6 Delcourt C, Cristol JP, Tessier F, Léger CL, Descamps B, Papoz L. Age-related macular degeneration and antioxidant status in the POLA study. POLA Study Group. *Pathologies Oculaires Liées à l'Age. Arch Ophthalmol* 1999;**117**:1384-90. [10532448]
 - 7 Talegawkar SA, Johnson EJ, Carithers TC, Taylor HA, Bogle ML, Tucker KL. Carotenoid intakes, assessed by food-frequency questionnaires (FFQs), are associated with serum carotenoid concentrations in the Jackson Heart Study: validation of the Jackson Heart Study Delta NIRI Adult FFQs. *Public Health Nutr* 2008;**11**:989-97. [18053294] [doi:10.1017/S1368980007001310]
 - 8 Strobel M, Tinz J, Biesalski HK. The importance of beta-carotene as a source of vitamin A with special regard to pregnant and breastfeeding women. *Eur J Nutr* 2007;**46**:11-20. [17665093] [doi:10.1007/s00394-007-1001-z]
 - 9 Singhal S, Gupta R, Goyle A. Comparison of antioxidant efficacy of vitamin E, vitamin C, vitamin A and fruits in coronary heart disease: a controlled trial. *J Assoc Physicians India* 2001;**49**:327-31. [11291971]
 - 10 Nykamp D, Kavanaugh ED, Wenker AP. Vitamins: the wise choice for women with cardiovascular disease. *Consult Pharm* 2007;**22**:490-502. [17713997]
 - 11 Biousse V, Bousser MG, Gaudric A. Age-related macular degeneration and risk of stroke. *J Fr Ophthalmol* 2008;**31**:111-25. [18401309] [doi:10.1016/S0181-5512(08)70343-7]
 - 12 Lornejad-Schäfer MR, Lambert C, Breithaupt DE, Biesalski HK, Frank J. Solubility, uptake and biocompatibility of lutein and zeaxanthin delivered to cultured human retinal pigment epithelial cells in tween40 micelles. *Eur J Nutr* 2007;**46**:79-86. [17225922] [doi:10.1007/s00394-006-0635-6]
 - 13 Mares JA. Potential value of antioxidant-rich foods in slowing age-related macular degeneration. *Arch Ophthalmol* 2006;**124**:1339-40. [16966632] [doi:10.1001/archophth.124.9.1339]
 - 14 McFadden SA, Howlett MH, Mertz JR, Wallman J. Acute effects of dietary retinoic acid on ocular components in the growing chick. *Exp Eye Res* 2006;**83**:949-61. [16797531] [doi:10.1016/j.exer.2006.05.002]
 - 15 Porter L, Reynolds N, Ellis JD. Total parenteral nutrition, vitamin E, and reversible macular dysfunction morphologically mimicking age related macular degeneration. *Br J Ophthalmol* 2005;**89**:1531-2. [16234469] [doi:10.1136/bjo.2005.074195]
 - 16 Koto T, Nagai N, Mochimaru H, Kurihara T, Izumi-Nagai K, Satofuka S, Shinoda H, Noda K, Ozawa Y, Inoue M, Tsubota K, Oike Y, Ishida S. Eicosapentaenoic acid is anti-inflammatory in preventing choroidal neovascularization in mice. *Invest Ophthalmol Vis Sci* 2007;**48**:4328-34. [17724224] [doi:10.1167/iov.06-1148]
 - 17 Weber U, Michaelis L, Wirth KE. Serum lipids and vitamins in senile macular degeneration. *Ophthalmologie* 1993;**90**:486-9. [8219637]
 - 18 Chua B, Flood V, Rochtchina E, Wang JJ, Smith W, Mitchell P. Dietary fatty acids and the 5-year incidence of age-related maculopathy. *Arch Ophthalmol* 2006;**124**:981-986. [16832021] [doi:10.1001/archophth.124.7.981]
 - 19 Robman L, Vu H, Hodge A, Tikellis G, Dimitrov P, McCarty C, Guymer R. Dietary lutein, zeaxanthin, and fats and the progression of age-related macular degeneration. *Can J Ophthalmol* 2007;**42**:720-6. [17724493] [doi:10.3129/107-116]
 - 20 SanGiovanni JP, Chew EY, Clemons TE, Davis MD, Ferris FL 3rd, Gensler GR, Kurinij N, Lindblad AS, Milton RC, Seddon JM, Sperduto RD. Age-Related Eye Disease Study Research Group. The relationship of dietary lipid intake and age-related macular degeneration in a case-control study: AREDS Report No. 20. *Arch Ophthalmol* 2007;**125**:671-9. [17502507] [doi:10.1001/archophth.125.5.671]
 - 21 Delcourt C. Application of nutrigenomics in eye health. *Forum Nutr* 2007;**60**:168-75. [17684413] [doi:10.1159/000107176]
 - 22 Lakshminarayana R, Raju M, Krishnanantha TP, Baskaran V. Lutein and zeaxanthin in leafy greens and their bioavailability: olive oil influences the absorption of dietary lutein and its accumulation in adult rats. *J Agric Food Chem* 2007;**55**:6395-400. [17602649]
 - 23 Fletcher AE, Bentham GC, Agnew M, Young IS, Augood C, Chakravarthy U, de Jong PT, Rahu M, Seland J, Soubrane G, Tomazzoli L, Topouzis F, Vingerling JR, Vioque J. Sunlight exposure, antioxidants, and age-related macular degeneration. *Arch Ophthalmol* 2008;**126**:1396-403. [18852418] [doi:10.1001/archophth.126.10.1396]
 - 24 Seddon JM, George S, Rosner B. Cigarette smoking, fish consumption, omega-3 fatty acid intake, and associations with age-related macular degeneration: the US Twin Study of Age-Related Macular Degeneration. *Arch Ophthalmol* 2006;**124**:995-1001. [16832023] [doi:10.1001/archophth.124.7.995]
 - 25 Clemons TE, Kurinij N, Sperduto RD. AREDS Research Group. Associations of mortality with ocular disorders and an intervention of high-dose antioxidants and zinc in the Age-Related Eye Disease Study: AREDS Report No. 13. *Arch Ophthalmol* 2004;**122**:716-26. [15136320] [doi:10.1001/archophth.122.5.716]
 - 26 Parekh N, Chappell RJ, Millen AE, Albert DM, Mares JA. Association between vitamin D and age-related macular degeneration in the Third National Health and Nutrition Examination Survey, 1988 through 1994. *Arch Ophthalmol* 2007;**125**:661-9. [17502506] [doi:10.1001/archophth.125.5.661]
 - 27 Coleman H, Chew E. Nutritional supplementation in age-related macular degeneration. *Curr Opin Ophthalmol* 2007;**18**:220-3. [17435429] [doi:10.1097/ICU.0b013e32814a586b]
 - 28 Kaushik S, Wang JJ, Flood V, Tan JS, Barclay AW, Wong TY, Brand-Miller J, Mitchell P. Dietary glycemic index and the risk of age-related macular degeneration. *Am J Clin Nutr* 2008;**88**:1104-10. [18842800]
 - 29 Jampol LM, Ferris FL 3rd. Antioxidants and zinc to prevent progression of age-related macular degeneration. *JAMA* 2001;**286**: 2466-8 [1759670] [doi:10.1001/jama.286.19.2466]
 - 30 Morris MS, Jacques PF, Chylack LT. Intake of zinc and antioxidant micronutrients and early age-related maculopathy lesions. *Ophthalmic Epidemiol* 2007;**14**:288-98. [17994438] [doi:10.1080/09286580601186759]
 - 31 Gottlieb JL. Age-Related Macular Degeneration. *JAMA* 2002;**288**: 2233-2236. [12425683] [doi:10.1001/jama.288.18.2233]
 - 32 Chiu CJ, Milton RC, Gensler G, Taylor A. Association between dietary glycemic index and age-related macular degeneration in nondiabetic participants in the Age-Related Eye Disease Study. *Am J Clin Nutr* 2007;**86**:180-8. [17616779]

- 33 Kowluru RA, Kanwar M, Chan PS, Zhang JP. Inhibition of retinopathy and retinal metabolic abnormalities in diabetic rats with AREDS-based micronutrients. *Arch Ophthalmol* 2008;**126**:1266-72. [18779489] [doi:10.1001/archophpt.126.9.1266]
- 34 Mares JA, LaRowe TL, Snodderly DM, Moeller SM, Gruber MJ, Klein ML, Wooten BR, Johnson EJ, Chappell RJ; CAREDS Macular Pigment Study Group and Investigators. Predictors of optical density of lutein and zeaxanthin in retinas of older women in the Carotenoids in Age-Related Eye Disease Study, an ancillary study of the Women's Health Initiative. *Am J Clin Nutr* 2006;**84**:1107-22. [17093164]
- 35 Nguyen CT, Bui BV, Sinclair AJ, Vingrys AJ. Dietary omega 3 fatty acids decrease intraocular pressure with age by increasing aqueous outflow. *Invest Ophthalmol Vis Sci* 2007;**48**:756-62. [17251475] [doi:10.1167/iovs.06-0585]
- 36 Coleman AL, Stone KL, Kodjebacheva G, Yu F, Pedula KL, Ensrud KE, Cauley JA, Hochberg MC, Topouzis F, Badala F, Mangione CM; Study of Osteoporotic Fractures Research Group. Glaucoma risk and the consumption of fruits and vegetables among older women in the study of osteoporotic fractures. *Am J Ophthalmol* 2008;**145**:1081-9. [18355790] [doi:10.1016/j.ajo.2008.01.022]
- 37 Krinsky NI, Landrum JT, Bone RA. Biologic mechanisms of the protective role of lutein and zeaxanthin in the eye. *Annu Rev Nutr* 2003;**23**:171-201. [12626691] [doi:10.1146/annurev.nutr.23.011702.073307]
- 38 Beatty S, Boulton M, Henson D, Koh HH, Murray IJ. Macular pigment and age-related macular degeneration. *Br J Ophthalmol* 1999;**83**:867-77. [10381676]
- 39 Bernstein PS, Zhao DY, Wintch SW, Ermakov IV, McClane RW, Gellermann W. Resonance Raman measurement of macular carotenoids in normal subjects and in age-related macular degeneration patients. *Ophthalmology* 2002;**109**:1780-7. [12359594] [doi:10.1016/S0161-6420(02)01173-9]
- 40 Moeller SM, Parekh N, Tinker L, Ritenbaugh C, Blodi B, Wallace RB, Mares JA; CAREDS Research Study Group. Associations between intermediate age-related macular degeneration and lutein and zeaxanthin in the Carotenoids in Age-Related Eye Disease Study (CAREDS): ancillary study of the Women's Health Initiative. *Arch Ophthalmol* 2006;**124**:1151-62. [16908818] [doi:10.1001/archophpt.124.8.1151]
- 41 Mares-Perlman JA, Fisher AI, Klein R, Palta M, Block G, Millen AE, Wright JD. Lutein and zeaxanthin in the diet and serum and their relation to age-related maculopathy in the third national health and nutrition examination survey. *Am J Epidemiol* 2001;**153**:424-32. [11226974] [doi:10.1093/aje/153.5.424]
- 42 Rhone M, Basu A. Phytochemicals and age-related eye diseases. *Nutr Rev* 2008;**66**:465-72. [18667008]
- 43 Age-Related Eye Disease Study Research Group, SanGiovanni JP, Chew EY, Clemons TE, Ferris FL 3rd, Gensler G, Lindblad AS, Milton RC, Seddon JM, Sperduto RD. The relationship of dietary carotenoid and vitamin A, E, and C intake with age-related macular degeneration in a case-control study: AREDS Report No. 22. *Arch Ophthalmol* 2007;**125**:1225-32. [17846363] [doi:10.1001/archophpt.125.9.1225]
- 44 Khachik F, de Moura FF, Chew EY, Douglass LW, Ferris FL 3rd, Kim J, Thompson DJ. The effect of lutein and zeaxanthin supplementation on metabolites of these carotenoids in the serum of persons aged 60 or older. *Invest Ophthalmol Vis Sci* 2006;**47**:5234-42. [17122108] [doi:10.1167/iovs.06-0504]
- 45 Obana A, Hiramitsu T, Gohto Y, Ohira A, Mizuno S, Hirano T, Bernstein PS, Fujii H, Iseki K, Tanito M, Hotta Y. Macular carotenoid levels of normal subjects and age-related maculopathy patients in a Japanese population. *Ophthalmology* 2008;**115**:147-57. [18166409] [doi:10.1016/j.ophtha.2007.02.028]
- 46 Wang W, Connor SL, Johnson EJ, Klein ML, Hughes S, Connor WE. Effect of dietary lutein and zeaxanthin on plasma carotenoids and their transport in lipoproteins in aged-related macular degeneration. *Am J Clin Nutr* 2007;**85**:762-9. [17344498]
- 47 Kalariya NM, Ramana KV, Srivastava SK, van Kuijk FJ. Carotenoid derived aldehydes-induced oxidative stress causes apoptotic cell death in human retinal pigment epithelial cells. *Exp Eye Res* 2008;**86**:70-80. [17977529] [doi:10.1016/j.exer.2007.09.010]
- 48 Trumbo PR, Ellwood KC. Lutein and zeaxanthin intakes and risk of age-related macular degeneration and cataracts: an evaluation using the Food and Drug Administration's evidence-based review system for health claims. *Am J Clin Nutr* 2006;**84**:971-4. [17093145]
- 49 HR Jahadi Hosseini, M Aminlari, MR Khalili. Prevention of Selenite-induced Cataract by L-Cysteine and Vitamin C in Rats. *Iranian Red Crescent Med J* 2008;**10**:281-287.
- 50 Berendschot TT, Broekmans WM, Klopping-Ketelaars IA, Kardinaal AF, Van Poppel G, Van Norren D. Lens aging in relation to nutritional determinants and possible risk factors for age-related cataract. *Arch Ophthalmol* 2002;**120**:1732-7. [12470150]
- 51 Hammond BR Jr. Possible role for dietary lutein and zeaxanthin in visual development. *Nutr Rev* 2008;**66**:695-702. [19019038] [doi:10.1111/j.1753-4887.2008.00121.x]
- 52 Harikumar KB, Nimita CV, Preethi KC, Kuttan R, Shankaranarayana ML, Deshpande J. Toxicity profile of lutein and lutein ester isolated from marigold flowers (*Tagetes erecta*). *Int J Toxicol* 2008;**27**:1-9. [18293208] [doi:10.1080/10915810701876265]
- 53 Christen WG, Liu S, Glynn RJ, Gaziano JM, Buring JE. Dietary carotenoids, vitamins C and E, and risk of cataract in women: a prospective study. *Arch Ophthalmol* 2008;**126**:102-9. [18195226] [doi:10.1001/archophpt.126.1.102]
- 54 Fernandez MM, Afshari NA. Nutrition and the prevention of cataracts. *Curr Opin Ophthalmol* 2008;**19**:66-70. [18090901] [doi:10.1097/ICU.0b013e3282f2d7b6]
- 55 Age Related Eye Disease Study Research Group. A randomized, placebo-controlled clinical trial of high-dose supplementation with vitamin C and E and beta carotene for age-related cataract and vision loss: AREDS report no.9. *ARCH Ophthalmol* 2001;**119**:1439-1452. [11594943]
- 56 Son HY, Kim H, H Kwon Y. Taurine prevents oxidative damage of high glucose-induced cataractogenesis in isolated rat lenses. *J Nutr Sci Vitaminol (Tokyo)* 2007;**53**:324-30. [17934237] [doi:10.3177/jnsv.53.324]
- 57 Higgins GT, Khan J, Pearce IA. Glycaemic control and control of risk factors in diabetes patients in an ophthalmology clinic: what lessons have we learned from the UKPDS and DCCT studies? *Acta Ophthalmol Scand* 2007;**85**:772-6. [17944626] [doi:10.1111/j.1600-0420.2007.00944.x]
- 58 Townend BS, Townend ME, Flood V, Burlutsky G, Rohtchina E, Wang JJ, Mitchell P. Dietary macronutrient intake and five-year incident cataract: the Blue Mountains eye study. *Am J Ophthalmol* 2007;**143**:932-939. [17459316] [doi:10.1016/j.ajo.2007.03.006]
- 59 Lu M, Taylor A, Chylack LT Jr, Rogers G, Hankinson SE, Willett WC, Jacques PF. Dietary linolenic acid intake is positively associated with five-year change in eye lens nuclear density. *J Am Coll Nutr*

- 2007;**26**:133-40. [17536124]
- 60 Moeller SM, Voland R, Tinker L, Blodi BA, Klein ML, Gehrs KM, Johnson EJ, Snodderly DM, Wallace RB, Chappell RJ, Parekh N, Ritenbaugh C, Mares JA; CAREDS Study Group; Women's Health Initiative. Associations between age-related nuclear cataract and lutein and zeaxanthin in the diet and serum in the Carotenoids in the Age-Related Eye Disease Study, an Ancillary Study of the Women's Health Initiative. *Arch Ophthalmol* 2008; **126**:354-64. [18332316] [doi:10.1001/archophth.126.3.354]
- 61 Clinical Trial of Nutritional Supplements and Age-Related Cataract Study Group, Maraini G, Sperduto RD, Ferris F, Clemons TE, Rosmini F, Ferrigno L. A randomized, double-masked, placebo-controlled clinical trial of multivitamin supplementation for age-related lens opacities. Clinical trial of nutritional supplements and age-related cataract report no. 3. *Ophthalmology* 2008; **115**:599-607. [18387406] [doi:10.1016/j.ophtha.2008.01.005]
- 62 Miljanović B, Trivedi KA, Dana MR, Gilbard JP, Buring JE, Schaumberg DA. Relation between dietary n-3 and n-6 fatty acids and clinically diagnosed dry eye syndrome in women. *Am J Clin Nutr* 2005; **82**:887-93. [16210721]
- 63 Tomlinson A, Pearce EI, Simmons PA, Blades K. Effect of oral contraceptives on tear physiology. *Ophthalmic Physiol Opt* 2001;**21**:9-16. [11220045]
- 64 Blades KJ, Patel S, Aidoo KE. Oral antioxidant therapy for marginal dry eye. *Eur J Clin Nutr* 2001;**55**:589-97. [11464232] [doi:10.1038/sj.ejcn.1601186]
- 65 Age-Related Eye Disease Study Research Group, SanGiovanni JP, Chew EY, Clemons TE, Ferris FL 3rd, Gensler G, Lindblad AS, Milton RC, Seddon JM, Sperduto RD. The relationship of dietary carotenoid and vitamin A, E, and C intake with age-related macular degeneration in a case-control study: AREDS Report No. 22. *Arch Ophthalmol* 2007;**125**:1225-32. [17846363] [doi:10.1001/archophth.125.9.1225]
- 66 SanGiovanni JP, Chew EY, Agrón E, Clemons TE, Ferris FL 3rd, Gensler G, Lindblad AS, Milton RC, Seddon JM, Klein R, Sperduto RD; Age-Related Eye Disease Study Research Group. The relationship of dietary omega-3 long-chain polyunsaturated fatty acid intake with incident age-related macular degeneration: AREDS report no. 23. *Arch Ophthalmol* 2008;**126**:1274-9. [18779490] [doi:10.1001/archophth.126.9.1274]
- 67 Kuehn BM. Studies probe diet's role in eye disease. *JAMA* 2005;**294**:32-3. [15998878] [doi:10.1001/jama.294.1.32]
- 68 Evans J. Antioxidant supplements to prevent or slow down the progression of AMD: a systematic review and meta-analysis. *Eye* 2008;**22**:751-60. [18425071] [doi:10.1038/eye.2008.100]
- 69 Morris MS, Jacques PF, Chylack LT, Hankinson SE, Willett WC, Hubbard LD, Taylor A. Intake of zinc and antioxidant micronutrients and early age-related maculopathy lesions. *Ophthalmic Epidemiol* 2007; **14**:288-98. [17994438] [doi:10.1080/09286580601186759]
- 70 Kowluru RA, Menon B, Gierhart DL. Beneficial effect of zeaxanthin on retinal metabolic abnormalities in diabetic rats. *Invest Ophthalmol Vis Sci* 2008;**49**:1645-51. [18385086] [doi:10.1167/iovs.07-0764]
- 71 Huang LL, Coleman HR, Kim J, de Monasterio F, Wong WT, Schleicher RL, Ferris FL 3rd, Chew EY. Oral supplementation of lutein/zeaxanthin and omega-3 long chain polyunsaturated fatty acids in persons aged 60 years or older, with or without AMD. *Invest Ophthalmol Vis Sci* 2008;**49**:3864-9. [18450596] [doi:10.1167/iovs.07-1420]
- 72 Age-Related Eye Disease Study Research Group. A Randomized, Placebo-Controlled, Clinical Trial of High-Dose Supplementation With Vitamins C and E, Beta Carotene, and Zinc for Age-Related Macular Degeneration and Vision Loss: AREDS Report No. 8. *Arch Ophthalmol* 2001;**119**: 1417-1436. [11594942]
- 73 Lee AY, Brantley MA Jr. CFH and LOC387715/ARMS2 genotypes and antioxidants and zinc therapy for age-related macular degeneration. *Pharmacogenomics* 2008;**9**:1547-50. [18855541] [doi:10.2217/14622416.9.10.1547]
- 74 Doven schuyler. Can diet and supplement preserve vision? *Eurotimes* 2008;**13**:10-11.
- 75 Chen X, Mikhail SS, Ding YW, Yang G, Bondoc F, Yang CS. Effects of vitamin E and selenium supplementation on esophageal adenocarcinogenesis in a surgical model with rats. *Carcinogenesis* 2000;**21**:1531-6. [10910955] [doi:10.1093/carcin/21.8.1531]
- 76 Wright ME, Virtamo J, Hartman AM, Pietinen P, Edwards BK, Taylor PR, Huttunen JK, Albanes D. Effects of alpha-tocopherol and beta-carotene supplementation on upper aerodigestive tract cancers in a large, randomized controlled trial. *Cancer* 2007;**109**:891-8. [17265529] [doi:10.1002/cncr.22482]