The Relationship between Geographic Tongue and Stress

H Ebrahimi¹, S Pourshahidi¹, A Andisheh Tadbir²*, S Bakhshi Shyan³

¹Department of Oral Medicine, ²Department of Oral Pathology, School of Dentistry, Shiraz University of Medical Science, Shiraz, Iran

Abstract

Background: Geographic tongue (GT) is a fairly common and usually asymptomatic disorder that is often detected on routine dental examinations. Characteristic lesions appear as multiple, well-demarcated zones of erythema, concentrated at the tip and lateral borders of tongue. Although the etiology of the condition is unknown, it may be related to stress. The purpose of this study was to evaluate the association between GT and stress.

Method: This study was performed on 60 patients with GT and 60 patients without GT. All the patients filled a questionnaire and a psychologist evaluted them.

Results: The total mean score of stress was 19.8 in GT group and 15.8 in the control group.

Conclusion: According to the results, there is an association between stress and GT. Decreasing stress in GT patients can lead to the healing of the lesion. Other factors that may be associated with GT should be studied in future.

Keywords: Geographic tongue; Stress; Perceived Stress Scale

Introduction

Geographic tongue or erythema migrans is a common benign condition that primarily affects the tongue. It is often detected on routine examination of the oral mucosa. The characteristic lesions of erythema migrans are seen on the anterior two thirds of the dorsal tongue mucosa. They appear as multiple, welldemarcated zones of erythema, concentrated at the tip and lateral borders of tongue. This erythema is due to atrophy of the filliform papilla, and these atrophic areas are typically surrounded at least partially by a slightly elevated, yellowish-white, serpentine or scalloped border. The lesions appear quickly in one area, healing within a few days or weeks, and then they develop in a very different area. Frequently, the lesion begins as a small white patch, which then develops a central erythematous atrophic zone and enlarges centrifugally. The lesions are usually asymptomatic, although a burning sensation or sensitivity to hot or spicy foods may be noted when the lesions are active.¹

Very infrequently, erythema migrans may occur on oral mocusal sites other than the tongue. In this instance, the condition is called ectopic geographic tongue.¹

Even though erythema migrans has been documented for many years, the etiopathogenesis is still unknown. Many risk factors have been proposed for GT including hormonal disturbances and oral contraceptive use,² psychological findings,³ diabetes mellitus,⁴ allergic conditions such as atopy,⁵ hay fever and rhinitis,⁶ dermatological diseases such as psoriasis,⁷ and Reiter's Syndrome.⁸ There is also a reported correlation with Down Syndrome⁹ and fissured tongue.¹⁰ A family history has also been reported to be associated with GT¹¹ which may be genetic and linked to major histocompatibility complex.

Psychosomatic factors appear to play a significant role in the etiology of geographic tongue.¹² It has

^{*}Correspondence: Azadeh Andisheh Tadbir, DMD, MSc, Assistant professor of Department of Oral Pathology, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran. Tel: +98-711-6263193-4, Fax: +98-711-6270325, e-mail: andisheh@sums.ac.ir, andisheh202003@yahoo.com Received: June 2, 2009 Accepted: October 6, 2009

Ebrahimi et al.

been reported that lesions arise in connection with pronounced emotional stress. Redman *et al.* found a higher prevalence of geographic tongue in mentally ill patients than in university students. They also noted that under emotional stress the student group with geographic tongue tended to have more severe lesions. These findings support the possible role of psychological factors in the etiology of geographic tongue.³ The purpose of this study was to evaluate the association between GT and stress.

Materials and Methods

Sixty patients (38 women and 22 men) with GT were selected for this study. They were referred to the Department of Oral Medicine in Tabriz University of Medical Sciences from May 2008 to July 2008. A specialist in the Oral Medicine Department made the diagnosis mainly on clinical features. The control group consisted of 60 healthy persons without GT among whom there were some patients who had referred to the department for dental screen and check up; we used simple random sampling. The patient and control groups filled the Perceived Stress Scale (PSS) questionnaire¹³ A psychologist evaluted the results. All the subjects were informed about the research characteristics and agreed to participate in the study by signing the free and informed consent form (ethical issue). We used Chi-square test and SPSS software (Version 15, Chicago, IL, USA) to analyze the data.

Results

The mean age of the GT group was 27.5 and that of the control group was 24.6 years. There were 77 women (64.2%) and 43 men (35.8%); 38 women and 22 men were in the GT group and 39 women and 21 men were in the control group. The total mean score of sterss in the GT group was 19.8 and that in the control group was 15.9.

The difference was statistically significant (p<0.001). There was neither a significant relationship between age and GT (p=0.217) nor between sex and the disorder (p=0.465) but stress had a significant relationship with GT (p<0.001). Adding one score to the final score of stress, risk of GT increased 1.16 times (OR=1.16)

Discussion

Geographic tongue is a common lesion that mostly appears on the dorsal and lateral bordre of the tongue.¹ No study could mention a specific ethiology for GT; yet there are several factors such as stress, allergy, and genetic and systemic diseases.¹ According to the results of our study, GT is associated with sterss; this is consistent with Redman's reports.³

Shulman and Carpenter¹⁴ studied the prevalence and risk factors associated with geographic tongue among US adults. Their findings showed no significant relationship between stress and GT. One explanation for this discrepancy may be due to differences in the recording of stress. There are no physiologic measures of stress such as serum cortisol levels, or questionairs directly addressing stress.

Jainkittivong and Laglais¹⁵ investigated the clinical characteristics and assessed other factors associated with geographic tongue in Thailand. They found no significant difference in the incidence of medical problems between the subjects and controls with geographic tongue. The incidence of stress-related condition in the subject group was not significantly higher than that in the control group. So, stress was not an etiologic factor for GT. This might, of course, be due to few samples in that study.

Zegareli¹⁶ studied 57 patients sufferring from burning mouth and found that psychiatric factors were the most common cause. There was no association between GT and psychologic disorders. This can be because of the use of a small number of samples and insuficient assessment of stress.

In Bánóczy *et al's*.¹⁷study GT had an association with emotional sterss. They studied GT's clinical and histopathological features. Women were more affected (65.7%) and the patients were mostly in their 5^{th} decade of life or older. GT in 20% of cases had co-incidence with other anomalies of the tongue and in other 20% it was associated with emotional stress.

In the present case control study, we evaluated the association of emotional stress with GT, using a standard questionnaire, but other etiologic factors such as strength and limitation were not evaluated. We recommend other studies to evaluate the role of other factors in the etiology of GT. According to the results, sterss has an association with GT and decreasing stress in GT patients can be helpful in healing the lesions.

Conflict of interest: None declared.

References

- Nevill BW, Damm DD, Allen CM, Bouguot JE. Oral and Maxillofacial Pathology, 3rd ed. W.B. Saunders Company. Philadelphia, 2009; pp.677-9.
- Waltimo J. Geographic tongue during a year of oral contraceptive cycles. Br Dent J 1991;171:94-6. [188 8590] [doi:10.1038/sj.bdj.4807618]
- 3 Redman RS, Vance FL, Gorlin RJ, Peagler FD, Meskin LH. Psychological component in the etiology of geographic tongue. J Dent Res 1966;45:1403-8. [4380992]
- 4 Wysocki GP, Daley TD. Benign migratory glossitis in patients with juvenile diabetes. Oral Surg Oral Med Oral Pathol 1987;63:68-70. [3468467] [doi:10.1016/0030-4220 (87)90342-2]
- 5 Marks R, Simons MJ. Geographic tongue--a manifestation of atopy. Br J Dermatol 1979;101:159-62. [486 322] [doi:10.1111/j.1365-2133.1979. tb05601.x]
- 6 Marks R, Czarny D. Geographic tongue: sensitivity to the environment. Oral Surg Oral Med Oral Pathol 1984;58:156-9. [6592509] [doi: 10.1016/0030-4220(84)90130-0]

- 7 Daneshpazhooh M, Moslehi H, Akhyani M, Etesami M. Tongue lesions in psoriasis: a controlled study. BMC Dermatol 2004;4:16. [15527508] [doi:10.1186/1471-5945-4-16]
- 8 Weathers DR, Baker G, Archard HO, Burkes EJ Jr. Psoriasiform lesions of the oral mucosa (with emphasis on "ectopic geographic tongue"). Oral Surg Oral Med Oral Pathol 1974;37:872-88. [4524888] [doi:10.1016/0030-4220(74)90440-X]
- 9 Ercis M, Balci S, Atakan N. Dermatological manifestations of 71 Down syndrome children admitted to a clinical genetics unit. *Clin Genet* 1996;**50**:317-20. [9007317]
- 10 Eidelman E, Chosack A, Cohen T. Scrotal tongue and geographic tongue: polygenic and associated traits. Oral Surg Oral Med Oral Pathol 1976;42:591-6. [1068416] [doi:10.1016/0030-4220(76)90210-3]
- 11 Fenerli A, Papanicolaou S, Papanicolaou M, Laskaris G. Histocompatibility antigens and geographic tongue. Oral Surg Oral Med Oral Pathol 1993;76:476-9. [8233428] [doi:10.1016/0030-4220(93)90015-V]

- 12 Saprio SM, Shklar G. Stomatitis areata migrans. Oral Surg Oral Med Oral Pathol 1973;36:28-33. [4514522] [doi:10.1016/0030-4220(73)90263-6]
- Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. J Health Soc Behav 1983; 24:385-96. [6668417] [doi:10.2307/ 2136404]
- 14 Shulman JD, Carpenter WM. Prevalence and risk factors associated with geographic tongue among US adults. *Oral Dis* 2006;12:381-6. [16792723] [doi:10.1111/j.1601-08 25.2005.01208.x]
- 15 Jainkittivong A, Langlais RP. Geographic tongue: clinical characteristics of 188 cases. *J Contemp Dent Pract* 2005;6:123-35. [15719084]
 16 Zegarelli DJ. Burning mouth: an
- 16 Zegarelli DJ. Burning mouth: an analysis of 57 patients. Oral Surg Oral Med Oral Pathol 1984;58:34-8. [6589575] [doi:10.1016/0030-4220 (84)90360-8]
- Bánóczy J, Szabó L, Csiba A. Migratory glossitis. A clinical-histologic review of seventy cases. Oral Surg Oral Med Oral Pathol 1975;39:113-21. [1053669] [doi:10.1016/0030-4220(75)90401-6]