

Exercise-Induced ST-Segment Elevation in Lead aVR as a Predictor of LCx Stenosis

Dear Editor,

Exercise-induced ST-segment depression, although the most common accepted criterion for detecting coronary artery disease (CAD) has limited sensitivity and is unable to localize ischemic territory.¹⁻⁴ Recently ST-segment elevation in lead aVR has been studied for the detection of CAD.¹⁻⁷ This ECG finding is usually associated with ST-segment depression in other leads, so the question is whether this ECG finding during exercise tolerance test (ETT) adds any further diagnostic information over the presence of ST-segment depression alone or not? Different studies have reported exercise-induced ST-segment elevation in lead aVR to be associated with proximal left anterior descending artery (LAD), left main coronary artery (LMCA), left circumflex artery (LCx) or even right coronary artery (RCA) stenosis in various clinical settings.¹⁻⁹ Considering these inconclusive data, we decided to perform a cross-sectional study on patients who had positive ETT associated with ST segment elevation in lead aVR.

A total of 560 patients with symptoms resembling angina were referred to ETT room of two outpatient clinics from January 2007 through March 2009. Fifty patients with positive ETT who had co-existing ST segment elevation in lead aVR constituted the first group of study. Demographic data as well as CAD risk factors including history of hypertension, diabetes, previous history of CAD, smoking, hyperlipidemia, BMI and family history of premature CAD were obtained. Another 50 patients with positive ETT but without ST segment elevation in lead aVR who were matched one by one with the first group of patients regarding demographic data and CAD risk factors, constituted the second group of study. The study was approved by Ethical Committee of Shiraz University

of Medical Sciences. *P* values of 0.05 or less were considered significant.

Regarding the number of vessels involved (i.e. single, two or three vessel disease), there was no statistical significance (Chi-Square test, *p* value = 0.740), but regarding the type of coronary vessel, it was shown that LCx was involved significantly higher in the first group than the second group (Chi-Square test, *p* value = 0.016, Table 1). This finding was in accordance with the finding of Gaitonde *et al.* that significant proximal LCx stenosis should cause ST-segment elevation in both lead aVR and V1, reflecting posterior wall ischemia. Using vectorcardiology, they proposed ECG changes in the posterolateral wall to be reflected in lead aVR.¹⁰ Our study provided further evidences in support of this hypothesis although not supported by other investigators.

This study was unique in the way that it was designed in the form of a cross-sectional study only focusing on ST-segment elevation in lead aVR. So, further studies are needed to confirm this finding. It was also demonstrated that exercise induced ST-segment elevation in lead aVR had a sensitivity of 64%, specificity of 50%, positive predictive value of 32% and negative predictive value of 21% in predicting LCx stenosis. These numbers may not be comparable with the results reported in other studies, but considering the fact that the stenosis of LCx was one of the main causes of false negative results in ETT, its importance may become more apparent.

Some studies have concentrated on specific groups of patients, either during or shortly after myocardial infarction while our selection of study population was performed on a large group of patients with a broad spectrum of demographic data and therefore more applicable to actual clinical setting. One of the limita-

Table 1: Angiographic findings of the patients

Vessel involvement	ST-segment elevation in lead aVR (Group 1) (%)	No ST-segment elevation in lead aVR (Group 2) (%)	<i>P</i> value
LMCA	-	2 (4.0)	0.495
LAD	24 (48)	22 (44)	0.841
LCx	30 (60)	17 (34)	0.016
RCA	21 (42)	20 (40)	0.673
Normal angiography	18 (36)	18 (36)	

tions of our study was the low number of patients. Another limitation is that the sensitivity, specificity, negative and positive predictive value provided were calculated among patients with positive ETT and so can not be extended to the whole population.

Keywords: ST-segment elevation; aVR lead; Exercise tolerance test; Coronary angiography

Conflict of interest: None declared.

MA Ostovan¹, AA Zolghadrasli^{2*}

¹Department of Cardiology, School of Medicine,
²Student Research Center, Shiraz University of
Medical Sciences, Shiraz, Iran

*Correspondence: Abdolali Zolghadrasli, MD, Student Research Center, Shiraz University of Medical Sciences, Shiraz, Po Box 71345-1975, Iran. Tel/Fax: +98-711-2122390, e-mail: zolghadr@sums.ac.ir
Received: May 20, 2011 Accepted: August 12, 2011

References

- 1 Baine KR, Kalia N, Carter D, Hrynychyn G, Kasza L, Lee TK, Wirzba B, Senaratne MP. Right precordial leads and lead aVR at exercise electrocardiography: does it change test results? *Ann Noninvasive Electrocardiol* 2006;**11**:247-52. [16846440] [<http://dx.doi.org/10.1111/j.1542-474X.2006.00111.x>]
- 2 Michaelides AP, Psomadaki ZD, Dilaveris PE, Richter DJ, Andrikopoulos GK, Aggeli KD, Stefanadis CI, Toutousaz PK. Improved detection of coronary artery disease by exercise electrocardiography with the use of right precordial leads. *N Engl J Med* 1999;**340**:340-5. [9929523] [<http://dx.doi.org/10.1056/NEJM199902043400502>]
- 3 Neill J, Shannon HJ, Morton A, Muir AR, Harbinson M, Adgey JA. ST segment elevation in lead aVR during exercise testing is associated with LAD stenosis. *Eur J Nucl Med Mol Imaging* 2007;**34**:338-45. [17019610] [<http://dx.doi.org/10.1007/s00259-006-0188-1>]
- 4 Michaelides AP, Psomadaki ZD, Richter DJ, Dilaveris PE, Andrikopoulos GK, Stefanadis C, Gialafos JE, Toutouzas PK. Significance of exercise-induced simultaneous ST-segment changes in lead aVR and V5. *Int J Cardiol* 1999;**71**:49-56. [10522564] [[http://dx.doi.org/10.1016/S0167-5273\(99\)00115-1](http://dx.doi.org/10.1016/S0167-5273(99)00115-1)]
- 5 Ozmen N, Yiginer O, Uz O, Kardesoglu E, Aparci M, Isilak Z, Cingozbay BY, Cebeci BS, Kocum HT. ST elevation in the lead aVR during exercise treadmill testing may indicate left main coronary artery disease. *Kardiol Pol* 2010;**68**:1107-11. [20967704]
- 6 Rostoff P, Wnuk M, Piwowska W. Clinical significance of exercise-induced ST-segment elevation in lead aVR and V1 in patients with chronic stable angina pectoris and strongly positive exercise test results. *Pol Arch Med Wewn* 2005;**114**:1180-9. [16789487]
- 7 Tuna Katircibaşı M, Tolga Koçum H, Tekin A, Erol T, Tekin G, Baltali M, Muderrisoglu H. Exercise-induced ST-segment elevation in leads aVR and V1 for the prediction of left main disease. *Int J Cardiol* 2008;**128**:240-3. [17658634] [<http://dx.doi.org/10.1016/j.ijcard.2007.05.022>]
- 8 Yamaji H, Iwasaki K, Kusachi S, Murakami T, Hirami R, Hamamoto H, Hina K, Kita T, Sakakibara N, Tsuji T. Prediction of acute left main coronary artery obstruction by 12-lead electrocardiography. ST segment elevation in lead aVR with less ST segment elevation in lead V(1). *J Am Coll Cardiol* 2001;**38**:1348-54. [11691506] [[http://dx.doi.org/10.1016/S0735-1097\(01\)01563-7](http://dx.doi.org/10.1016/S0735-1097(01)01563-7)]
- 9 Kosuge M, Kimura K, Ishikawa T, Ebina T, Hibi K, Toda N, Umemura S. ST-segment depression in lead aVR: a useful predictor of impaired myocardial reperfusion in patients with inferior acute myocardial infarction. *Chest* 2005;**128**:780-6. [16100167] [<http://dx.doi.org/10.1378/chest.128.2.780>]
- 10 Gaitonde RS, Sharma N, Ali-Hasan S, Miller JM, Jayachandran JV, Kalaria VG. Prediction of significant left main coronary artery stenosis by the 12-lead electrocardiogram in patients with rest angina pectoris and the withholding of clopidogrel therapy. *Am J Cardiol* 2003;**92**:846-8. [14516891] [[http://dx.doi.org/10.1016/S0002-9149\(03\)00898-1](http://dx.doi.org/10.1016/S0002-9149(03)00898-1)]