

# The Role of Vitamin C in Prevention of Preterm Premature Rupture of Membranes

Nayereh Ghomian<sup>1</sup>, Leili Hafizi<sup>1\*</sup>, Zahra Takhti<sup>1</sup>

<sup>1</sup> Department of Obstetrics and Gynecology, Imam Reza Hospital, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, IR Iran

\* Corresponding author: Leili Hafizi, Department of Obstetrics and Gynecology, Women's Health Research Center, Imam Reza Hospital, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, IR Iran. Tel: +98-5118022608, Fax: +98-5118525305, E-mail: hafizil@mums.ac.ir.

## ABSTRACT

**Background:** Preterm premature rupture of membranes (PPROM) is one of the most important complications of the pregnancy and cause perinatal morbidity and mortality. History of PPRM is a risk factor of recurrent PPRM. Vitamin C plays an important role in collagen metabolism and increases resistance maintenance of the chorioamniotic membranes.

**Objectives:** The aim of this study is to evaluate the role of vitamin C supplementation in prevention of PPRM in women with a positive history of PPRM.

**Patients and Methods:** This clinical trial study was performed on 170 pregnant women with the history of PPRM, with singleton pregnancy and gestational age 14 weeks in Imam-Reza Hospital, Mashhad University of Medical Sciences during 2008 to 2010. They were randomly divided into two groups. The case patients received 100 mg vitamin C daily from 14th weeks of gestation. PPRM occurrence was compared between two groups as an indicator of the protective effect of vitamin C supplements.

**Results:** PPRM occurred in 44.7% of controls and 31.8% of cases ( $P < 0.05$ ). PROM occurred in 34.1% of controls and 18.8% of cases ( $P < 0.05$ ). Pregnancy was terminated at term gestation in 21.2% of controls and 49.4% of cases ( $P < 0.05$ ). Rupture of membranes was significantly decreased in the case group.

**Conclusions:** Vitamin C supplementations after 14th weeks of gestation can prevent from PPRM in women with the history of PPRM.

**Keywords:** Ascorbic Acid; Pregnancy; Fetal Membranes, Premature Rupture; Prevention and Control

Copyright © 2013, Iranian Red Crescent Medical Journal; Published by Kowsar Corp.

## 1. Background

Premature rupture of membranes (1) defined as leakage of amniotic fluid through ruptured chorioamniotic membranes that occur before starting the labor pain at any gestational age. It is one of the most common problems in obstetrics and affects 10-20% of all pregnancies. (2,3) If PROM occurs before term pregnancy (37 gestational weeks), it is named as preterm premature rupture of

membranes (PPROM). Although PPRM involves 3% of pregnancies, it is the cause of one third of all preterm births with increased rates of neonatal and maternal morbidity and mortality (4). Its pathophysiologic mechanism is little known. It has been reported that premature ruptured membranes have less collagen content, which is necessary to provide mechanical strength to fetal mem-

► Article type: Research Article; Received: 11 Apr 2012, Revised: 12 Jul 2012, Accepted: 08 Jan 2013; DOI: 10.5812/ircmj.5138

► Implication for health policy/practice/research/medical education:

Premature rupture of membranes defined as amniotic fluid leakage before starting the labor pain. If PROM occurred before term pregnancy it is named as preterm premature rupture of membranes (PPROM). PPRM is associated with neonatal and maternal morbidity and mortality. It has been reported that premature ruptured membranes have less collagen content. Vitamin C is involved in collagen synthesis, secretion, and collagenolysis processes. Some studies have shown that there were lower levels of vitamin C in serum and amniotic fluid of cases with PPRM. The aim of this study is to evaluate the role of vitamin C in prevention of PPRM.

► Please cite this paper as:

Hafizi Lotfabad L. The Role of Vitamin C in Prevention of Preterm Premature Rupture of Membranes. Iran Red Cres Med J. 2013;15(2): 113-6. DOI: 10.5812/ircmj.5138

► Copyright © 2013, Iranian Red Crescent Medical Journal; Published by Kowsar Corp.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

branes and therefore tension resistance is decreased (5). However, PPRM should be considered important because of intrauterine infection risk (6). Smoking, previous PPRM, intrauterine infection, bacterial vaginosis, multifetal gestations, cervical shortening, hydramnios, and inadequate availability of some nutrients during pregnancy such as copper, zinc, magnesium,  $\beta$ -carotenes, vitamin E, and vitamin C have been identified as risk factors for PPRM and low birth-weight (7-9). Vitamin C is involved in collagen synthesis, collagen secretion, and collagenolysis processes (10). Occurrence of PPRM has been associated with changed patterns of collagen synthesis and decreased concentration of vitamin C at 28th weeks of gestation (11). It is reported that PROM could be used as a functional test of vitamin C status during pregnancy (12). Some studies have shown that there were lower levels of ascorbic acid (vitamin C) in serum, leucocytes, and amniotic fluid of cases with PPRM as compared to the control group (13). But they have provided little information about the relationship between vitamin C intake and its role in PPRM. The aim of this study is to evaluate the role of vitamin C in prevention of PPRM.

## 2. Objectives

The aim of this study is to evaluate the role of vitamin C supplementation in prevention of PPRM in women with a positive history of PPRM.

## 3. Patients and Methods

One hundred and seventy pregnant women with 14 weeks gestational age and the history of at least one PPRM with singleton pregnancy were enrolled in this clinical trial, after being completely explained about study conditions and signing informed consent. The study was approved by the Ethics Committee of Mashhad University of Medical sciences, and was performed in Imam-Reza University Hospital during a time period of 2008 to 2010. Inclusion criteria was the history of at least one PPRM in previous pregnancies, body mass index (BMI) of 18.5-30 kg/m<sup>2</sup>, singleton pregnancy, normal fetus and normal amniotic fluid in sonography, mother age of 18-35yrs, normal cervix length (more than 25mm), no Tobacco usage, and no consumption of vitamin C supplements. Patients were randomly divided into two groups:

For 85 women in the case group, 400 $\mu$ g folic acid daily was prescribed in the first trimester, then iron tablet containing 30 mg elemental iron and chewing tablet of 100 mg vitamin C (Iran-Darufactory) daily were added from 14th weeks of gestation (2) and was continued up to 37th weeks.

In control group, 85 patients were treated similar except for chewing tablet of placebo instead of same shape of vitamin C tablets. A questionnaire was completed for each woman including age, weight, history of previous disease, history of previous pregnancies, and gestational age of PPRM at previous pregnancy. The patients were

evaluated for bacterial vaginosis in the first visit, moreover anytime during the survey whenever the patient has been complaining of vaginosis signs, and if discovered it was treated medially. Sonography was performed for all of the cases during 12-14 weeks to evaluate the length of cervix and number of fetuses. All of the patients were evaluated monthly during the second trimester. At the end of second trimester, they were evaluated with sonography for the volume of amniotic fluid and fetal anomalies. The fetal membrane rupture was obtained by sterile speculum and observing amniotic fluid existing from the cervix, fern test or Nitrazin test. After delivery, the questionnaire was completed by data such as the date and cause of referring to the hospital, date and time of rupture of membranes, gestational age, neonate's sex, birth-weight, and five minute Apgar score. Data was analyzed by SPSS software (version 11). For quantitative variables, T test, and variance analysis were used if the variables were parametric in Kolmogorov-Smirnov test. Otherwise, Mann-Whitney and Cross-Calvaistests were used. Chi-square test was used for qualitative variables.  $P \leq 0.05$  was considered statistically significant.

## 4. Results

There was no significant difference between case and control groups in basic interventional factors (Table 1). Rupture of membranes was significantly decreased in the case group in the shape of PPRM, PROM and during term labor. Mean latency period was significantly ( $P = 0.002$ ) higher in case group ( $19.02 \pm 6.1$  hours) vs. control group ( $13.42 \pm 6.3$  hours). Both groups were statistically different in the view of gestational age, birth-weight, and neonatal Apgar score (Table 2). In the case group, 61 patients (71.8%) had vaginal delivery and 24 patients (28.2%) underwent cesarean section. it was similar in the control group, (60 women (70.6%) underwent vaginal delivery and 25 women (29.4%) cesarean section). ( $P = 1.007$ ). In the next step we compared neonatal outcomes between case and control groups in specific patient who were involved PPRM or PROM. The mean birth-weight of women with PROM was significantly higher in the case group (3068 gram) compared to the control (2968 gram) ( $P = 0.002$ ). The mean neonatal Apgar score in patients with PROM was 9.0 in the case group and 8.3 in the control group ( $P = 0.03$ ). The mean gestational age at delivery was 38.1 weeks in case group and 37.7 weeks in control group ( $P = 0.02$ ). Mean of latency period was  $3.6 \pm 2.9$  hours in case group and  $3.3 \pm 3.6$  hours in control group ( $P \leq 0.001$ ). The mean birth-weight of women with PPRM was 2370 grams in case and 2102 grams in control group ( $P \leq 0.001$ ). The mean neonatal Apgar score was 8 in patients with PPRM in case group and 7.3 in control group ( $P \leq 0.001$ ). The mean of latency period was  $59.2 \pm 38.1$  hours in case group and  $27.5 \pm 14.6$  hours in control group. ( $P \leq 0.001$ ) The mean gestational age at the time of delivery was  $34.7 \pm 1.2$  weeks in case group and  $33.3 \pm 1.4$  weeks in

control group ( $P \leq 0.001$ ). All the mentioned differences were statistically significant.

**Table 1.** Comparison of Interventional Factors Between Case and Control Groups Interventional

Interventional Factors, Mean	Case Group	Control Group	P value
Age, y	29.8	29	0.132
Number of Pregnancy	2.5	2.6	0.99
BMI, Kg/ m <sup>2</sup>	21.4	20.8	0.4
Gestational age at PPROM in previous pregnancy	31.05	31.4	1.70
Number of Dead neonates	0.3	0.6	1.88
Number of Abortion	0.14	0.05	0.62

**Table 2.** Comparison of Pregnancy and Neonatal Outcomes Between Case and Control Groups

Outcome	Case Group	Control Group	P value
<b>Pregnancy, No. (%)</b>			
PPROM	27 (31.8)	38 (44.7)	0.000
PROM	16 (18.8)	29 (34.1)	0.000
Normal <sup>a</sup>	42 (49.4)	18 (21.2)	0.000
<b>Neonatal, Mean <math>\pm</math> SD</b>			
Mean gestational age at delivery (week)	37.1 $\pm$ 1.9	35.9 $\pm$ 2.8	0.000
Mean birth weight, gram	2840 $\pm$ 382	2630 $\pm$ 529	0.000
Neonatal Apgar score	8.4 $\pm$ 0.7	7.83 $\pm$ 0.7	0.000

<sup>a</sup> Rupture of membrane during term delivery

## 5. Discussion

PPROM has been known as the main cause of preterm delivery and associated with increased rates of neonatal and maternal morbidity and mortality (14-16). Although it has different causes, collagen metabolism is considered as the main factor in premature rupture of membranes. Vitamin C usage during pregnancy can modulate the collagen metabolism and cause the strength of amniochorion membranes. There are many factors that affect the availability of vitamin C during pregnancy. Simhan et al. reported decreased level of vitamin C in women with premature rupture of membranes (17). The results of this study showed that vitamin C usage in case group significantly increased the gestational age at delivery, neonatal Apgar score, birth-weight, and latency period. This finding was confirmed by the study of Barret et al. He concluded that the administration of 100mg of vitamin C in pregnant women after 20th weeks of gestation can significantly decrease the incidence of PROM and PPROM (18). Siega et al. showed that the rates of membranes rupture before 37 weeks is increased with decrease, although the relation was not statistically significant (1). In addition, Hajifoghaha et al. reported that the usage of vitamin C supplements after 20th weeks of gestation prevents of PPROM (19). Vermilion et al. performed a study that one group of pregnant women with the history of PPROM received vitamin C along with ferrous sulfate. Although the

incidence of iron shortage anemia was decreased, no difference was observed in term of PPROM between case and control group (20). Also, Casaneuva et al. reported no significant difference between two groups in the view of vitamin C intake and PPROM (21). These different results from the results of the present study may be due to the lower volume sample in their study. In Borna et al. study no statistically significant difference was reported between vitamin C supplements and placebo groups in terms of sepsis incidence, but resemble to our study, neonatal Apgar score and birth-weight was different between two groups (22). Vitamin C is an essential nutrient, involved in several biochemical functions. It is an antioxidant that blocks the damaging effects of oxidative stress in vitro (23). Therefore, vitamin C can prevent premature rupture of membranes through its role as an antioxidant or in collagen synthesis and maintenance (24,25). The present study had some limitations, we aimed to study the independent effect of vitamin C, but because the serum level of vitamin C was not assayed, isolation of the effect of a single nutrient is difficult. We propose a relationship between low vitamin C intake and an increased risk of preterm premature rupture of membranes. Vitamin C supplement is recommended to be administered for pregnant women with the history of PPROM during pregnancy to prevent PPROM. Nevertheless, further studies with larger sample size and fewer limitations are needed to best clarify the role of vitamin C in

prevention of PPRM especially in women with other risk factors of PPRM.

## Acknowledgements

None declared.

## Financial Disclosure

None declared.

## Funding Support

None declared.

## References

1. Siega-Riz Anna Maria, Promislow Joanne HE, Savitz David A, Thorp John M, McDonald Thad. Vitamin C intake and the risk of preterm delivery. *American journal of obstetrics and gynecology*.2003;**189**(2):519-525.
2. Cunningham FG, Leveno KJ, Bloom SL, Hauth JC, Rouse DJ, Spong CY. *Management of preterm labor*. Williams obstetrics. 23th ed. New-york: MC Graw Hill; 2010.p. 232-47.
3. Mercer BM. Premature rupture of the membrane. In: Petraglia F, StrauSS GF, Gabbe SG, Wises G, editors. *Complicated Pregnancy*. 4th ed. London: Informa health care; 2007. p. 713-27.
4. Parry S, Strauss JF, 3rd. Premature rupture of the fetal membranes. *N Engl J Med*.1998;**338**(10):663-70.
5. Malak TM, Ockleford CD, Bell SC, Dalgleish R, Bright N, Macvicar J. Confocal immunofluorescence localization of collagen types I, III, IV, V and VI and their ultrastructural organization in term human fetal membranes. *Placenta*.1993;**14**(4):385-406.
6. Pasquier JC, Picaud JC, Rabilloud M, Claris O, Ecochard R, Moret S, et al. Neonatal outcomes after elective delivery management of preterm premature rupture of the membranes before 34 weeks' gestation (DOMINOS study). *Eur J Obstet Gynecol Reprod Biol*.2009;**143**(1):18-23.
7. Institute of Medicine Food and Nutrition Board. *Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids*. Washington DC: National Academy Press; 2000.
8. Khadem N, Mohammadzadeh A, Farhat AS, Valaee L, Khajedaluae M, Parizadeh SM. Relationship between Low Birth Weight Neonate and Maternal Serum Zinc Concentration. *Iran Red Crescent Med J*.2012;**14**(4):240-4.
9. Tejero E, Perichart O, Pfeffer F, Casanueva E, Vadillo-Ortega F. Collagen synthesis during pregnancy, vitamin C availability, and risk of premature rupture of fetal membranes. *Int J Gynaecol Obstet*.2003;**81**(1):29-34.
10. Pfeffer F, Casanueva E, Kamar J, Guerra A, Perichart O, Vadillo-Ortega F. Modulation of 72-kilodalton type IV collagenase (Matrix metalloproteinase-2) by ascorbic acid in cultured human amnion-derived cells. *Biol Reprod*.1998;**59**(2):326-9.
11. Bryant-Greenwood GD. The extracellular matrix of the human fetal membranes: Structure and function. *Placenta*.1998;**19**(1):1-11.
12. Casanueva E, Ripoll C, Tolentino M, Morales RM, Pfeffer F, Vilchis P, et al. Vitamin C supplementation to prevent premature rupture of the chorioamniotic membranes: a randomized trial. *Am J Clin Nutr*.2005;**81**(4):859-63.
13. Hadley CB, Main DM, Gabbe SG. Risk factors for preterm premature rupture of the fetal membranes. *Am J Perinatol*.1990;**7**(4):374-9.
14. Mercer BM, Rabello YA, Thurnau GR, Miodovnik M, Goldenberg RL, Das AF, et al. The NICHD-MFMU antibiotic treatment of preterm PROM study: impact of initial amniotic fluid volume on pregnancy outcome. *Am J Obstet Gynecol*.2006;**194**(2):438-45.
15. Martin RJ, Fanaroff AA, Walsh ME. *Fanaroff and Martins neonatal perinatal medicine*. 8th ed. Philadelphia: Frank Polizzano; 2006.p. 1097.
16. Petralia GF, Strauss GF, Gabbe SG, Wises G. *Complicated pregnancy*. 4th ed. London: In forma health care; 2007. p. 852-85.
17. Simhan HN, Canavan TP. Preterm premature rupture of membranes: diagnosis, evaluation and management strategies. *BJOG*.2005;**112** Suppl 1:32-7.
18. Barrett BM, Sowell A, Gunter E, Wang M. Potential role of ascorbic acid and beta-carotene in the prevention of preterm rupture of fetal membranes. *Int J Vitam Nutr Res*.1994;**64**(3):192-7.
19. Hajifoghaha M, Keshavarz T. Vitamin C supplementation and PROM. *Iran J Obstet Gynecol Infertil*.2008;**11**(2):33-9.
20. Vermillion ST, Kooba AM, Soper DE. Amniotic fluid index values after preterm premature rupture of the membranes and subsequent perinatal infection. *Am J Obstet Gynecol*.2000;**183**(2):271-6.
21. Casanueva E, Magana L, Pfeffer F, Baez A. Incidence of premature rupture of membranes in pregnant women with low leukocyte levels of vitamin C. *Eur J Clin Nutr*.1991;**45**(8):401-5.
22. Borna S, Borna H, Khazardoost S, Hantoushzadeh S. Perinatal outcome in preterm premature rupture of membranes with Amniotic fluid index < 5 (AFI < 5). *BMC Pregnancy Childbirth*.2004;**4**(1):15.
23. Gopalani S, Krohn M, Meyn L, Hitti J, Crombleholme WR. Contemporary management of preterm premature rupture of membranes: determinants of latency and neonatal outcome. *Am J Perinatol*.2004;**21**(4):183-90.
24. Fortunato SJ, Menon R. Distinct molecular events suggest different pathways for preterm labor and premature rupture of membranes. *Am J Obstet Gynecol*.2001;**184**(7):1399-405.
25. Plessinger MA, Woods JR, Jr, Miller RK. Pretreatment of human amnion-chorion with vitamins C and E prevents hypochlorous acid-induced damage. *Am J Obstet Gynecol*.2000;**183**(4):979-85.