

# Genetic Analysis of *cagA* and *vacA* Genes in *Helicobacter Pylori* Isolates and Their Relationship with Gastroduodenal Diseases in the West of Iran

Negar Souod<sup>1,\*</sup>, Mohammad Kargar<sup>2</sup>, Abbas Doosti<sup>3</sup>, Reza Ranjbar<sup>4</sup>, Meysam Sarshar<sup>4</sup>

<sup>1</sup> Young Researchers and Elite Club, Jahrom Branch, Islamic Azad University, Jahrom, IR Iran

<sup>2</sup> Department of Microbiology, Jahrom Branch, Islamic Azad University, Jahrom, IR Iran

<sup>3</sup> Department of Biotechnology, Shahrekord Branch, Islamic Azad University, Shahrekord, IR Iran

<sup>4</sup> Molecular Biology Research Center, Baqiyatallah University of Medical Sciences, Tehran, IR Iran

\*Corresponding author: Negar Souod, Young Researchers and Elite Club, Jahrom Branch, Islamic Azad University, Jahrom, IR Iran Tel: +98-3813361045, Fax: +98-3813361064, E-mail: negarsouod@yahoo.com.

## ABSTRACT

**Background:** *Helicobacter pylori* have different virulence factors which are associated with several gastroduodenal diseases; however, this association is variable in different geographical regions. Data of genotypes of Iranian *H. pylori* isolates are few.

**Objectives:** The aim of the current study was to investigate the *cagA/vacA* genotypes of *Helicobacter pylori* isolates and determine the relationship between these genotypes with respect to different gastric disorders in patients of Chaharmahal Bakhtiarian.

**Materials and Methods:** In this cross-sectional study, gastric biopsies were taken from 200 patients with gastroduodenal diseases. Histopathological features were recognized by specialist. The samples were subjected to PCR for detection and identification of *ureC*, *cagA* and *vacA* genes.

**Results:** The frequency of the *vacA* genotypes, *sa1/m1*, *sa1/m1b*, *sa1/m2*, *s1b/m1a*, *s1b/m1b*, *s1b/m2*, *s1c/m1a*, *s1c/m1b*, *s1c/m2*, *s2/m1a*, *s2/m1b* and *s2/m2* were 27(6.6%), 8(4.3%), 45(28.04%), 7(3.7%), 5(2.5%), 10 (6.1%), 12 (7.4%), 4 (2.5%), 18(11%), 6(3.7%), 0 and 22(13.5%) respectively. The *cagA* gene was detected in 92% of strains. Based on our findings, it seemed that *cagPAI* and *vacA s1* genotypes were associated with some gastric disorders in patients with *H. pylori*. In this region, the isolates carrying *sa1/m2* were the most prevalent.

**Conclusions:** We found considerable relationship between *sa1/m1a*, *sa1/m2*, *s2/m2* and *s1c/m1a* and some gastric disorders. Further studies about the role of *H. pylori* virulence factors and gastric disorders were recommended.

**Keywords:** *Helicobacter Pylori*; *VacA* Protein, *Helicobacter Pylori*; *CagA* protein, *Helicobacter Pylori*

Copyright © 2013, Iranian Red Crescent Medical Journal; Published by Kowsar Corp.

## 1. Background

*Helicobacter pylori* is major causes of chronic gastritis. They are also involved in the pathogenesis of several diseases including gastric and duodenal ulcers, gastric adenocarcinoma and mucosa-associated lymphoid tissue lymphoma (1). Several *H. pylori* virulent genes have been identified to contribute to the severity of these diseases.

The pathogenicity island (PAI) is the most important virulent factor of the bacterium that encodes a type IV secretion apparatus. The *cagA* gene is located at the end of the *cag* PAI (2-5). Another important virulence factor of *H. pylori* is a vacuolating cytotoxin (*VacA*), which is associated with injury of epithelial cells. The *vacA* gene is present in nearly all strains of *H. pylori* but it is poly-

►Article type: Research Article; Received: 10 Dec 2011; Revised: 09 Jan 2013; Accepted: 20 Jan 2013; Epub: 05 May 2013; Ppub: 05 May 2013

►Implication for health policy/practice/research/medical education:

There are geographic variations in *H. pylori* strains and different strains may cause distinct gastric diseases in different areas. Unfortunately, existing data is contradictory and cannot explain the pathogenic role of this bacterium in the development of different gastric diseases. Furthermore, it might be useful to know the genetic diversity of *H. pylori* strains in Chaharmahal Bakhtiari, one of the risky provinces of Iran with regard to eradicate dangerous strains.

►Please cite this paper as:

Souod N, Kargar M, Doosti A, Ranjbar R, Sarshar M. Genetic Analysis of *cagA* and *vacA* Genes in *Helicobacter Pylori* Isolates and Their Relationship with Gastroduodenal Diseases in the West of Iran. *Iran Red Cres Med J*. 2013; 15(5): 371-5. DOI: 10.5812/ircmj.3732

►Copyright © 2013, Iranian Red Crescent Medical Journal; Published by Kowsar Corp.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

morphic, comprising variable signal regions (type s1 or s2) and mid-regions (type m1 or m2). The type s1/m1 *vacA* strains causes more epithelial cell damage than type s1/m2, whereas type s2/m2 and the rare s2/m1 are non-toxic due to the presence of a short 12-residue hydrophilic extension on the s2 form (3, 6). The s-region is classified into s1 and s2 types and the m-region into m1 and m2 types. The s1 type is further classified into s1a, s1b and s1c subtypes, and the m1 into m1a and m1b subtypes. The mosaic combination of s and m-region allelic types determines the particular cytotoxic and, consequently, the pathogenicity of the bacterium (7, 8). There is a geographical variation in the *vacA* genotypes (9, 10). For example, studies have consistently shown that *vacA* s1a strains predominate in northern Europe, s1b in Central and South America, Spain and Portugal, s1a and s1b in the USA and s1c in East Asia (10). These differentiations may cause variations in the prevalence of gastric diseases in these areas. Thus, existing data are contradictory and cannot explain the pathogenic role of *H. pylori* in the development of different gastric diseases. Furthermore, it might be useful to know the genetic diversity of *H. pylori* strains in Chaharmahal Bakhtiari, one of the risky provinces of Iran.

## 2. Objectives

The aim of the current study was to investigate the *cagA/vacA* genotypes of *Helicobacter pylori* isolates and determine the association between these genotypes with different gastric disorders in Chaharmahal Bakhtiarian patients.

## 3. Materials and Methods

### 3.1. Collection of Patient Samples

From June to November 2009, 200 consecutive patients with dyspeptic symptoms attending the endoscopy suite of gastroenterology section of Hospital of Shahrekord University of Medical Sciences (SUMS) enrolled in the study. The questionnaires, including medical history and demographic data, were recorded for each patient. All studied patients signed an informed consent form before endoscopy and declared their willingness to allow the application of their anonymous data for research purposes. For each patient, two biopsy specimens were taken from the antrum using a disinfected endoscope. One piece of each specimen was examined by Rapid Urease Test (RUT) for detection of *H. pylori*. RUT was performed with a Gastro urease kit (Baharafshan, Iran). The second piece from positive samples in RUT was placed in 0.1 ml of sterile saline solution and was sent to Biotechnology Research Center of Islamic Azad University, Shahrekord Branch for further studies.

### 3.2. Genomic DNA Extraction and Polymerase Chain Reaction

DNA was isolated from biopsy specimens using Genomic DNA purification kit (DNPTM, CinnaGen, Iran) according to the recommendations of manufacture Primers sequences used for the PCR include as follows: ET-2U (5'-CCCTCACGC-CATCAGTCCCAAAA-3') and ET-2L (5'-AAGAAGTCAAAAAC-GCCCCAAAAC-3') (4). Primers used for PCR assays of *vacA* and *cagA* genes are listed in (Table 1) (11, 12).

**Table 1.** Primers Used for PCR Analysis of *vacA* and *cagA* Genes (,)

Region	Primer	Sequence (5'-3')	Size and location of PCR Product
ureC (glmM)	GlmM1-R GlmM1-F	GCTTACTTTCTAACACTAACGCGC GGATAAGCTTTAGGGGT-GTTAGGGG	296bp
s1a	<i>vacA</i> s1a-F VA1-R	CTC TCG CTT TAG TAG GAG C CTG CTT GAA TGC GCC AAA C	213 bp (843-1055)
s1b	SS3-F VA1-R	AGC GCC ATA CCG CAA GAG CTG CTT GAA TGC GCC AAA C	187 bp (869-1055)
s1c	<i>vacA</i> s1c-F VA1-R	CTC TCG CTT TAG TGG GGY T CTG CTT GAA TGC GCC AAA C	213 bp (843-1055)
s2	SS2-F VA1-R	GCT AAC ACG CCA AAT GAT CC CTG CTT GAA TGC GCC AAA C	199 bp (433-631)
m1a	VA3-F VA3-R	GGT CAA AAT GCG GTC ATG G CCA TTG GTA CCT GTA GAA AC	290 bp (2741-3030)
m1b	VAm-F3 VAm-R3	GGC CCC AAT GCA GTC ATG GA GCT GTT AGT GCC TAA AGA AGC AT	291 bp (2741-3031)
m2	VA4-F VA4-R	GGA GCC CCA GGA AAC ATT G CAT AAC TAG CGC CTT GCA	352 bp (976-1327)
cagA	cagA-U cagA-L	GGA ATA CCA AAA ACG CAA AAA CCA CCC CAC AAT ACA CCA GCA AAA CT	300bp

DNA samples from *H. pylori* (D0008, Genekam, Germany) were used as a positive control of *cagA* and *vacA* genes, and sterile distilled water was used as a negative

control. PCR was done in 20 µL (for *H. pylori*) or 25 µL (for *vacA* and *cagA*) of total reaction volume containing 1.5 mM MgCl<sub>2</sub> (2.0 mM for *cagA*), 50 mM KCl, 10 mM Tris-HCl

(pH 9.0), 0.1% Triton X-100, 200 μM dNTPs each (Fermentas), 0.4 μM primers, 0.3 U of Taq DNA polymerase (Fermentas), and 2 μL (40-260 ng/μL) of DNA. PCR was performed in a DNA Thermal Cycler (Eppendorf Mastercycler 5330, Eppendorf-Nethel-Hinz GmbH, Hamburg, Germany), with 40 cycles for ET2 primer and 35 cycles for *vacA* and *cagA* primers. Each cycle consisted of denaturation at 95°C/45 seconds; annealing at 59°C/30 seconds for ET2, 52°C/45 seconds for *vacA*, and 58°C/45 seconds for *cagA*; and extension at 72°C/45 seconds (11). There was another time extension (6 minute) at 72°C. PCR products were visualized by electrophoresis in 1% agarose gel, stained with ethidium bromide, and examined under ultraviolet illumination.

### 3.3. Statistical Analysis

The data were analyzed by SPSS software (Version 17.spss Inc, USA) and P value was calculated using Chi-square and Fisher's exact tests to find the significant relationship. P value less than 0.05 was statistically significant.

## 4. Results

Out of 200 gastric biopsy specimens, 164 (82%) were confirmed to be *H. pylori* infection positive by RUT. Of all patients studied, 79 (48.1%) were male and 85 (51.8%) were female, with a mean age of 47 ± 17 years (Range 15 to 88 years old). Sixteen patients (11.8%) had Gastric ulcers, 22 (16.2%) had Duodenal ulcers, 160 (97.5%) had Gastritis, 3 (2.2%) had Gastric cancer and 3 (2.2%) had Duodenitis. Possible combinations of *vacA* s and m regions were determined in Iranian population. Overall 63 samples were classified as *vacA* s1/m1, 73 samples as s1/m2, 22 as s2/m2 and 6 as s2/m1 genotypes. Out of 135 s1 strains, all the samples were successfully sub-typed using s1a, s1b and s1c specific primers. Among them, 79 (48.1%) were s1a positive, 21 (12.8%) were s1b positive and 35 (21.3%) were s1c positive. In the case of m1 sub-typing, the distribution of m1a and m1b was 31.7% and 9.14% respectively. M2 was found in 58.5% of the cases (Table 2).

**Table 2.** Correlation Between *vacA* Subtypes and Clinical Outcomes

<i>vacA</i> genotypes	G.U <sup>a</sup> (n = 16), No. (%)	D.U <sup>a</sup> (n = 22), No. (%)	G.C <sup>a</sup> (n = 3), No. (%)	C.G <sup>a</sup> (n = 160), No. (%)	DUO <sup>a</sup> (n = 3), No. (%)	Total (n = 164), No. (%)
S1a	9 (56.2)	11 (50)	1 (33.3)	76 (47.5)	1 (33.3)	79 (48.1)
S1b	3 (18.7)	1 (4.5)	1 (33.3)	20 (12.5)	1 (33.3)	21 (12.8)
S1c	3 (18.7)	3 (13.6)	0	34 (21.2)	0	35 (21.3)
S2	1 (6.2)	7 (31.8)	1 (33.3)	28 (17.5)	1 (33.3)	29 (17.6)
M1a	4 (25)	9 (40.9)	2 (66.6)	52 (32.5)	0	52 (31.7)
M1b	3 (18.7)	3 (13.6)	0	15 (9.3)	0	15 (9.1)
M2	9 (56.2)	10 (45.4)	1 (33.3)	92 (57.5)	3 (100)	97 (59.1)
S1a/m1a	1 (6.2)	4 (18.1)	1 (33.3)	27 (16.8)	0	27 (16.4)
S1a/m1b	2 (12.5)	2 (9.09)	0	7 (4.3)	0	8 (4.8)
S1a/m2	6 (37.5)	2 (9.09)	0	41 (25.6)	1 (33.3)	45 (27.4)
S1b/m1a	2 (12.5)	0	0	6 (3.7)	0	7 (4.2)
S1b/m1b	0	0	0	4 (2.5)	0	5 (3.04)
S1b/m2	1 (6.2)	1 (4.5)	1 (33.3)	10 (6.2)	1 (33.3)	10 (6.09)
S1c/m1a	1 (6.2)	1 (4.5)	0	12 (7.5)	0	12 (7.3)
S1c/m1b	1 (6.2)	1 (4.5)	0	4 (2.5)	0	4 (2.4)
S1c/m2	1 (6.2)	1 (4.5)	0	18 (11.2)	0	18 (10.9)
S2/m1a	0	1 (4.5)	1 (33.3)	6 (3.7)	0	6 (3.6)
S2/m1b	0	0	0	0	0	0
S2/m2	1 (6.2)	6 (27.2)	0	22 (13.7)	1 (33.3)	22 (13.4)

<sup>a</sup> Abbreviation: C.G, Chronic gastritis; D.U, Duodenal ulcer; DUO, Duodenitis; G.C, Gastric cancer; G.U, Gastric ulcer

One hundred fifty one (92%) out of 164 ureC- positive samples carried *cagA* gene. One hundred twenty eight (78.04%) strains with *vacA* s1 genotype were *cagA* positive while 23 (14.02%) strains with *vacA* s2 genotype were *cagA* positive, indicating that the presence of *cagA* gene was

significantly associated with the *vacA* s1 genotype (P = 0.004). In particular, most samples (95.5%) with the *vacA* s1/m2 genotypes were *cagA* positive. Also, the prevalence of *cagA* gene was not related to the clinical outcomes (Table 3).

**Table 3.** Correlations Between *cagA* Status and Clinical Outcomes

Patients groups	<i>H. pylori</i> -positive (n = 164), No. (%)	<i>cagA</i> -negative (n = 13), No. (%)	<i>cagA</i> -positive (n = 151), No. (%)
G.U <sup>a</sup>	16 (9.7)	2 (15.3)	14 (9.2)
D.U <sup>a</sup>	22 (13.4)	2 (15.3)	20 (13.2)
G.C <sup>a</sup>	3 (1.8)	1 (7.6)	2 (1.3)
C.G <sup>a</sup>	160 (97.5)	147 (94.2)	12 (7.5)
DUO <sup>a</sup>	3 (1.8)	0	3 (1.9)

<sup>a</sup> Abbreviation: C.G, Chronic gastritis; D.U, Duodenal ulcer; DUO, Duodenitis; G.C, Gastric cancer; G.U, Gastric ulcer

## 5. Discussion

*H. pylori* are one of the most genetically diverse bacterial species which may be involved in the complex variety of gastro duodenal diseases in infected patients all over the world (13-16). The geographic prevalence of distinct *H. pylori* genotypes remains largely unknown (15). For instance, in Japan, South America, Turkey and Pakistan, the prevalence is more than 80%, while in Scandinavia and England, the prevalence is between 20% and 40% (3). The prevalence of this bacterium in Iran is 60-90%, indicating that Iran is a highly risky region for *H. pylori* infection. The prevalence of this bacterium was 82% in our study indicating that our findings are consistent with previous reports in Iran (16-18). The *vacA* genotypes show considerable variability in different geographic regions (3). According to our results, 80% of samples had *vacA* s1a, b, c, also m2 genotypes and s1a/m2 was predominant in *H. pylori* isolates. This finding is somewhat similar to Europe and North America, where *vacA* s1a, s1b and m2 are predominating too. Our isolates were similar to those isolated from East Asian isolates where s1c is predominant (15). This study showed that Iranian *H. pylori* isolates are very diverse in genotype and contain the East and the West elements. *H. pylori* strains concluding *cagA* gene are more virulent than *cagA*-negative strains (3). The prevalence of *cagA*-positive *H. pylori* varied from one geographic region to another, e.g., 97% in Korea, 94% in Malaysia, 90% in China, 78% in Turkey and 53% in Kuwait (19). In this study, we found *cagA* gene in 92% of the *H. pylori*-positive population. This finding did not demonstrate the role of *cagA* as predictive marker for increased virulence feature of *H. pylori*, because of the high positivity of this gene in all *H. pylori* isolates. A strong association between the *cagA* and *vacA* status and peptic ulcer disease has been reported (20-22). Beil et al. suggested that the increased inhibitory effect of *cagA*-positive, cytotoxin-producing strains on mucin synthesis could be considered as a possible mechanism which is responsible for the increased risk of developing peptic ulceration with these *H. pylori* strains (9). Gzyl et al. found that *cagA* gene correlated with active gastritis in infected children and adults. They also found that the majority of *H. pylori* strains carrying s1/m2 *vacA* alleles were responsible for the higher levels of cytotoxin production (10). Our data were in agreement

with those of Gzyl et al. and Beil et al., which suggest that *H. pylori* strains with *cagA* and *vacA* s1 genotypes are associated with more severe gastritis (9, 10). Investigator's opinions about the association between *vacA* genotypes and gastric disorders were different. For example, in Iran, Jafari et al. found no correlation between them, (3) whereas Mohammadi et al. and Molaei et al. found that s1a allele were associated with more severe inflammation (15, 23). As results showed, we found an association between some diseases and some *vacA* genotypes but we couldn't introduce any allele as a marker of a disease. For instance, s2m2 strains that are non-toxicogenic in most regions of the world and are associated with NUD diseases were surprisingly more prevalent in PUD than in NUD patients and had direct association with duodenal ulcer in our evaluation. In Thailand, Japan, Korea, Colombia and America, no association had been found (21) whereas in Cuba, Lebanon, Hung Kung, China and most of the European countries, a significant association between s1 allele and PUD diseases had been reported (20, 21, 24, 25). Data analysis revealed a significant association between Duodenal ulcer and s1a/m1a ( $P = 0.04$ ), s1a/m2 ( $P = 0.02$ ) and surprisingly s2m2 ( $P = 0.05$ ) genotypes. We observed an association between Gastritis diseases and s1c/m2 ( $P = 0.01$ ). We didn't find any relationship between other diseases with regard to *vacA* genotypes. In conclusion, we found that the *cagA*-positive s1/m2 *H. pylori* were dominant genotypes in the patients under study. The *cagA* gene positivity rate was probably not closely associated with severity of the disease. *H. pylori* strains including *vacA* s1 genotype were associated with more severe gastritis.

## Acknowledgements

The authors would like to thank Mr. Momeni and Mr. Ramezani at the Biotechnology Research Center of the Islamic Azad University of Shahrekord and Endoscopy Unit of Hajar Hospital of Shahrekord, for their sincere technical and clinical support.

## Authors' Contribution

Souod N and Kargar M defined the research theme; Doosti A and Kargar M designed methods and experiments; Doosti A, Souod N and Sarshar M carried out the

laboratory experiments; Ranjbar R and Souod N analyzed the data, interpreted the results and wrote the paper.

### Financial Disclosure

All authors declare that there is no financial disclosure.

### Funding Support

The study was supported by Young researcher's club financially.

### References

1. Tanvir A, Khawar S, Muhammd R, Mukhtar M, Bilal R, Khanum A. Prevalence of Helicobacter pylori pathogenicity-associated *cagA* and *vacA* genotypes among Pakistani dyspeptic patients. *FEMS Immunol Med Microbiol*. 2009;55(1):34-8.
2. Iamaroon A, Chaimano S, Linpisarn S, Pongsiriwet S, Phornphutkul K. Detection of Helicobacter pylori in recurrent aphthous ulceration by nested PCR. *J Oral Sci*. 2003;45(2):107-10.
3. Jafari F, Shokrzadeh L, Dabiri H, Baghaei K, Yamaoka Y, Zojaji H, et al. *vacA* genotypes of Helicobacter pylori in relation to *cagA* status and clinical outcomes in Iranian populations. *Jpn J Infect Dis*. 2008;61(4):290-3.
4. Li C, Ha T, Ferguson DA, Jr, Chi DS, Zhao R, Patel NR, et al. A newly developed PCR assay of *H. pylori* in gastric biopsy, saliva, and feces. Evidence of high prevalence of *H. pylori* in saliva supports oral transmission. *Dig Dis Sci*. 1996;41(11):2142-9.
5. Peach HG, Pearce DC, Farish SJ. Helicobacter pylori infection in an Australian regional city: prevalence and risk factors. *Med J Aust*. 1997;167(6):310-3.
6. Argent RH, Thomas RJ, Letley DP, Rittig MG, Hardie KR, Atherton JC. Functional association between the Helicobacter pylori virulence factors *VacA* and *CagA*. *J Med Microbiol*. 2008;57(Pt 2):145-50.
7. Momtaz H, Souod N, Dabiri H. Comparison of the virulence factors of Helicobacter pylori isolated in stomach and saliva in Iran. *Am J Med Sci*. 2010;340(5):345-9.
8. Nahaei M, Sharifi Y, Taghi Akhi M, Ashghaezade M, Nahayei M, Fatahi E. Helicobacter pylori *cagA* and *vacA* genotypes and their relationship to peptic ulcer disease and non ulcer dysplasia. *Res J Microbiol*. 2008;3(5):386-94.
9. Beil W, Enss ML, Muller S, Obst B, Sewing KF, Wagner S. Role of *vacA* and *cagA* in Helicobacter pylori inhibition of mucin synthesis in gastric mucous cells. *J Clin Microbiol*. 2000;38(6):2215-8.
10. Gzyl A, Berg DE, Dzierzanowska D. Epidemiology of *cagA/vacA* genes in *H. pylori* isolated from children and adults in Poland. *J Physiol Pharmacol*. 1997;48(3):333-43.
11. Wang J, Chi DS, Laffan JJ, Li C, Ferguson DA, Jr, Litchfield P, et al. Comparison of cytotoxin genotypes of Helicobacter pylori in stomach and saliva. *Dig Dis Sci*. 2002;47(8):1850-6.
12. Yamazaki S, Yamakawa A, Okuda T, Ohtani M, Suto H, Ito Y, et al. Distinct diversity of *vacA*, *cagA*, and *cagE* genes of Helicobacter pylori associated with peptic ulcer in Japan. *J Clin Microbiol*. 2005;43(8):3906-16.
13. Farshad S, Alborzi A, Japoni A, Ranjbar R, Hosseini Asl K, Badiie P, et al. Antimicrobial susceptibility of Helicobacter pylori strains isolated from patients in Shiraz, Southern Iran. *World J Gastroenterol*. 2010;16(45):5746-51.
14. Izadi M, Fazel M, Sharubandi SH, Saadat SH, Farahani MM, Naseri MH, et al. Helicobacter species in the atherosclerotic plaques of patients with coronary artery disease. *Cardiovasc Pathol*. 2012;21(4):307-11.
15. Molaei M, Foroughi F, Mashayekhi R, Haghazali M, Zojaji H, Jafari F, et al. *CagA* status and *VacA* subtypes of Helicobacter pylori in relation to histopathologic findings in Iranian population. *Indian J Pathol Microbiol*. 2010;53(1):24-7.
16. Momtaz H, Souod N, Dabiri H, Sarshar M. Study of Helicobacter pylori genotype status in saliva, dental plaques, stool and gastric biopsy samples. *World J Gastroenterol*. 2012;18(17):2105-11.
17. Ghasemi A, Shirazi MH, Ranjbar R, Khorramizadeh MR, Daryani NE, Hosseini M. The prevalence of *cagA* and *cagE* genes in Helicobacter pylori strains isolated from different patient groups by polymerase chain reaction. *Pak J Biol Sci*. 2008;11(22):2579-83.
18. Hosseinzadeh M, Khosravi A, Saki K, Ranjbar R. Evaluation of Helicobacter pylori infection in patients with common migraine headache. *Arch Med Sci*. 2011;7(5):844-9.
19. Nahaei M, Sharifi Y, Taghi Akhi M, Ashghaezade M, Nahayei M, Fatahi E. Helicobacter pylori *cagA* and *vacA* genotypes and their relationship to peptic ulcer disease and non ulcer dysplasia. *Res J Microbiol*. 2008;3(5):386-94.
20. Torres L, Melián K, Moreno A, Alonso J, Sabatier C, Hernandez M, et al. Prevalence of *vacA*, *cagA* and *babA2* genes in Cuban Helicobacter pylori isolates. *World J Gastroenterol*. 2009;15(2):204-10.
21. Wong B, Yin Y, Berg DE, Xia HH, Zhang JZ, Wang WH, et al. Distribution of Distinct *vacA*, *cagA* and *iceA* Alleles in Helicobacter pylori in Hong Kong. *Helicobacter*. 2001;6(4):317-24.
22. Yamaoka Y, Kodama T, Gutierrez O, Kim JG, Kashima K, Graham DY. Relationship between Helicobacter pylori *iceA*, *cagA*, and *vacA* status and clinical outcome: studies in four different countries. *J Clin Microbiol*. 1999;37(7):2274-9.
23. Mohammadi M, Oghalaie A, Mohajerani N, Massarrat S, Nasiri M, Bennedsen M, et al. Prevalence of Helicobacter pylori vacuolating cytotoxin and its allelic mosaicism as a predictive marker for Iranian dyspeptic patients. *Bull Soc Pathol Exot*. 2003;96(1):3-5.
24. Khayat A, Soweid A, Kattar M, Tahwil A, El Hajj A, Azar C, et al. Prevalence and clinical relevance of Helicobacter pylori *cagA* and *vacA* genes in Lebanese patients with gastritis and peptic ulcer disease. *J Infect Develop Count*. 2007;1(1):55-61.
25. Wang J, van Doorn LJ, Robinson PA, Ji X, Wang D, Wang Y, et al. Regional variation among *vacA* alleles of Helicobacter pylori in China. *J Clin Microbiol*. 2003;41(5):1942-5.