

The Association of Health-Promoting Lifestyle With Quality of Life Among the Iranian Elderly

Tayebeh Rakhshani¹; Davood Shojaiezadeh^{1,*}; Kamran Bagheri Lankarani²; Fatemeh Rakhshani³; Mohammad Hossain Kaveh⁴; Najaf Zare⁵

¹Department of Health Education and Promotion, Tehran University of Medical Sciences, Tehran, IR Iran

²Health Research Center, Shiraz University of Medical Sciences, Shiraz, IR Iran

³Safety Promotion and Injury Prevention Research center, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

⁴Department of Health Education and Promotion, Shiraz University of Medical Sciences, Shiraz, IR Iran

⁵Department of Biostatistics, Shiraz University of Medical Sciences, Shiraz, IR Iran

*Corresponding Author: Davood Shojaiezadeh, Department of Health Education and Promotion, Tehran University of Medical Sciences, Tehran, IR Iran. Tel: +98-9196250473, E-mail: shojae5@yahoo.com

Received: February 23, 2014; Revised: April 26, 2014; Accepted: June 5, 2014

Background: As individuals live a longer life, health-promoting lifestyle (HPL) becomes even more essential, particularly with regard to maintaining functional independence and improving quality of life (QoL).

Objectives: This study aimed to determine the association between QoL and HPL in the Iranian elderly living in Shiraz.

Materials and Methods: This analytical cross-sectional study was conducted in 2013 at retirement centers of Shiraz City, Iran. The sample included 500 elderly who aged > 60 years. Proportional stratified random sampling was used to select the elderly from retirement centers. QoL was assessed by the Farsi version of Short Form Health Survey questionnaire (SF-36) and HPL was measured by health-promoting lifestyle profile (HPLP II). Data were analyzed using independent-samples t test, one-way ANOVA, Pearson's correlation, and stepwise multiple regression using SPSS 21.

Results: There were significant differences in QoL in terms of sex, age, education, and marital status. There were significant differences in HPL in terms of gender, age and education ($P < 0.05$). There was a statistically significant association between QoL and HPL in the elderly ($r = 0.42$, $P < 0.05$). Based on HPLP II constructs, the significant predicting factors of QoL in the elderly consisted of spiritual growth, stress management, and physical activity ($P < 0.05$).

Conclusions: Health providers should improve the QoL in the elderly by facilitating HPL through health-promoting interventions, which will maintain and increase physical activity, stress management, and spiritual growth.

Keywords: Quality of Life; Health-Promoting Lifestyle; the Elderly

1. Background

The world's population is aging (1). The improvement in the living conditions has led to the phenomena of aging in different societies (2, 3). Like other countries, Iran has experienced a shift in the population structure towards aging so that according to statistics in 2011, 8.19% of Iran's population were older than 60 years (4).

As aging is considered an important social issue worldwide, the biggest challenge is improving the quality of life (QoL) in the elderly (5, 6). QoL is a multidimensional concept that includes the individual's physical, psychological, and social performances (7). The increase of human's life span emphasizes the importance of the health-promoting behavior in maintaining and improving QoL (8-10). The World Health Organization has placed an emphasis on the importance of health-promoting behavior as a key strategy for maintaining a good QoL (11). Pender et al. classified the health-promoting lifestyle (HPL) into six subcategories of nutrition, physical activities, stress

management, health responsibility, interpersonal relations, and spiritual growth (12).

In Iran, researches have been conducted on different age groups, except the elderly, regarding the association between HPL and QoL (13, 14). In developed countries, the association of one or two subscales of HPL with QoL in the elderly have been investigated, which have mainly dealt with the aspect of nutrition and physical activity and the results have indicated that nutrition and physical activities are effective factors in QoL of the elderly (15-18).

A study in China indicated that interpersonal relations, spiritual growth, and physical activities were better predictors for QoL in older than 50 years of age retired people (19, 20). Despite some similarities between the elderly in Iran and the elderly in developed countries, the cultural, religious, and environmental issues should not be neglected (21). In Iran, the results of a study by Agha-nuri et al. has indicated that the quality of nutrition does

not have a significant association with the QoL in the elderly (22). On the contrary, researches in developed countries show a significant association between nutrition and QoL in the elderly (23-25). Knowledge about the factors that influence QoL is important as population aging becomes a worldwide reality (8). There is less evidence about the amount of influence of HPL on the health-related QoL (HRQoL) in the Iranian elderly and it should be considered that any designed and executed strategy for health promotion needs evidence-based science, because if there is no awareness and recognition of the status of the society's health factors, health programs may just impose high costs. Hence, the results of this study can help health specialists, managers, and policymakers to design and execute health-promoting strategies based on the evidences, and prepare a basis for further researches in the field of aging.

2. Objectives

This study aimed to investigate the association between HPL and its subscales (nutrition, physical activities, stress management, health responsibility, interpersonal relations, and spiritual growth) with QoL in the Iranian elderly. Moreover, this study familiarized us with the status of the elderly's QoL and HPL.

3. Materials and Methods

3.1. Design

This research was an analytical cross-sectional study to determine the association between QoL and HPL among the elderly in Shiraz, Iran, in 2013.

3.2. Participants and Setting

The research population included the retired elderly or their spouse that were members of the state retirement centers. The inclusion criteria were retired males or females with ≥ 60 years of age, who were able to speak, had no hearing impairment or any serious health problems, and were willing to participate in the study.

3.3. Sampling and Data Collection

A total of 45671 retired elderly were registered in the state retirement centers. The sample size was estimated at 500 by using Morgan formula. The sampling method was based on a proportionate stratified random sampling approach. We considered those retirement centers with the elderly members > 200 who had agreed to participate in the study as one stratum.

Considering the sample size, proportionate allocation sampling was used to identify a sampling fraction for each center. The participants were selected randomly from existed elderly members' list in each center. Participate in the study were interviewed face to face by trained interviewers with a standard questionnaires at the ap-

propriate time and place so there was no missing data in this study.

3.4. Measurement Tools

The Farsi version of Short Form Health Survey questionnaire (SF-36) was used to collect data on QoL. The total score of SF-36 ranges from zero to 100 and higher scores indicate a better condition. The reliability and validity of this scale has been approved for the elderly in some surveys (26, 27). The obtained Cronbach's alpha was 0.93 for QoL in this study. The Health-promoting Lifestyle Profile (HPLP II) is a 52-item rating scale and consists of six subscales to measure major aspects of a health promoting lifestyle including health responsibility, physical activity, nutrition, interpersonal relations, spiritual growth, and stress management. The mean can be obtained on the total scale as a measurement of HPL and a mean can be derived for each subscale. A mean score higher than the average score was considered as the better response by the authors. The reliability and validity of this scale has been approved for the elderly in some studies (27, 28). In this study, the specialist group assessed the content validity of the Farsi version of HPLP II scale and found it to be culturally relevant with the obtained alpha coefficient of 0.93.

3.5. Data Analysis

Frequencies, percentage, mean, and standard deviation were used to describe demographic characteristics, QoL, and HPL. Independent-samples t-test and ANOVA with post hoc test were used to determine the differences in QoL and HPL. Pearson's correlation coefficient was used to identify the degree of correlation between QoL with HPL and its subscales. In addition, stepwise multiple regressions were used to determine the predicting factors of QoL with respect to HPLP II subscales. A P value < 0.05 was considered as statistically significant. As part of the regression analysis, the assumption of normality, linearity, homoscedasticity, and independence of residuals were tested using residual scatter plots, correlation matrix, Durbin-Watson, tolerances, and variance inflation factor, respectively.

3.6. Ethical Considerations

The Ethic Committee of Tehran University of Medical Sciences approved the study protocol (date, 2013-05-19; ID, 240/32251). All participants gave their oral consents for interview. We kept the information of the participants confidential.

4. Results

Overall, 53.6% of the participants were male and 46.4% female. The highest frequencies belonged to the age group of 60 to 75 years (76.4%), the elementary level (39.4%), and the married (73%) in age groups, education level, and

Table 1. The Socio-Demographics Characteristics of the Elderly participants (n = 500)^a

Characteristics	Results
Sex	
Female	232 (46.4)
Male	268 (53.6)
Age	
< 75	382 (76.4)
≥ 75	118 (23.6)
Education	
Elementary	197 (39.4)
High School	190 (38)
Diploma	81 (16.2)
University Education	32 (6.4)
Marital Status	
Married	365 (73)
Single	17 (3.4)
Windowed/Divorced	118 (23.6)

^a Data are presented as No. (%).**Table 2.** The Range of Dimension of Health Promoting Lifestyle Profile in Elderly (n = 500)^{a,b}

Variables	Results	Median (Range)
Total HPLP II Score	122.78 ± 23.34	125.5 (52-208)
Physical Activity	14.66 ± 4.71	20 (8-32)
Stress Management	19.27 ± 4.4	20 (8-32)
Health Responsibility	19.6 ± 5.22	22.5 (9-36)
Nutrition	22.41 ± 4.93	22.5 (9-36)
Spiritual Growth	23.12 ± 5.6	22.5 (9-36)
Interpersonal Relationship	23.69 ± 4.62	22.5 (9-36)
Total SF-36 Score	50.88 ± 17.01	50 (0-100)

^a Data are presented as mean ± SD.^b Abbreviations: HPLP II; health-promoting lifestyle profile, SF-36; short form health survey questionnaire.**Table 3.** The Differences of Scores in Quality of Life and Health-Promoting Lifestyle Based on Demographic Variables in the Elderly (n = 500)^{a,b}

	Quality of Life (SF-36)			Health Promoting Lifestyle (PHLP II)		
	Mean ± SD	T a, F b Statistics	P Value	Mean ± SD	T a, F b Statistics	P Value
Sex		t = -6.66	0.00		t = -3.08	0.002
Male	55.4 ± 17.2			125.75 ± 23.96		
Female	45.7 ± 15.3			119.34 ± 22.16		
Age		t = 4.59	0.00		t = 2.37	0.01
< 75	52.79 ± 16.57			124.15 ± 22.54		
> 75	44.71 ± 17.02			118.34 ± 25.37		
Education		F = 17.34	0.00		F = 16.6	0.00
Elementary	45.5 ± 16.12			114.22 ± 20.03		
High School	52.65 ± 15.75			128.64 ± 21.42		
Diploma	56.63 ± 16.61			126.01 ± 26.75		
University Education	61.71 ± 18.66			134.5 ± 28.08		
Marital Status		F = 18.06	0.00		F = 2.33	0.09
Married	53.5 ± 16.63			124.14 ± 23.3		
Single	49.01 ± 24.51			118.64 ± 32.15		
Windowed/Divorced	43.06 ± 14.42			119.15 ± 21.71		
Total	50.88 ± 17.01			122.78 ± 23.34		

^a Acronyms: HPLP II, health-promoting lifestyle profile; and SF-36, short form health survey questionnaire.^b Independent-samples t-test (P < 0.05) and One-way ANOVA (P < 0.05) were used.

marital status, respectively (Table 1). Table 2 shows the possible ranges, means, and standard deviations of HPL, its subscales, and QOL. The mean QoL score of the participants was 50.8 ± 17.01 and the mean score of HPL was 122.78 ± 23.34. In addition, interpersonal relationship and spiritual growth had the highest score and physical activity had the lowest score in the HPLP II subscales of the el-

derly (Table 2). According to independent-samples t-test and ANOVA, there were significant differences in QoL and HPLP II scores in terms of sex, age, and education whereas marital status had significant differences on QoL (P < 0.05) not on HPLP II (P > 0.05). The means scores of QoL and HPLP II of male and age group < 75 years old were significantly higher than the means scores of QoL and HPLP

II of female and age group ≥ 75 years old ($P < 0.05$). Post hoc testing indicated that the means scores of QoL and HPLP II of those with lower education (Elementary education) were significantly lower than that of the other education levels in the elderly ($P < 0.05$). The mean score of QoL of married individuals was significantly higher than that of widowed/divorced ones ($P < 0.05$) (Table 3).

Pearson correlation coefficient was used to describe the magnitude and direction of the association between QoL and HPLP II subscales. There was a statistically significant correlation between QoL and HPLP II and its subscales in the elderly ($P < 0.001$) at the 0.01 level. Results are summarized in (Table 4).

Multiple linear regression analysis with stepwise method showed that three subscales of HPLP II including spiritual growth, stress management and physical activity were significant predicting factors of QoL in the elderly; they explained 31.4% of the variance of QoL while the other subscales of HPLP II explained 1% of it (adjusted $R^2 = 0.314$, R^2 change = 0.01). The level of statistical significance was at $P < 0.05$ (Table 5).

Table 4 . The Correlation Between Quality of Life and the Dimensions of Health Promotion Lifestyle Profile in the Elderly ($n = 500$)^a

Variable	QoL (SF-36 Score)	
	R ^b	P Value
Total HPLP II Score	0.47 ^c	0.00
Spiritual Growth	0.52 ^c	0.00
Stress Management	0.46 ^c	0.00
Physical Activity	0.41 ^c	0.00
Interpersonal Relationship	0.35 ^c	0.00
Health Responsibility	0.27 ^c	0.00
Nutrition	0.2 ^c	0.00

^a Abbreviations: QoL; quality of life, SF-36; short form health survey questionnaire, HPLP II; health-promoting lifestyle profile.

^b Pearson correlation coefficient.

^c $P < 0.01$.

Table 5. Predicting Factors of Quality of Life Among the Elderly by Stepwise Multiple Regression Analysis ($n = 500$)^a

Variable	Unstandardized Coefficients		Standardized Coefficients	t	P Value
	B	Std. Error	Beta		
Spiritual Growth	1.17	0.16	0.38	7.05	0.00
Stress Management	0.65	0.22	0.17	2.87	0.00
Physical Activity	0.59	0.17	0.16	3.45	0.00

^a $B_0 = 13.73$; $R = 0.56$; $R^2 = 0.32$; Adjusted $R^2 = 0.314$; R^2 change = 0.01; ($F = 58.19$, P value = 0.00).

5. Discussion

In general, the obtained scores of SF-36 in this study showed moderate QoL in the elderly. These result were compatible with the QoL of the elderly in Tehran (29). The obtained scores of HPLP II were a little lower than the moderate scores; the HPL condition in older than 50 years old Chinese people was similar to that of our study population (20). We could find no data on HPL condition among the elderly in the other cities of Iran; therefore, it is better to conduct further studies on the HPL status in the Iranian elderly.

In this study, the participants obtained fairly higher scores for spiritual growth as well as interpersonal relationship and fairly lower scores for nutrition, stress management, and health responsibility. In addition, the lowest score belonged to physical activity, which is emphasizing and contemplating. One possible reason may be that elderly do not consider health-promoting behavior as being necessary for healthy aging while the recent evidence supports that healthy lifestyle is an effective strategy for sound aging (30).

In the study on retired workers, the highest and lowest scores belonged to the interpersonal relations and health responsibility, respectively (20). A study in elderly

women determined that the lowest and highest scores belonged to physical activities and nutrition, respectively (31). With regard to these results, the elderly do not have enough health-promoting behavior, especially in physical activity, and further studies are needed to explore the cause of this problem. Considering the demographic features of the subjects, the QoL score was lower in females, older ages, single elderly, and those with lower educational level. These results were compatible with the results of many other studies in Iran and other countries (8, 29, 31-38). Moreover, HPLP II score was higher in males, < 75 years of age, and those with higher educational level. These results were compatible with the results in the other studies (19, 31). Therefore, it is better to pay attention seriously to the health status of the elderly that are female, older, and unmarried with lower education. Moreover, this study showed that HPL and all of its subscales are significantly associated with QoL in the elderly, which was consistent with those reported in China (19). These are similar with the results of other researches on other age groups in Iran (13, 14). The subscales with the highest correlations with HRQoL were spiritual growth, stress management, physical activity,

and interpersonal relations, consecutively. In addition, nutrition and health responsibility had the lowest correlations with HRQoL. These results are consistent with the results of the study by Zhang (19). The findings by Keller et al. in Canadian elderly indicated that nutrition is significantly associated with the QoL and suggested intervening studies for confirming these results (24). Considering different correlation degrees of HPL subscales with QoL, the stepwise multiple regression analysis showed that spiritual growth, stress management, and physical activity were stronger predictors of QoL. A longitudinal study in Florida introduced physical activity as an effective predictor of QoL in the elderly (39). Another study in Brazil concluded that physical activity was an effective factor for QoL (38). In Chinese elderly, health responsibility, spiritual growth, and physical activity were stronger predictors of QoL (20). Hence, physical activity is considered as a positive effective factor for QoL in elderly of different societies. A study in Spain is another proof to this claim that suggested education and training on physical activities for improving QoL in the elderly (40). Spiritual growth was one of the important factors for improving QoL and the score of spiritual growth was higher than average in this study. Hence, it can be concluded that the tendency of most Iranian elderly towards religious rites and Islamic ceremonies and spending most of their times in the mosques can be the reason for better scores of spiritual growth. Researches have shown that spiritual growth plays an important role on QoL in the elderly and is related to their health (41). Despite the advance of scientific literature, more studies are required about the influence of spiritual performance on health (42).

5.1. Final Conclusion

With regard to the low level of health-promoting behaviors, especially physical activities and stress management that were stronger predictors of QoL in this study, physical activities and stress management must be included in health interventions in order to improve QoL in the elderly. It is recommended that the health policymakers adopt more strategies for health promotion in the elderly women due to their low QoL as women have an important role in the development of the communities.

5.2. Limitation and Strong Points

This study was a cross-sectional study with small sample; therefore, results could not be generalized to the entire elderly population in Iran. Interventional or longitudinal studies are recommended to confirm these results. The strong point of the study was simultaneously determining the association of the six HPL subscales, i.e. health responsibility, physical activity, nutrition, interpersonal relations, spiritual growth, and stress management, with QoL in the elderly.

Acknowledgements

We appreciate the cooperation of retirement centers in Shiraz.

Financial Disclosure

The study was approved in May 2013 as thesis PhD and supported by Tehran University of Medical Sciences.

References

1. Noncommunicable Disease Prevention and Health Promotion Department. *Aging and Life Course*. Geneva: policy framework; 2002.
2. AK P. Determinants of Living Arrangements of Elderly in Orissa, India, Asia. *Pac Popul J*. 2010;25.
3. Kinsella K, Phillips DR. Global Aging: The Challenge of Success. *Population Bulletin*. *Popul Bulletin, Popul Reference Bureau*. 2005;1:1-5.
4. Statistical Yearbook of Islamic Republic of Iran Management and Planning Organization. Tehran. *Satake Center Ir J*. 2007.
5. Vameghi R, Niksirat Z, Hatamizadeh N, Kazemnejad A. Health-related Quality of Life in Retired Older People in Tehran City. *Journal of Applied Gerontology*. 2010;31(2):155-72.
6. Malekafzali H, Baradaran Eftekhari M, Hejazi F, Khojasteh T, Noot RH, Falahat K, et al. The Effectiveness of Educational Intervention in the Health Promotion in Elderly people. *Iran J Public Health*. 2010;39(2):18-23.
7. Revicki DA. Health-related quality of life in the evaluation of medical therapy for chronic illness. *J Fam Pract*. 1989;29(4):377-80.
8. Lee TW, Ko IS, Lee KJ. Health promotion behaviors and quality of life among community-dwelling elderly in Korea: a cross-sectional survey. *Int J Nurs Stud*. 2006;43(3):293-300.
9. Lucile C, Aure'lie M, Nicole C, Florence C, Ge'lie3 D, Ve'ronique D-P, et al. *Health Promotion In nursing practice*. Pender NJ MCPM editor. New Jersey: Prentice Hall PTR; 2005.
10. Resnick B. Health promotion practices of older adults: testing an individualized approach. *J Clin Nurs*. 2003;12(1):46-55.
11. World Health Organization. *Noncommunicable Diseases and Mental Health Cluster*. Geneva: a policy framework; 2002.
12. Pender NJ, Muraugh CL, Parsons MA. *Health Promotion In nursing practice*. New Jersey: Prentice Hall PTR; 2005.
13. Shankar R, Tuyethong N, Sallis JD. Atherogenesis. Mitigation of monocyte adhesion to arterial endothelium in hyperlipidemic rats by phosphocitrate, a phosphorylated polycarboxylic acid. *Atherosclerosis*. 1986;62(1):47-54.
14. Mohamadian H, Eftekhari H, Rahimi A, Mohamad HT, Shojaeizade D, Montazeri A. Predicting health-related quality of life by using a health promotion model among Iranian adolescent girls: a structural equation modeling approach. *Nurs Health Sci*. 2011;13(2):141-8.
15. Sohng KY, Sohng S, Yeom HA. Health-Promoting Behaviors of Elderly Korean Immigrants in the United States. *Public Health Nursing*. 2002;19(4):294-300.
16. Rejeski WJ, Mihalko SL. Physical Activity and Quality of Life in Older Adults. *Biological Sciences and Medical Sciences*. 2001;56(Supplement 2):23-35.
17. McNaughton SA, Crawford D, Ball K, Salmon J. Understanding determinants of nutrition, physical activity and quality of life among older adults: the Wellbeing, Eating and Exercise for a Long Life (WELL) study. *Health and Quality of Life Outcomes*. 2012;10(1):109.
18. Weiwen C, Claudio RN, Ian SP, Robert WM, Caroline H, Rod KD. fruit and vegetable consumption, and physical inactivity in a free living, multiethnic population in Hawaii. *International Journal of Behavioral Nutrition and Physical Activity*. 2010.
19. Zhang S, Tao F, Ueda A, Wei C, Fang J. The influence of health-promoting lifestyles on the quality of life of retired workers in a medium-sized city of Northeastern China. *Environmental Health and Preventive Medicine*. 2013;18(6):458-65.
20. Zhang SC, Tao FB, Ueda A, Wei CN, Fang J. The influence of health-

- promoting lifestyles on the quality of life of retired workers in a medium-sized city of Northeastern China. *Environ Health Prev Med*. 2013;**18**(6):458-65.
21. Sonay A. Quality of life for elderly people in iran with service design approach. 2012.
 22. Aghanuri A, Mahmoudi M, Asadi M, Mortaji F, Salehi H, Djafarian K, et al. Quality of life and its relationship with quality of diet among elderly people in urban areas of Markazi province, Iran, Arak . *Medic Uni J*. 2013;**15**(68):1-11.
 23. Kvamme JM, Olsen JA, Florholmen J, Jacobsen BK. Risk of malnutrition and health-related quality of life in community-living elderly men and women: the Tromso study. *Qual Life Res*. 2011;**20**(4):575-82.
 24. Keller HH, Ostbye T, Goy R. Nutritional risk predicts quality of life in elderly community-living Canadians. *J Gerontol A Biol Sci Med Sci*. 2004;**59**(1):68-74.
 25. Amarantos E, Martinez A, Dwyer J. Nutrition and Quality of Life in Older Adults. *Biol Sci Medic Sci*. 2001;**56**(Supplement 2):54-64.
 26. Montazeri A, Goshtasebi A, Vahdaninia M, Gandek B. The Short Form Health Survey (SF-36): Translation and validation study of the Iranian version. *Quality of Life Research*. 2005;**14**(3):875-82.
 27. Walker SN, Sechrist KR, Pender NJ. Health-Promoting lifestyle Profile II, University of , College of Nursing. *Neb Medic Center J*. 1995.
 28. Morowatisharifabad MA, Ghofranipour F, Heidarnia A, Ruchi G, Ehrampoush MH. Self-Efficacy and Health Promotion Behaviors of Older Adults in Iran. *Social Behavior and Personality*. 2006;**34**(7):759-68.
 29. Tajvar M, Arab M, Montazeri MA. Determinants of health-related quality of life in elderly in Tehran, Iran. *BMC Public Health*. 2008:8-323.
 30. Franklin NC, Tate CA. Lifestyle and Successful Aging: An Overview. *Americ Lifestyle Med J*. 2008;**3**(1):6-11.
 31. Pullen C, Walker SN, Fiandt K. Determinants of health-promoting lifestyle behaviors in rural older women. *Fam Community Health*. 2001;**24**(2):49-72.
 32. Nejati V, Shirinbayan P, Akbari Kamrani A, Foroughan M, Taheri P, Sheikhvatan M, et al. Quality of life in elderly people in Kashan, Iran. *Middel East J*. 2008;**5**(2).
 33. Ioanna TH, Dimitra TH, Dimitra K. The differences at quality of life and loneliness between elderly people. *JBE*. 2010;**6**(2).
 34. Canbaz S, Sunter AT, Dabak S, Peksen Y. The prevalence of chronic diseases and quality of Life in elderly people in Samsun. *Turk J Med Sci* . 2003;**33**(5):335-40.
 35. Borglin G, Jakobsson U, Edberg AK, Hallberg IR. Self-reported health complaints and their prediction of overall and health-related quality of life among elderly people. *Int J Nurs Stud*. 2005;**42**(2):147-58.
 36. Orfila F, Ferrer M, Lamarca R, Tebe C, Domingo-Salvany A, Alonso J. Gender differences in health-related quality of life among the elderly: the role of objective functional capacity and chronic conditions. *Soc Sci Med*. 2006;**63**(9):2367-80.
 37. Edit P, Ferenc B, Aranka K, Sandor B, Laszlo N. Determinants of health-promoting lifestyle behaviour in the rural areas of Hungary. *Health Promo Inter J*. 2010;**25**(3).
 38. Margareth GL, Marilisa BAB, Cheste RL, Luana C, Rozana MC. Health related quality of life among the elderly. Rio de Janeiro: based study using SF-36 survey. 2009. pp. 2159-67.
 39. Kahana E, Lawrence RH, Kahana B, Kercher K, Wisniewski A, Stoller E, et al. Long-term impact of preventive proactivity on quality of life of the old-old. *Psychosom Med*. 2002;**64**(3):382-94.
 40. Guallar-Castillón P, Santa-olalla Peralta P, Ramón Banegas J, López E, Rodríguez-Artalejo F. Actividad física y calidad de vida de la población adulta mayor en España. *Med Clínica J*. 2004;**123**(16):606-10.
 41. Dávid R, Hodge R, Bonifas P. Spirituality and Older Adults: Ethical Guidelines to Enhance Service Provision. *Advanc Soc Work*. 2010;**11**(1).
 42. Gillum F, Griffith DM. Prayer and Spiritual Practices for Health Reasons among American Adults: The Role of Race and Ethnicity. *Religion Health J*. 2009;**49**(3):283-95.