

Study of the Effects of Diazinon on Fetal Liver in BALB/c Mice

Fatemeh Saraei,¹ Mehrangiz Sadoughi,¹ Gholamreza Kaka,^{2,*} Seyed Homayoon Sadraie,³ and Mohsen Foaddodini⁴

¹Department of Biology, Islamic Azad University, North Tehran Branch, Tehran, IR Iran

²Neurosciences Research Center, Baqiyatallah University of Medical Sciences, Tehran, IR Iran

³Department of Anatomy, Baqiyatallah University of Medical Sciences, Tehran, IR Iran

⁴Department of Physiology, Birjand University of Medical Sciences, Birjand, IR Iran

*Corresponding author: Gholamreza Kaka, Neurosciences Research Center, Baqiyatallah University of Medical Sciences, Tehran, IR Iran. Tel/Fax: +98-2126127286, E-mail: gh_kaka@yahoo.com

Received 2015 February 23; Revised 2015 June 15; Accepted 2015 August 09.

Abstract

Background: Diazinon is an organophosphate that is broadly used as a pesticide to control insects and environmental pollutions. This toxic material is absorbed via inhalation, contact, or digestion and affects different tissues.

Objectives: This research was a histomorphometric and immunohistochemical study of the fetal liver of mice after exposure to Diazinon.

Materials and Methods: Twenty-five pregnant BALB/c mice (25-30 gr) were divided into five equal groups in the animal lab of Baqiyatallah University of Medical Sciences, Tehran, Iran. The normal group was without any intervention, and two sham groups received an emulsifier as 0.52 and 5.2 $\mu\text{L}/\text{volume}$ (5000 cc in desiccator) and two experimental groups received Diazinon 1.3 and 13 $\mu\text{L}/\text{volume}$ from the seventh to eighteenth days of pregnancy every other day via forty minutes of inhalation. The pregnant mice were killed on the eighteenth day of gestation and their fetuses were removed and evaluated for fetal growth and liver development. Five fixed fetuses were dehydrated through a series of graded ethanol, embedded in paraffin wax and their whole bodies were sectioned sagittally and stained via the hematoxylin-eosin method. Quantitative computer-assisted morphometric studies were done on the fetal liver tissues occupied by hepatocytes, blood islands, liver sinusoids, and apoptosis.

Results: The mean crown-rump of the fetuses and their mean weight were increased in the experimental group as compared to the sham and normal groups, but the differences were not significant. The mean percentage of the hepatocyte area significantly increased in the experimental group as compared to the sham and control groups ($P < 0.0001$). However, the mean sinusoid area significantly decreased in the experimental group as compared to the sham and control groups. The mean percentage of the area occupied by apoptotic hepatocytes in the experimental group -13 $\mu\text{L}/\text{volume}$ (8.6143 ± 1.00945) and 1.3 $\mu\text{L}/\text{volume}$ (6.1091 ± 0.93093) - significantly increased as compared to the normal and sham groups ($P < 0.0001$).

Conclusions: Our data showed that inhalation of Diazinon during pregnancy increased the hepatocyte area and hepatocyte apoptosis while it decreased the sinusoid area of the fetal liver.

Keywords: Diazinon, Fetal Liver, Apoptosis, Histomorphology

1. Background

Organophosphate (OP) pesticides have been widely used in agriculture as a broad-spectrum pesticide based on its action as an inhibitor of acetylcholine esterase. Several studies have indicated widespread exposure to OP pesticides among some susceptible populations, including pregnant women and children (1, 2). Diazinon is one of the most effective organophosphorus insecticides to control insects and agricultural products (3).

Diazinon is a persistent and dangerous chemical compound. Organophosphates pesticides are commonly used as insecticides and they are generally the most toxic pesticides for the animal species, especially vertebrate animals (4).

Some studies have reported increased exposure to pesticides on women and children and suggested an association between environmental exposure to certain agricultural pesticides, like organophosphorus (OP) compounds, and adverse reproductive outcomes in men and women working on or living near farms (5). Studies reported an association between pesticide exposure and changes in histomorphological parameters and the spermatogenesis of rat testes (6).

The toxicity of OP combinations has been shown in different organs, including the safety and genitourinary systems. It also causes biochemical changes in the blood (7).

During the past few years, a good number of epidemiological studies have been conducted to show the effects of exposure to OPs and other compounds on pregnant

women, but there is a paucity of literature on animal models concerning the effects of OPs during different stages of pregnancy (8). Peiris-John et al. suggested that high OP pesticide level might adversely affect the duration of gestation (5). Some investigators also revealed evidence of the impairment of fetal growth and development brought about by prenatal exposure to OPs. A similar effect was noted with Parathion (9, 10). The deleterious effects due to pre- or post-conception exposure to OP pesticides include menstrual disorders, sterility, fetal toxicity, abortion, stillbirth, and developmental defects (11, 12).

Eskenazi reported that in utero exposure to OP pesticides may decrease fetal growth, shortening the gestational period (13), and Czeizel also demonstrated an increase in congenital malformations after exposure to OP (14). Kalender reported that Diazinon exposure increased levels of aspartate aminotransferase (AST), alanine aminotransferase (ALT), and alkaline phosphatase (ALP). In addition, exposure of hepatocytes to Diazinon accelerates hepatocyte apoptosis and loss of ATP (15).

2. Objectives

Since data concerning the effects of OPs, such as Diazinon, on fetal hepatic tissue have not been shown, this research was carried out to study the histomorphometric and immunohistochemical evaluation of the fetal liver of mice after maternal exposure to Diazinon.

3. Materials and Methods

3.1. Diazinon and Emulsifier Preparations

This OP has 60% Diazinon and 40% emulsifiers as a solvent of Diazinon preparation (Mahan Co. Ltd, Iran).

3.2. Animals and Diazinon Administration

All procedures were approved by the institutional review board of our medical school. The transportation, treatment and experimentation upon the animals was carried out in strict accordance with the guidelines of the animal care and use ethics committee of Baqiyatallah University of Medical Sciences (Tehran, Iran) for experimental study under Code 33 in 2014. The animals were obtained from the animal breeding unit of Baqiyatallah University and the guiding principles for the care and use of laboratory animals were strictly adhered throughout the study.

The mice, weighing 30 ± 5 gr, were maintained on a 12-hour light-dark cycle. The animals were housed at $21 \pm 0.5^\circ\text{C}$ with a relative humidity of $50 \pm 10\%$ and were given ad libitum access to food and water. Males were housed individually with estrous females for approximately 16

hours. Mating was confirmed by the presence of a copulatory plug. The day that the plug was detected was considered day 0 of pregnancy. Twenty-five pregnant mice were separated and then randomly divided into five equal groups (five mice for each group). 1, normal group, without any intervention; 2, sham group, receiving emulsifier as solvent of Diazinon of 0.52 and $5.2 \mu\text{L}/\text{volume}$ (5000 cc in desiccator) and then two experimental groups that received 1.3 and $13 \mu\text{L}/\text{volume}$ of Diazinon respectively (Mahan, Tehran chemical Co. Ltd) for 40 minutes every other day by inhalation from the seventh to the eighteenth day of pregnancy.

3.3. Macroscopic Assessments of Fetuses

All the pregnant mice on Day 18 of gestation were sacrificed via an overdose of chloroform (Merk Co.). The uterine horns were cut and the fetuses and their placentas were removed from the uteruses. The fetuses, randomly selected from four to five litters per group, were examined for congenital malformations using a stereomicroscope. Fetal body weights were assessed using laboratory balance equipment (Sartorius, Japan) and crown-rump lengths were measured using a digital vernier caliper (Germany) and recorded.

3.4. Tissue Preparation and Morphometric Studies

Each fetus was fixed overnight in 10% formalin and processed in paraffin for sectioning. The paraffin blocks of the whole bodies of the fetuses were sagittally sectioned at $5 \mu\text{m}$ thickness and the sections were stained by the hematoxylin and eosin method. The surfaces of the sinusoid and blood island areas were directly measured using a Motic hardware and software system (version 1.2) with a microscopic magnification of $40\times$ of the objective lens. For each section, three measurements were randomly taken from non-adjacent points of the fetal liver. Twenty tissue sections were randomly selected at predetermined intervals for each of the fetuses and histomorphometric studies and cell counting were performed in the following manner - the percentage area that was occupied by hepatocytes per unit area (counting frame $23 \times 20 \mu\text{m}$ equal to $460 \mu\text{m}^2$) was counted with the aid of a rectangular calibrated ocular micrometer using a $100\times$ objective lens. Only hepatocytes that completely fell within the counting frame were counted. Three visual fields were counted for each section. The percentage of the area occupied by hepatocytes, blood islands, and liver sinusoids was measured by Motic software.

3.5. Detection of Apoptotic Hepatocytes by TUNEL Assay

Apoptotic hepatocytes were detected using the technique of terminal-transferase UTP Nick End Labeling

(TUNEL Apoptag plus peroxidase in situ Apoptosis detection kit, S7101, Chemicon). The liver of the fetuses was collected, fixed in 10% formalin, and then processed and embedded in paraffin. Blocks were sectioned at five microns. Sections were mounted on slides and a proteinase k digestion (20 $\mu\text{g}/\text{mL}$) was carried out for 15 minutes. Endogenous hydrogen peroxidase activity was quenched by 3% hydrogen peroxide. The nucleotides contained in the Reaction Buffer were enzymatically added to the DNA by terminal deoxynucleotidyl transferase (TdT). The incubation was carried out for 60 minutes and the labeled DNA was detected using anti-digoxigenin-peroxidase for 30 minutes. The chromogen, diaminobenzidine tetra hydrochloride (DAB), resulted in a brown reaction product that was evaluated by light microscopy and cells were counted. Positive and negative controls were carried out on slides from the same block. Incubation without TdT served as the negative control.

3.6. Statistical Analysis

Data are presented as mean \pm standard deviation (SD). Randomly selected animals ($n = 5$) were used for data analysis. Statistical analyses were performed using SPSS 13 for windows (SPSS Inc., USA). Data were tested for normal distribution using the Kolmogorov-Smirnov test. For data with normal distribution, one-way ANOVA by Tukey's test was used to compare the groups. The samples were randomly allocated to the study group by a randomized block procedure of size $n = 5$. $P < 0.05$ are usually reported as statistically significant.

4. Results

4.1. Fetal Body Weights, Crown-to-Rump Lengths

No significant differences in the indicators of embryo toxicity were found between the sham and normal groups. The mean crown-to-rump lengths of the fetuses in the experimental groups were not significantly different as compared to the normal and sham groups ($P < 0.924$). Likewise, the mean body weight of the fetuses increased in the experimental groups as compared to the normal and sham groups. However, the difference was not statistically significant ($P < 0.803$) (Table 1).

4.2. Morphometric Findings

The mean percent-area occupied by sinusoids in the experimental groups [doses 1.3, (20.85 \pm 8.72) and 13, (16.86 \pm 7.55)] significantly decreased when compared with the normal (37.00 \pm 16.79) and sham groups [doses 0.52, (35.38 \pm 7.53) and 5.2, (36.12 \pm 4.40)] ($P < 0.0001$). On the other hand, the mean percent-area occupied by blood islands

increased in the experimental groups [doses 1.3, (25.47 \pm 16.19) and 13(27.05 \pm 13.02)] as compared to the normal (22.80 \pm 19.95) and sham groups [doses 0.52, (25.36 \pm 8.32) and 5.2, (22.77 \pm 5.88)]. However, the difference was not statistically significant (Figure 1).

The mean percent-area occupied by hepatocytes in the experimental groups at doses of 1.3 (53.69 \pm 17.12) and 13 (56.09 \pm 14.90) $\mu\text{L}/\text{volume}$ of Diazinon significantly increased when compared to the normal (40.20 \pm 4.76) and sham groups [doses 0.52, (39.26 \pm 3.66) and 5.2, (41.11 \pm 1.15)] ($P < 0.0001$) (Figures 1 and 2).

Table 1. Fetal Crown-Rump Lengths (Crl) (Mm) and Fetal Weights (G), Are Shown in Different Groups

Groups	Crown-Rump Lengths Fetuses (CRL)	Weight of Fetuses
Normal	20.76 \pm 4.059	1.040 \pm 0.458
Sham 0.52	20.602 \pm 0.899	1.040 \pm 0.105
Sham 5.2	20.65 \pm 1.311	1.045 \pm 0.080
Exp 1.3	20.49 \pm 0.937	1.085 \pm 0.075
Exp 13	20.97 \pm 1.908	1.092 \pm 0.177
P-Value (between groups)	0.924	0.803

4.3. Apoptotic Hepatocyte Results

The mean number of apoptotic hepatocytes in the experimental groups at doses of 1.3(6.109 \pm 3.087) and 13(8.614 \pm 3.777) $\mu\text{L}/\text{volume}$ of Diazinon significantly increased when compared to the normal (2.970 \pm 2.152) and sham groups [doses 0.52, (3.100 \pm 2.612) and 5.2 (3.300 \pm 3.103)] ($P < 0.0001$) (Figures 3 and 4).

5. Discussion

Diazinon is a non-systemic insecticide used in agriculture to control soil and foliage insects and pests on a variety of fruit, vegetables, nuts and field crops (16). In addition, Diazinon is one of the most applicable organophosphorus insecticides to control insects and agricultural products (3). Reproductive abnormalities caused by organophosphates (OP) have been observed in many animals (17-19). Jorsaraei showed that Diazinon had a significant effect on the structure of rat testes (6). Diazinon induced lipid peroxidation and increased oxygen free radicals in the reproductive tissues in both genders, so the gonads in male and female rats were vulnerable to oxidative stress and its damages, including infertility (20). Sargazi showed that, considering the effect of oxidative stress in multiple physiological processes, from oocyte maturation





Figure 1. The Mean Percent-Area Occupied by Hepatocytes, Sinusoids, and Blood Islands in 18 Day Fetuses in Different Groups

Figure 2. Photomicrographs of the Livers of 18 Day Fetuses



A, Normal; B, Exp 1.3 (at dose of 1.3 μ L/volume of Diazinon); C, Exp 13 (at dose of 13 μ L/volume of Diazinon); H, hepatocyte; S, sinusoids; BI, blood islands) (H and E: 400 \times)

to fertilization, embryo development, and the pathophysiology of infertility, it can be concluded that female rats are more vulnerable to oxidative stress and its consequences, including infertility. Therefore, it is necessary to prevent further toxin entry into the body which causes gonadal dysfunction (20).

Our results showed a non-significant increase in fetal weight after maternal exposure to Diazinon. Our findings are in agreement with previously published research examining prenatal developmental toxicity in pregnant rats which were fed Diazinon doses up to 100 mg/kg/day during days six through fifteen of gestation which showed




Figure 3. The Mean Number of Apoptotic Hepatocytes in 18 Day Fetuses in Different

fetal weights increasing at the highest dose. However, the number of live fetuses decreased and pre- and post-implantation losses were increased at the highest dose (21). In contrast to our results, some researchers have shown that if rat embryos are exposed to Diazinon in culture, the neonatal weights of the experimental group significantly decreased as compared to the control group (22). It has been shown that the teratogenic effects of a drug or chemical agents directly depend on the amount or timing of drug concentration in maternal plasma, but the most important external factor concerning the impact of the drug on the fetus depends on the toxin concentration and duration of the fetal body (23, 24). In addition, our results demonstrated that Diazinon decreased sinusoidal area while it increased hepatocyte and blood island surface area in the fetal liver. On the other hand, our findings also revealed that the mean number of apoptotic hepatocytes in the experimental groups significantly increased when compared to the normal and sham groups.

Some other studies have shown the evacuation of liver tissues, cell infiltration, irritation, vein enlargement, glomeruli degeneration, glomeruli removal, and compression by Diazinon in the treated group (25). Their results are in agreement with our findings about fetal liver tissue. Since the liver is one of the most complex and vital organs and its primary function is detoxification of absorbed substances from the digestive system before their distribution into the systemic circulatory system (26), Diazinon's in vitro and in vivo degradation rate of metabolites in the liver requires further research (27). The exposure to OPCs during pregnancy is an important factor because it affects two organisms - the mother and the fetus (28-30). Moreover, it has been shown that Diazinon can also act as a sub-

strata for glutathione peroxidase (GST) and GPx enzymes that neutralize poisons (16, 31-33).

In addition, antioxidants are scavengers that detoxify excessive ROS and have an important role in maintaining oxidant/antioxidant balance in the body (34).

On the other hand, Yilmaz reported that oxidative stress contributes to Diazinon-induced brain toxicity and the vitamins E and C in combination may have a protective effect against this toxicity (35). Khan et al. also showed that consumption of cypermethrin and malathion by desert mice decreases liver glutathione (36). Fujita and colleagues (1993) reported that the need for oxygen during hypothermia due to high liver metabolism makes the liver hepatocytes susceptible to the negative effects of hypoxia and anoxia (37). It has been shown that OP combinations cause toxicity in different organs, including the safety and genitourinary systems. It also causes biochemical changes in the blood (7). Since most of the OP compounds in the body are converted to the form of an active metabolite through the hepatic cytochrome P450 system and by oxidative desulfurization (38) and this combination is rapidly absorbed over a few hours through the intestine and converted to metabolized diazoxon in liver (39), it seems the fetal liver in our study was affected by Diazinon and suffered apoptotic changes.


Many of these effects have no relation to the control of the acetyl cholinesterase enzyme, but they are induced by other cell mechanisms (40, 41). One of these mechanisms is the production of free radicals and the disorder of the antioxidant systems. Naturally, there is a balance between the production and removal of free radicals. An imbalance of these processes causes oxidative stress. If this stress continues, it can produce serious cell damage (42). One strength of our study was that exposure of Diazinon to pregnant mice occurred via inhalation. Histomorphometric evaluations and immunohistochemical studies were also some of the advantages of our research. In contrast, one of the limitations of our study was that we could not measure how much of Diazinon was absorbed into the bodies of the pregnant mice. Nor could monitor the function of the liver in newborn mice.

In conclusion, the data suggest that prenatal Diazinon exposure has toxic effects on fetal development, as well as increasing apoptosis in fetal hepatocytes.

Acknowledgments

We thank Baqiyatallah University of Medical Sciences and the faculty of biology, Islamic Azad University, north Tehran for their support.

Figure 4. Photomicrographs



A, Normal; B, Exp 1.3 (at dose of 1.3 $\mu\text{L}/\text{volume}$ of Diazinon); C, Exp 13 (at dose of 13 $\mu\text{L}/\text{volume}$ of Diazinon); H, hepatocyte; AP, apoptosis (Tunel: 1000 \times).

Footnotes

Authors' Contribution: Designing the method of study; collection, validation, and analysis of the data; drafting the manuscript, and final revision, Fatemeh Saraei; validation and analysis of the data and final revision, Mehrangiz Sadoughi, Gholamreza Kaka, Seyed Homayoon Sadraie and Mohsen Foadoddin.

Conflicts of Interest: The authors have no conflicts of interest.

Funding/Support: This research project was funded by the center for neuroscience research, Baqiyatallah University of Medical Sciences (Code: 33).

References

- Whyatt RM, Rauh V, Barr DB, Camann DE, Andrews HF, Gar nkel R, et al. Prenatal insecticide exposures and birth weight and length among an urban minority cohort. *Environ Health Perspect.* 2004;**112**(10):1125–32. [PubMed: 15238288].
- Ye X, Pierik FH, Angerer J, Meltzer HM, Jaddoe VW, Tiemeier H, et al. Levels of metabolites of organophosphate pesticides, phthalates, and bisphenol A in pooled urine specimens from pregnant women participating in the Norwegian Mother and Child Cohort Study (MoBa). *Int J Hyg Environ Health.* 2009;**212**(5):481–91. doi: 10.1016/j.ijheh.2009.03.004. [PubMed: 19394271].
- Hariri AT, Moallem SA, Mahmoudi M, Memar B, Hosseinzadeh H. Sub-acute effects of diazinon on biochemical indices and specific biomarkers in rats: protective effects of crocin and safranal. *Food Chem Toxicol.* 2010;**48**(10):2803–8. doi: 10.1016/j.fct.2010.07.010. [PubMed: 20637253].
- Shah MD, Iqbal M. Diazinon-induced oxidative stress and renal dysfunction in rats. *Food Chem Toxicol.* 2010;**48**(12):3345–53. doi: 10.1016/j.fct.2010.09.003. [PubMed: 20828599].
- Peiris-John RJ, Wickremasinghe R. Impact of low-level exposure to organophosphates on human reproduction and survival. *Trans R Soc Trop Med Hyg.* 2008;**102**(3):239–45. doi: 10.1016/j.trstmh.2007.11.012. [PubMed: 18242652].
- Jorsaraei SGA, Firoozjaee A, Pasha YY, Marzony ET, Sarabi E. Histopathological effects of single dose treatment of diazinon on testes structure in rat. *Yakhteh Med J.* 2010;**12**(1):39–42.
- Chambers J, Oppenheimer SF. Organophosphates, serine esterase inhibition, and modeling of organophosphate toxicity. *Toxicol Sci.* 2004;**77**(2):185–7. [PubMed: 14992203].
- Eskenazi B, Bradman A, Castorina R. Exposures of children to organophosphate pesticides and their potential adverse health effects. *Environ Health Perspect.* 1999;**107** Suppl 3:409–19. [PubMed: 10346990].
- Weitman SD, Vodcink MJ, Lech JJ. Influence of pregnancy on parathion toxicity and disposition. *Toxicol Appl Pharmacol.* 1983;**71**(2):215–24. [PubMed: 6636186].
- Weitman SD, Vodcink MJ, Lech JJ. Mechanism of enhanced parathion/paraoxon toxicity during pregnancy in the mouse. *Toxicological Sciences.* 1986;**6**(1):155–61.
- Sever LE, Arbuckle TE, Sweeney A. Reproductive and developmental effects of occupational pesticide exposure: the epidemiologic evidence. *Occup Med.* 1997;**12**(2):305–25. [PubMed: 9220488].
- Arbuckle TE, Sever LE. Pesticide exposures and fetal death: a review of the epidemiologic literature. *Crit Rev Toxicol.* 1998;**28**(3):229–70. doi: 10.1080/10408449891344218. [PubMed: 9631282].
- Eskenazi B, Harley K, Bradman A, Weltzien E, Jewell NP, Barr DB, et al. Association of in utero organophosphate pesticide exposure and fetal growth and length of gestation in an agricultural population. *Environ Health Perspect.* 2004;**112**(10):1116–24. [PubMed: 15238287].
- Czeizel AE, Elek C, Gundy S, Metneki J, Nemes E, Reis A, et al. Environmental trichlorfon and cluster of congenital abnormalities. *Lancet.* 1993;**341**(8844):539–42. [PubMed: 8094783].
- Kalender S, Ogutcu A, Uzunhisarcikli M, Acikgoz F, Durak D, Ulusoy Y, et al. Diazinon-induced hepatotoxicity and protective effect of vitamin E on some biochemical indices and ultrastructural changes. *Toxicology.* 2005;**211**(3):197–206. doi: 10.1016/j.tox.2005.03.007. [PubMed: 15925023].
- Flynt E, Dupuy AJ, Kennedy C, Bennett S. Solid-phase microextraction of organophosphate pesticides in source waters for drinking water treatment facilities. *J Chromatogr Sci.* 2006;**44**(8):484–8. [PubMed: 16959124].
- Dutta H, Meijer H. Sublethal effects of diazinon on the structure of the testis of bluegill, *Lepomis macrochirus*: a microscopic analysis. *Environ Pollut.* 2003;**125**(3):355–60.
- Contreras HR, Bustos-Obregon E. Morphological alterations in mouse testis by a single dose of malathion. *J Exp Zool.* 1999;**284**(3):355–9. [PubMed: 10404127].
- Sobarzo C, Bustos-Obregon E. Sperm quality in mice acutely treated with parathion. *Asian J Androl.* 2000;**2**(2):147–50. [PubMed: 11232794].

20. Sargazi Z, Nikravesh M, Jalali M, Sadeghnia H, Anbarkeh FR, Mohammadzadeh L. Gender-related differences in sensitivity to diazinon in gonads of adult rats and the protective effect of vitamin E. *IJWHR*. 2015;**3**(1):40-7.
21. Wright CG, Leidy RB, Dupree HJ. Chlorpyrifos in the air and soil of houses eight years after its application for termite control. *Bull Environ Contam Toxicol*. 1994;**52**(1):131-4. [PubMed: 7510550].
22. Trimmer GW, Freeman JJ, Priston RA, Urbanus J. Results of chronic dietary toxicity studies of high viscosity (P70H and P100H) white mineral oils in Fischer 344 rats. *Toxicol Pathol*. 2004;**32**(4):439-47. doi: 10.1080/01926230490465865. [PubMed: 15204967].
23. Schreinemachers DM, Everson RB. Aspirin use and lung, colon, and breast cancer incidence in a prospective study. *Epidemiology*. 1994;**5**(2):138-46. [PubMed: 8172988].
24. Smith WL, Lands WE. Stimulation and blockade of prostaglandin biosynthesis. *J Biol Chem*. 1971;**246**(21):6700-2. [PubMed: 5132680].
25. Cakici O, Akat E. Effects of oral exposure to diazinon on mice liver and kidney tissues: biometric analyses of histopathologic changes. *Anal Quant Cytopathol Histopathol*. 2013;**35**(1):7-16. [PubMed: 23469619].
26. Erbas O, Akseki HS, Solmaz V, Aktug H, Taskiran D. Fatty liver-induced changes in stereotypic behavior in rats and effects of glucagon-like peptide-1 analog on stereotypy. *Kaohsiung J Med Sci*. 2014;**30**(9):447-52. doi: 10.1016/j.kjms.2014.05.007. [PubMed: 25224767].
27. Larkin DJ, Tjeerdema RS. Fate and effects of diazinon. *Rev Environ Contam Toxicol*. 2000;**166**:49-82. [PubMed: 10868076].
28. Abu-Qare AW, Abdel-Rahman A, Brownie C, Kishk AM, Abou-Donia MB. Inhibition of cholinesterase enzymes following a single dermal dose of chlorpyrifos and methyl parathion, alone and in combination, in pregnant rats. *J Toxicol Environ Health A*. 2001;**63**(3):173-89. doi: 10.1080/15287390151101529. [PubMed: 11405414].
29. Abu-Qare AW, Abdel-Rahman AA, Kishk AM, Abou-Donia MB. Placental transfer and pharmacokinetics of a single dermal dose of [¹⁴C]methyl parathion in rats. *Toxicol Sci*. 2000;**53**(1):5-12. [PubMed: 10653515].
30. Abu-Qare AW, Abou-Donia MB. Inhibition and recovery of maternal and fetal cholinesterase enzyme activity following a single cutaneous dose of methyl parathion and diazinon, alone and in combination, in pregnant rats. *J Appl Toxicol*. 2001;**21**(4):307-16. [PubMed: 11481665].
31. Dringen R. Metabolism and functions of glutathione in brain. *Prog Neurobiol*. 2000;**62**(6):649-71. [PubMed: 10880854].
32. Amirkabirian N, Teimouri F, Esmaily H, Mohammadirad A, Aliahmadi A, Abdollahi M. Protection by pentoxifylline of diazinon-induced toxic stress in rat liver and muscle. *Toxicol Mech Methods*. 2007;**17**(4):215-21. doi: 10.1080/15376510600943783. [PubMed: 20020971].
33. Winkler BS, Orselli SM, Rex TS. The redox couple between glutathione and ascorbic acid: a chemical and physiological perspective. *Free Radic Biol Med*. 1994;**17**(4):333-49. [PubMed: 8001837].
34. Agarwal A, Aponte-Mellado A, Premkumar BJ, Shaman A, Gupta S. The effects of oxidative stress on female reproduction: a review. *Reprod Biol Endocrinol*. 2012;**10**:49. doi: 10.1186/1477-7827-10-49. [PubMed: 22748101].
35. Yilmaz N, Yilmaz M, Altuntas I. Diazinon-induced brain toxicity and protection by vitamins E plus C. *Toxicol Ind Health*. 2012;**28**(1):51-7. doi: 10.1177/0748233711404035. [PubMed: 21543467].
36. Khan SM, Sobti RC, Kataria L. Pesticide-induced alteration in mice hepato-oxidative status and protective effects of black tea extract. *Clin Chim Acta*. 2005;**358**(1-2):131-8. doi: 10.1016/j.cccn.2005.02.015. [PubMed: 15885683].
37. Fujita S, Hamamoto I, Nakamura K, Tanaka K, Ozawa K. Evaluation of oxygen necessity during hypothermic liver perfusion. *Nihon Geka Hokan*. 1993;**62**(5):228-40. [PubMed: 8031217].
38. Worek F, Diepold C, Eyer P. Dimethylphosphoryl-inhibited human cholinesterases: inhibition, reactivation, and aging kinetics. *Arch Toxicol*. 1999;**73**(1):7-14. [PubMed: 10207609].
39. Garrett SJ, Jones K, Mason HJ, Cocker J. Exposure to the organophosphate diazinon: data from a human volunteer study with oral and dermal doses. *Toxicology letters*. 2002;**134**(1):105-13.
40. Stallones L, Beseler C. Pesticide illness, farm practices, and neurological symptoms among farm residents in Colorado. *Environ Res*. 2002;**90**(2):89-97. [PubMed: 12483798].
41. Storm JE, Rozman KK, Doull J. Occupational exposure limits for 30 organophosphate pesticides based on inhibition of red blood cell acetylcholinesterase. *Toxicology*. 2000;**150**(1-3):1-29. [PubMed: 10996660].
42. Michiels C, Raes M, Toussaint O, Remacle J. Importance of S-glutathione peroxidase, catalase, and Cu/Zn-SOD for cell survival against oxidative stress. *Free Radic Biol Med*. 1994;**17**(3):235-48. [PubMed: 7982629].