

Association of Body Mass Index with Dyslipidemia among the Government Staff of Kermanshah, Iran: A Cross-Sectional Study

Ebrahim Shakiba,¹ Nahid Khademi,² Malihe Khoramdad,^{3,*} Yousef Alimohamadi,^{4,5} and Neda Izadi⁶

¹MD-PhD, Faculty of Health, Kermanshah University of Medical Sciences, Kermanshah, Iran

²MD, Faculty of Health, Kermanshah University of Medical Sciences, Kermanshah, Iran

³MSc of Epidemiology, Faculty of Health, Kermanshah University of Medical Sciences, Kermanshah, Iran

⁴Pars Advanced And Minimally Invasive Medical Manners Research Center, Pars Hospital, Iran University of Medical Sciences, Tehran, Iran

⁵Noor Research Center for Ophthalmic Epidemiology, Noor Eye Hospital, Tehran, Iran

⁶PhD student of Epidemiology, Department of Epidemiology, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

*Corresponding author: Malihe Khoramdad, Faculty of Health, Kermanshah University of Medical Sciences, Kermanshah, Iran. E-mail: m.khoramdad@yahoo.com

Received 2016 December 03; Revised 2017 March 30; Accepted 2017 May 07.

Abstract

Background: There is some evidence suggesting that Body Mass Index (BMI) may increase plasma lipoprotein levels.

Objectives: This study aimed to evaluate the possible association between BMI and dyslipidemia among the government staff of Kermanshah.

Methods: This descriptive-analytical cross-sectional study was performed on the staff of 13 governmental organizations in Kermanshah (a city located in the west of Iran) in 2012. The staff was recruited using census method. We obtained information of 1496 staff aged 22 - 69 through the standardized stepwise questionnaire for NCDs risk factor surveillance. The staff was then physically examined by trained persons. Finally, their blood samples were obtained.

Results: The prevalence of dyslipidemia was 16.6% in the study group, with the prevalence of 18.2% in men and 7.9% in women. There were also 53% (55.8% in men and 7.9% in women) and 17.8% (17.6% in men and 38.1% in women) prevalence rates for overweight and obesity, respectively. Dyslipidemia was significantly associated with factors such as BMI (P value = 0.004), gender (P value < 0.001), marital status (P value = 0.01), cigarette smoking (P value = 0.008), and hookah smoking (P value = 0.002), but not with passive smoking, age, education level, physical activity, FBS (fast blood sugar), WHR (waist to hip ratio), hypertension, waist size, and hip size. In the adjusted model, there was a link merely between obesity and dyslipidemia while no statistically significant association was found between dyslipidemia and overweight. Moreover, a non-linear dose-response association was observed between dyslipidemia and BMI.

Conclusions: Since dyslipidemia was significantly prevalent among men, fat people, and smokers, a particular attention is crucial to be paid to these groups.

Keywords: Dyslipidemia, Body Mass Index, Staff, Kermanshah, Iran

1. Background

Cardiovascular diseases have been accounted for a remarkable rate of mortality and disability especially in societies with low/moderate economic status. Although developed countries have successfully controlled risk factors related to cardiovascular diseases (1), nowadays, hyperlipidemia is a key contributing factor to these disorders (2). Through preventing hyperlipidemia, we will be able to reduce the mortality caused by cardiovascular diseases. Notably, a reduction of 2% - 3% in coronary diseases' risk has been reported following a 1% decrease in serum levels of lipids (3). However, the reduced serum lipid mainly depends on the identification of etiological factors, like obesity (4, 5). One of the frequently used tools to measure obesity is body mass index (6). This factor may result in elevated (not necessarily in a linear mode) levels of serum lipids (7). A remarkable relationship has been re-

ported between BMI and hyperlipidemia in various studies. Nonetheless, when conducting multivariate analysis, BMI in these studies had been converted into a categorical variable (8).

In this study, we attempted to investigate the dose-response link between the BMI continuous variance and the frequency of hyperlipidemia among the government staff of Kermanshah.

2. Methods

This study was a part of a project entitled "investigation of risk factors related to uncommunicable diseases in staff of Kermanshah, Iran". This descriptive-analytical cross-sectional study was performed on the staff of 13 governmental organizations in Kermanshah (a city located in the west of Iran) in 2012. The staff was recruited using census method. We obtained information of 1496 staff aged

between 22 and 69 years. All individuals who previously filled an informed consent form voluntarily participated in this study. In addition, this study gave an ethical code numbered KUMS.ARBC.1394.245. The details of study process have been published elsewhere (9).

2.1. Data Collection and Measurements

The standardized stepwise questionnaire for NCDs risk factor surveillance, suggested by the world health organization, was used. The validated questionnaire was utilized by Meysamie et al. in a national study (9). In addition, reliability of the questionnaire was determined as 0.85 using Cronbach's alpha. The subjects were weighted on a calibrated scale while wearing light clothes and no shoes. The height of the subjects was measured using a tape measure while they were standing on a smooth and flat surface. Subsequently, the measurement of waist size in the narrowest part and hip size in the widest part was conducted using a tape measure, while wearing minimum clothes. In addition, blood pressure of all participants was measured after 5 min rest using a mercury pressure-meter mounted on their right hand. Venous blood samples of fasting subjects were collected to measure fast blood sugar (FBS), as well as blood lipid profile including total cholesterol (TC), high-density cholesterol (HDL), low-density cholesterol (LDL), and triglycerides (TG).

2.2. Definitions

Dyslipidemia: An abnormal body condition in which the amount of total cholesterol is ≥ 240 mg/dL and/or triglycerides ≥ 200 and/or low-density lipoprotein ≥ 140 , and/or high-density lipoprotein cholesterol ≤ 35 (10). BMI: An index in which the individual's weight in kilograms is divided by the square of height (per meters). Our study described normal weight with $18.5 \leq \text{BMI} \leq 24.9$ kg/m², overweight with $25 \leq \text{BMI} \leq 29.9$ kg/m², and obese with $\text{BMI} \geq 30$ (11). Abdominal obesity (WHR) was defined as the waist circumference (cm) divided by hip circumference based on cm. The WHR of < 0.95 in men and < 0.80 in women was considered normal (12). People who had a legal spouse were considered as married, while those who were not married, got divorced, or whose spouse had died were considered as singles. Active smoker: a person who consumes at least one cigarette a day. Hookah smoker: a person with hookah consumption, at least once a day. Passive smoker: non-smoker persons who are exposed to the smoke of cigarettes, hookah, and pipe at their office or their house at least 5 minutes a day (13). Physical activity was determine according to the frequency and the amount of body activity at the time of working and break (metabolic scale/min) expressed as an adjusted standard

unit. Finally, a physical activity less than 600 MET- minutes/week was regarded as low level. High blood pressure: Persons who mentioned a history of high blood pressure or had a recent blood pressure. An FBS of ≥ 126 was considered abnormal (14, 15).

2.3. Statistical Analysis

We used the STATA and SPSS to analyze our data. Distribution of demographic characteristics among participants with/without hyperlipidemia was calculated by Chi-square test, evaluating categorical variables, and logistic regression, evaluating quantitative variables. We used the adjusted logistic regression model to check the correlation of BMI with dyslipidemia. Furthermore, restricted cubic spline method was utilized to evaluate potential non-linear relationship between BMI and hyperlipidemia. Variables with a P value less than 0.05 entered into the multi-variable regression. The level of significance of all 2-tail statistical tests was considered < 0.05 .

3. Results

The prevalence of dyslipidemia was 16.6% (18.2% in men and 7.9% in women). There was also a BMI-based prevalence of 53% (55.8% in men and 38.1% in women) and 17.8% (17.6% in men and 19.2% in women) for overweight and obesity, respectively. Tables 1 and 2 illustrate demographic features of staff working in Kermanshah province. These tables show a statistically significant association between dyslipidemia and gender, cigarette smoking, hookah smoking, marital status, and BMI, while such association was not found for age, education level, FBS, waist size, hip size, WHR, hypertension, physical activity, and passive smoking.

Table 3 indicates the relationship of dyslipidemia with BMI in crude and adjusted models. With regard to this table, in both models, there was a link between obesity and dyslipidemia, as in the crude model the rate of dyslipidemia in fat people was two-times higher than that of normal cases. In model I and II, the possibility of dyslipidemia occurrence was 0.93 higher compared to normal persons.

The dose-response relationship between BMI and dyslipidemia is shown in Figure 1. This figure implies a significant nonlinear dose-response association between BMI and increased risk of dyslipidemia (P value: 0.023), regarding the reference, the ORs were 1.01 (0.86 - 1.17) for BMI at 25.39 kg/m², 1.05 (0.67 - 1.65) for BMI at 28.04 kg/m², and 0.93 (2.37 - 2.79) for BMI at 33.05 kg/m² (Figure 1).

Table 1. Demographic Features and Their Association with Dyslipidemia Among Government Staff of Kermanshah in 2012 (n = 1496)^a

Variable	N	Non-Dyslipidemia (n = 1248)	Dyslipidemia (n = 248)	χ^2	P Value ^b
Gender				15.31	< 0.001
Male	1257	1028 (82.4)	229 (92.3)		
Female	239	220 (17.6)	19 (7.7)		
Educational level				1.12	0.89
Under diploma	231	192 (15.4)	39 (15.7)		
Diploma	347	287 (23)	60 (24.2)		
Associated degree	229	194 (15.5)	35 (14.1)		
BSc	561	465 (37.3)	96 (38.7)		
Msc and above	128	110 (8.8)	18 (7.3)		
Marital status				5.96	0.01
Single	177	159 (12.7)	18 (7.3)		
Married	1319	1089 (87.3)	230 (92.7)		
Smoking				6.93	0.008
No	1372	1155 (92.5)	217 (87.5)		
Yes	124	93 (7.5)	31 (12.5)		
Hookah smoking				9.27	0.002
No	1351	1140 (91.3)	211 (85.1)		
Yes	145	108 (8.7)	37 (14.9)		
Passive smoking				0.03	0.85
No	1316	1097 (87.9)	219 (88.3)		
Yes	180	151 (12.1)	29 (11.7)		
Physical activity				0.44	0.50
600 < MET- minutes/week	671	555 (44.5.9)	116 (46.8)		
600 \geq MET- minutes/week	825	693 (55.5)	132 (53.2)		
Central obesity, (Waist Hip Ratio)				1.32	0.24
No	947	798 (63.9)	149 (60.1)		
Yes	549	450 (36.1)	99 (39.9)		
Hypertension				0.73	0.39
No	1194	1001 (80.2)	193 (77.8)		
Yes	302	247 (19.8)	55 (22.2)		
High FBS				3.69	0.05
No	1433	1201 (96.2)	232 (93.5)		
Yes	63	47 (3.8)	16 (6.5)		

^aValues are expressed as No. (%).^bSignificance level: P value < 0.05.

4. Discussion

With regard to the strong adverse effect of cardiovascular diseases on societies, deep consideration of risk factors is highly recommended. Particularly, abnormalities

of blood lipid profile, as one of the most important risk factors, have been thoroughly investigated (16, 17). In this study, the overall prevalence of hyperlipidemia was reported as 16.6%. Muhammadbeigi et al. estimated this prevalence to be 51.8% (18). Similarly, in a meta-analysis

Table 2. Demographic Features and Their Association with Dyslipidemia Among Government Staff of Kermanshah in 2012 (n = 1496)

Variable	B	S.E.	Wald	P Value	OR (95% CI) ^a
Age, y	0.17	0.008	4.34	0.037	1.01 (1.00 - 1.03)
BMI, kg/m ²	0.05	0.018	8.20	0.004	1.05 (1.01 - 1.09)
Waist, cm	0.006	0.007	0.60	0.40	1.006 (0.99 - 1.02)
Hip, cm	-0.005	0.009	0.35	0.55	0.99 (0.97 - 1.01)

^aThe level of significance was tested by logistic regression.

Table 3. Logistic Regression Analysis for the Influence of BMI on Dyslipidemia among Government Staff of Kermanshah in 2012 (n = 1496)

Model	BMI	B	S.E.	Wald	P Value ^a	OR (95% CI)
Univariate	18.5 to 24.9					1.00
	25 to 29.9	0.29	0.17	2.93	0.08	1.34 (0.95 - 1.88)
	≥ 30	0.69	0.20	11.66	0.001	2.00 (1.34 - 2.99)
Model I ^b	18.5 to 24.9					1.00
	25 to 29.9	0.20	0.17	1.45	0.22	1.23 (0.87 - 1.73)
	≥ 30	0.66	0.20	10.28	0.001	1.93 (1.29 - 2.89)
Model II ^c	18.5 to 24.9					1.00
	25 to 29.9	0.22	0.17	1.66	0.19	1.25 (0.88 - 1.77)
	≥ 30	0.66	0.20	10.08	0.001	1.93 (1.28 - 2.91)

^aSignificance level: P value < 0.05.

^bAdjusted for gender.

^cAdjusted for gender, marital status, smoking, hookah smoking.




Figure 1. The Dose-Response Relationship of BMI with Dyslipidemia Based on the Restricted Cubic Spline Model (BMI Was Coded Using an RCS Function with Four Knots Located at 21.07 kg/m², 25.39 kg/m², 28.04 kg/m², and 33.05 kg/m²).

study, Tabatabaei et al. reported the prevalence of different types of dyslipidemia including hypercholesterolemia, hypertriglyceridemia, and LDL as 41.6%, 46%, and 35%, respectively (19). In addition, the prevalence of hyperlipidemia in China has been reported to be 51.09%. There was also a frequency of 8.68% in rural and 5.74% in urban areas of India (8, 20).

In this study, the prevalence of obesity, according to BMI, was estimated to be 17.8%, being more than that of USA and Europe (21, 22) and less than that of South Africa (23). The reason for these differences can be due to the difference in statistical populations, race, and life style.

Dyslipidemia was found to be more common in men than in women. In contrast, in Germany and United States, women are involved more in the disease (24, 25). Most studies have reported results similar to ours (26, 27). This might be due to the higher smoking rate in men, whose physical activity is lower and overall/abdominal obesity is higher, resulting in the increased levels of blood lipids (28).

We also found a significant relationship between smoking and dyslipidemia, similar to studies conducted by Mouhamed and Brucket (29, 30). No significant association was found between dyslipidemia and exposing to smoke. Conversely, Hirata et al. found a raised level of HDL in cases exposed to smoke (31). Furthermore, a cross-sectional study conducted on children has shown a decreased level of HDL and increased level of LDL (32). Moreover, a case-control study on 164 patients with 30 - 64 years of age showed a significantly decreased level of serum lipids in people exposed to smoke, when compared to non-

smokers (14).

Our results also highlighted the capability of obesity in increasing the rate of dyslipidemia, similar to results reported by Lai et al. (33), Kenneth et al. (34) and Sullivan et al. (35). Similarly, a Chinese study showed a significant nonlinear association between BMI and raised risk of hyperlipidemia, with a significantly increased trend of odds ratio following 1 kg/m² increase in BMI (8).

The relationship between education level and hyperlipidemia has been investigated in some studies. Mahley and Pradeepa showed that people with high education level represent a raised rate of dyslipidemia (36, 37), whereas Wang and Al-Kaabba have proposed a decreased rate of dyslipidemia with high levels of education (38, 39).

In this study, the relationship between hyperlipidemia and elevated levels of blood sugar was not significant, whereas Wang showed a significant correlation between dyslipidemia and high blood sugar (38). Similarly, Luo reported higher levels of HDL in people with high blood sugar (40). Furthermore, we did not observe any statistically significant relationship between dyslipidemia and high blood pressure, while different studies have suggested that lipids metabolism varies (41) (as also proved in prior studies) in people with high blood pressure, compared to normal cases (42). The reason for this diversity may be due to either difference in conditions wherein study conducted or difference in study cases.

Some advantages could be mentioned for this study. First, the blood samples of all staff were collected and second, dose-response analysis was conducted to understand whether there is any association between BMI and dyslipidemia. In contrast, there are some disadvantages: 1) there might be some confounders affecting our results and, 2) we just collected samples of staff, and no sample was taken from the rest of society.

4.1. Conclusions

According to the growing rate of cardiovascular diseases, paying more attention to risk factors is urgently needed to be able to decrease the incidence of these diseases. Hyperlipidemia, in particular, is an important factor that may result in serious problems if ignored. Men, obese, and smoker cases are at the highest risk. Consequently, preventive strategies should be focused mostly on these groups.

Acknowledgments

We would like to appreciate all the staff at the governmental organizations of Kermanshah, and all those researchers who helped us conduct this study.

Footnote

Financial Disclosure: This work was supported by Kermanshah University of Medical Sciences. The funding agency did not play any role in the planning, execution, or reporting process, nor in the decision to submit the paper for publication.

References

1. Yusuf S, Wood D, Ralston J, Reddy KS. The World Heart Federation's vision for worldwide cardiovascular disease prevention. *Lancet*. 2015;**386**(9991):399-402. doi: [10.1016/S0140-6736\(15\)60265-3](https://doi.org/10.1016/S0140-6736(15)60265-3). [PubMed: [25892680](https://pubmed.ncbi.nlm.nih.gov/25892680/)].
2. Nelson RH. Hyperlipidemia as a risk factor for cardiovascular disease. *Prim Care*. 2013;**40**(1):195-211. doi: [10.1016/j.pop.2012.11.003](https://doi.org/10.1016/j.pop.2012.11.003). [PubMed: [23402469](https://pubmed.ncbi.nlm.nih.gov/23402469/)].
3. Marrie RA, Yu BN, Leung S, Elliott L, Caetano P, Warren S, et al. Rising prevalence of vascular comorbidities in multiple sclerosis: validation of administrative definitions for diabetes, hypertension, and hyperlipidemia. *Mult Scler*. 2012;**18**(9):1310-9. doi: [10.1177/1352458512437814](https://doi.org/10.1177/1352458512437814). [PubMed: [22328682](https://pubmed.ncbi.nlm.nih.gov/22328682/)].
4. Chang CJ, Jian DY, Lin MW, Zhao JZ, Ho LT, Juan CC. Evidence in obese children: contribution of hyperlipidemia, obesity-inflammation, and insulin sensitivity. *PLoS One*. 2015;**10**(5):e0125935. doi: [10.1371/journal.pone.0125935](https://doi.org/10.1371/journal.pone.0125935). [PubMed: [26011530](https://pubmed.ncbi.nlm.nih.gov/26011530/)].
5. Canale MP, Manca di Villahermosa S, Martino G, Rovella V, Noce A, De Lorenzo A, et al. Obesity-related metabolic syndrome: mechanisms of sympathetic overactivity. *Int J Endocrinol*. 2013;**2013**:865965. doi: [10.1155/2013/865965](https://doi.org/10.1155/2013/865965). [PubMed: [24288531](https://pubmed.ncbi.nlm.nih.gov/24288531/)].
6. Ogden CL, Carroll MD. Prevalence of overweight, obesity, and extreme obesity among adults: United States, trends 1960-1962 through 2007-2008. *Nat Center Health Statistics*. 2010;**6**(1):1-6.
7. Hirakawa Y, Lam TH, Welborn T, Kim HC, Ho S, Fang X, et al. The impact of body mass index on the associations of lipids with the risk of coronary heart disease in the Asia Pacific region. *Prev Med Rep*. 2016;**3**:79-82. doi: [10.1016/j.pmedr.2015.12.012](https://doi.org/10.1016/j.pmedr.2015.12.012). [PubMed: [26844192](https://pubmed.ncbi.nlm.nih.gov/26844192/)].
8. Rao W, Su Y, Yang G, Ma Y, Liu R, Zhang S, et al. Cross-Sectional Associations between Body Mass Index and Hyperlipidemia among Adults in Northeastern China. *Int J Environ Res Public Health*. 2016;**13**(5) doi: [10.3390/ijerph13050516](https://doi.org/10.3390/ijerph13050516). [PubMed: [27213419](https://pubmed.ncbi.nlm.nih.gov/27213419/)].
9. Khademi N, Babanejad M, Najafi F, Nikbakht MR, Hamzeh B, Mohammadi N. Tobacco Use and its Relationship with Health Complaints Among Employees of Kermanshah Province, Iran. *Int J Prev Med*. 2016;**7**:71. doi: [10.4103/2008-7802.181757](https://doi.org/10.4103/2008-7802.181757). [PubMed: [27226894](https://pubmed.ncbi.nlm.nih.gov/27226894/)].
10. Navaii L, Mehrabi Y, Azizi F. An epidemiologic study of hyperlipidemia, obesity, and hypertension in Tehran villages. *Iran J Endocrinol Metab*. 2000;**2**(4):253-62.
11. Garcia-Alvarez A, Serra-Majem L, Ribas-Barba L, Castell C, Foz M, Uauy R, et al. Obesity and overweight trends in Catalonia, Spain (1992-2003): gender and socio-economic determinants. *Public Health Nutr*. 2007;**10**(11A):1368-78. doi: [10.1017/S1368980007000973](https://doi.org/10.1017/S1368980007000973). [PubMed: [17988408](https://pubmed.ncbi.nlm.nih.gov/17988408/)].
12. Hedayati M. Central obesity as a reliable predictor for hypertension and dyslipidemia: Tehran Lipid Glucose Study. *Iran J Endocrinol Metab*. 2010;**12**(3):251-9.
13. Mohammadifard N, Sarafzadegan N, Nouri N, Bagheri M. The effect of passive smoking on acute myocardial infarction and its relationship with fibrinogen and serum lipids. ; 2004.
14. Biderafsh A, Karami M, Faradmal J, Poorolajal J, Esmailnasab N. The pattern of hypertension and the population attributable proportion of hypertension-related stroke in Hamadan province from 2005 to 2009. *Iran J Epidemiol*. 2014;**10**(3):54-64.

15. Yazdanpanah L, Shahbazian HB, Moravej Aleali A, Jahanshahi A, Ghanbari S, Latifi SM. Prevalence, awareness and risk factors of diabetes in Ahvaz (South West of Iran). *Diabetes Metab Syndr*. 2016;**10**(2 Suppl 1):S114-8. doi: [10.1016/j.dsx.2016.03.007](https://doi.org/10.1016/j.dsx.2016.03.007). [PubMed: 27312373].
16. Carr MC, Brunzell JD. Abdominal obesity and dyslipidemia in the metabolic syndrome: importance of type 2 diabetes and familial combined hyperlipidemia in coronary artery disease risk. *J Clin Endocrinol Metab*. 2004;**89**(6):2601-7. doi: [10.1210/jc.2004-0432](https://doi.org/10.1210/jc.2004-0432). [PubMed: 15181030].
17. Manninen V, Tenkanen L, Koskinen P, Huttunen JK, Manttari M, Heinonen OP, et al. Joint effects of serum triglyceride and LDL cholesterol and HDL cholesterol concentrations on coronary heart disease risk in the Helsinki Heart Study. Implications for treatment. *Circulation*. 1992;**85**(1):37-45. [PubMed: 1728471].
18. Mohammadbeigi A, Moshiri E, Mohammadsalehi N, Ansari H, Ahmadi A. Dyslipidemia Prevalence in Iranian Adult Men: The Impact of Population-Based Screening on the Detection of Undiagnosed Patients. *World J Mens Health*. 2015;**33**(3):167-73. doi: [10.5534/wjmh.2015.33.3.167](https://doi.org/10.5534/wjmh.2015.33.3.167). [PubMed: 26770936].
19. Tabatabaei-Malazy O, Qorbani M, Samavat T, Sharifi F, Larijani B, Fakhrzadeh H. Prevalence of dyslipidemia in iran: a systematic review and meta-analysis study. *Int J Prev Med*. 2014;**5**(4):373-93. [PubMed: 24829725].
20. Raj A, Sivakumar K, Sujatha K. Prevalence of dyslipidemia in South Indian adults: an urban-rural comparison. *Int J Commun Med Public Health*. 2017;**3**(8):2201-10.
21. Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM. Prevalence of overweight and obesity among US children, adolescents, and adults, 1999-2002. *JAMA*. 2004;**291**(23):2847-50. doi: [10.1001/jama.291.23.2847](https://doi.org/10.1001/jama.291.23.2847). [PubMed: 15199035].
22. Berghofer A, Pischon T, Reinhold T, Apovian CM, Sharma AM, Willich SN. Obesity prevalence from a European perspective: a systematic review. *BMC Public Health*. 2008;**8**:200. doi: [10.1186/1471-2458-8-200](https://doi.org/10.1186/1471-2458-8-200). [PubMed: 18533989].
23. Puoane T, Steyn K, Bradshaw D, Laubscher R, Fourie J, Lambert V, et al. Obesity in South Africa: the South African demographic and health survey. *Obes Res*. 2002;**10**(10):1038-48. doi: [10.1038/oby.2002.141](https://doi.org/10.1038/oby.2002.141). [PubMed: 12376585].
24. McDonald M, Hertz RP, Unger AN, Lustik MB. Prevalence, awareness, and management of hypertension, dyslipidemia, and diabetes among United States adults aged 65 and older. *J Gerontol A Biol Sci Med Sci*. 2009;**64**(2):256-63. doi: [10.1093/gerona/gln016](https://doi.org/10.1093/gerona/gln016). [PubMed: 19181717].
25. Scheidt-Nave C, Du Y, Knopf H, Schienkiewitz A, Ziese T, Nowossadeck E, et al. [Prevalence of dyslipidemia among adults in Germany: results of the German Health Interview and Examination Survey for Adults (DEGS 1)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitschutz*. 2013;**56**(5-6):661-7. doi: [10.1007/s00103-013-1670-0](https://doi.org/10.1007/s00103-013-1670-0). [PubMed: 23703484].
26. Erem C, Hacıhasanoglu A, Deger O, Kocak M, Topbas M. Prevalence of dyslipidemia and associated risk factors among Turkish adults: Trabzon lipid study. *Endocrine*. 2008;**34**(1-3):36-51. doi: [10.1007/s12020-008-9100-z](https://doi.org/10.1007/s12020-008-9100-z). [PubMed: 19003544].
27. Lee MH, Kim HC, Ahn SV, Hur NW, Choi DP, Park CG, et al. Prevalence of Dyslipidemia among Korean Adults: Korea National Health and Nutrition Survey 1998-2005. *Diabetes Metab J*. 2012;**36**(1):43-55. doi: [10.4093/dmj.2012.36.1.43](https://doi.org/10.4093/dmj.2012.36.1.43). [PubMed: 22363921].
28. Gastaldelli A, Folli F, Maffei S. Impact of tobacco smoking on lipid metabolism, body weight and cardiometabolic risk. *Curr Pharm Des*. 2010;**16**(23):2526-30. [PubMed: 20550509].
29. Haj Mouhamed D, Ezzaher A, Neffati F, Gaha L, Douki W, Najjar MF. Association between cigarette smoking and dyslipidemia. *Immuno-analyse & Biologie Specialisee*. 2013;**28**(4):195-200. doi: [10.1016/j.immbio.2013.03.004](https://doi.org/10.1016/j.immbio.2013.03.004).
30. Bruckert E, Jacob N, Lamaire L, Truffert J, Percheron F, de Gennes JL. Relationship between smoking status and serum lipids in a hyperlipidemic population and analysis of possible confounding factors. *Clin Chem*. 1992;**38**(9):1698-705. [PubMed: 1526001].
31. Hirata K, Yamano Y, Suzuki H, Miyagawa S, Nakadate T. Passive smoking is associated with lower serum HDL-C levels in school children. *Pediatr Int*. 2010;**52**(2):252-6. doi: [10.1111/j.1442-200X.2009.02957.x](https://doi.org/10.1111/j.1442-200X.2009.02957.x). [PubMed: 19744225].
32. Neufeld EJ, Mietus-Snyder M, Beiser AS, Baker AL, Newburger JW. Passive cigarette smoking and reduced HDL cholesterol levels in children with high-risk lipid profiles. *Circulation*. 1997;**96**(5):1403-7. [PubMed: 9315524].
33. Lai SW, Ng KC, Lin HF, Chen HL. Association between obesity and hyperlipidemia among children. *Yale J Biol Med*. 2001;**74**(4):205-10. [PubMed: 11697478].
34. Thorpe KE, Florence CS, Howard DH, Joski P. The impact of obesity on rising medical spending. *Health Aff (Millwood)*. 2004;**Suppl Web Exclusives**:W4-480-6. doi: [10.1377/hlthaff.w4.480](https://doi.org/10.1377/hlthaff.w4.480). [PubMed: 15496437].
35. Sullivan PW, Ghushchyan VH, Ben-Joseph R. The impact of obesity on diabetes, hyperlipidemia and hypertension in the United States. *Qual Life Res*. 2008;**17**(8):1063-71. doi: [10.1007/s1136-008-9385-7](https://doi.org/10.1007/s1136-008-9385-7). [PubMed: 1877200].
36. Mahley RW, Palaoglu KE, Atak Z, Dawson-Pepin J, Langlois AM, Cheung V, et al. Turkish Heart Study: lipids, lipoproteins, and apolipoproteins. *J Lipid Res*. 1995;**36**(4):839-59. [PubMed: 7616127].
37. Pradeepa R, Deepa R, Rani SS, Premalatha G, Saroja R, Mohan V. Socioeconomic status and dyslipidaemia in a South Indian population: the Chennai Urban Population Study (CUPS 11). *Natl Med J India*. 2003;**16**(2):73-8. [PubMed: 12816185].
38. Wang S, Xu L, Jonas JB, You QS, Wang YX, Yang H. Prevalence and associated factors of dyslipidemia in the adult Chinese population. *PLoS One*. 2011;**6**(3):e17326. doi: [10.1371/journal.pone.0017326](https://doi.org/10.1371/journal.pone.0017326). [PubMed: 21423741].
39. F. Al-Kaabba A, A. Al-Hamdan N, El Tahir A, M. Abdalla A, A. Saeed A, A. Hamza M. Prevalence and Correlates of Dyslipidemia among Adults in Saudi Arabia: Results from a National Survey. *Open J Endocrine Metab Dis*. 2012;**02**(04):89-97. doi: [10.4236/ojemd.2012.24014](https://doi.org/10.4236/ojemd.2012.24014).
40. Luo Y, Li J, Zhang J, Xu Y. Low HDL cholesterol is correlated to the acute ischemic stroke with diabetes mellitus. *Lipids Health Dis*. 2014;**13**:171. doi: [10.1186/1476-511X-13-171](https://doi.org/10.1186/1476-511X-13-171). [PubMed: 25395241].
41. Lithell H. Hypertension and hyperlipidemia. A review. *Am J Hypertens*. 1993;**6**(1 Pt 2):303S-8S. [PubMed: 8297536].
42. Hunt SC, Wu LL, Hopkins PN, Stults BM, Kuida H, Ramirez ME, et al. Apolipoprotein, low density lipoprotein subfraction, and insulin associations with familial combined hyperlipidemia. Study of Utah patients with familial dyslipidemic hypertension. *Arteriosclerosis*. 1989;**9**(3):335-44. [PubMed: 2497719].