



# Seroprevalence of Hepatitis C Virus among Blood Donors in Middle Eastern Countries: A Systematic Review and Meta-Analysis

Hossein Ghaderi-Zefrehi,<sup>1,2</sup> Heidar Sharafi,<sup>1,3</sup> Farzin Sadeghi,<sup>4</sup> Mohammad Gholami-Fesharaki,<sup>5</sup> Alireza Farasat,<sup>6</sup> Fatemeh Jahanpeyma,<sup>7</sup> and Seyed Moayed Alavian<sup>1,3,\*</sup>

<sup>1</sup>Baqiyatallah Research Center for Gastroenterology and Liver Diseases, Baqiyatallah University of Medical Sciences, Tehran, IR Iran

<sup>2</sup>Department of Clinical Biochemistry, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

<sup>3</sup>Middle East Liver Diseases (MELD) Center, Tehran, IR Iran

<sup>4</sup>Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, IR Iran

<sup>5</sup>Department of Biostatistics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, IR Iran

<sup>6</sup>Young Researchers and Elite Club, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, IR Iran

<sup>7</sup>Department of Medical Biotechnology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, IR Iran

\*Corresponding author: Seyed Moayed Alavian, Baqiyatallah Research Center for Gastroenterology and Liver Diseases, Baqiyatallah University of Medical Sciences, Mollasadra Ave, Vanak Sq, P.O. Box 14155-3651, Tehran, IR Iran. E-mail: alavian@thc.ir

Received 2017 July 13; Revised 2017 October 14; Accepted 2017 November 01.

## Abstract

**Context:** Hepatitis C virus (HCV) has been the main cause of post transfusion hepatitis. Long-term hepatitis can ultimately result in cirrhosis and hepatocellular carcinoma. Viral hepatitis, especially HCV infection, is a major health concern in the Middle East. The current study carried out a systematic search concerning HCV seroprevalence among blood donors in Middle Eastern countries.

**Evidence Acquisition:** Articles were identified through searching databases including PubMed, Scopus, and Web of Science. We reviewed systematically all studies reporting HCV seroprevalence among blood donors in Middle Eastern countries.

**Results:** A total of 5662 relevant records were identified by the electronic search, of which a total of 47 studies were identified as eligible papers that were meta-analyzed for the pooled seroprevalence of HCV among blood donors. Overall, HCV seroprevalence among blood donors in Middle Eastern countries was estimated 0.88% (95%CI = 0.79% - 0.97%). The highest pooled HCV seroprevalence rate was related to Egypt (5.76% [95%CI = 3.30% - 8.22%]), Libya (1.56% [95%CI = 0.99% - 2.13%]), and Yemen (1.09% [95%CI = 0.69% - 1.50%]) while the lowest pooled seroprevalence rate was in Iran and Israel with the rates of 0.14% (95%CI = 0.12% - 0.17%) and 0.16% (95%CI = 0.06% - 0.25%), respectively. Results of annual HCV seroprevalence suggest that there is a decrease in seroprevalence rate of HCV over time among blood donors in Egypt, Iran, Oman, Lebanon, Libya, and Saudi Arabia.

**Conclusions:** Our results showed that the trend of HCV seroprevalence over time among Middle Eastern blood donors was decreasing. It suggests that recent safety measures implemented in Middle Eastern countries have been effective.

**Keywords:** Blood safety, Epidemiology, Hepatitis C, Middle East, Systematic review

## 1. Context

First found in 1989, hepatitis C is caused by hepatitis C virus (HCV) (1) from the Flaviviridae family and it is the main reason of post transfusion hepatitis (2). HCV is most commonly transmitted through direct contact with blood as in blood transfusion (3). Screening of donated bloods for hepatitis C has been implemented in blood transfusion services since 1990s in most Middle Eastern countries (4). The post-transfusion hepatitis C has been a main cause of mortality and morbidity in poly-transfused patients such as patients with thalassemia and hemophilia in Middle Eastern countries (5). Viral hepatitis, especially HCV, is a major health concern in the Middle East, which is a geographical region with diverse ethnic groups including Arabs, Turks, Persians, etc. With an over 3.5% prevalence of

HCV infection, Middle East is considered a hotspot for this disease (6, 7). There has been inadequate research on HCV prevalence and prevention methods in the Middle East (8). Moreover, prevalence of HCV infection among blood donors has not been discussed in some countries of this region. Therefore, this paper reviews the conducted studies in order to provide the current estimate of HCV seroprevalence among blood donors in Middle Eastern countries.

## 2. Evidence Acquisition

### 2.1. Data Resources and Search Strategies

Literatures on HCV prevalence among blood donors in Middle Eastern countries, such as Bahrain, Cyprus, Egypt, Iran, Iraq, Israel, Jordan, Kuwait, Lebanon, Libya, Oman,

Palestine, Qatar, Saudi Arabia, Syria, Turkey, UAE, and Yemen, were acquired through searching PubMed, Scopus, and Web of Science in the time span of 1996 to 2016. Our last search was conducted on October 05, 2016. In order to search and include related studies as many as possible, we used the following terms: "Hepatitis C virus", "HCV", "prevalence", "blood donor", and name of the Middle Eastern countries.

## 2.2. Eligibility Criteria

Published studies were regarded qualified for analysis if they met the following criteria: 1- Studies with full text of the paper available in English language; 2- Studies with a sample size more than 100; and 3- Studies that reported anti-HCV antibody (HCV Ab) prevalence among blood donors in the Middle Eastern countries. Conversely, the followings were exclusion criteria: 1- Non-English reports on the seroprevalence of HCV; 2- Studies that reported seroprevalence of HCV among paid blood donors; 3- Studies which all of the participants were first-time blood donors; 4- Studies with overlapped time or place of donation; 5- Studies that used first-generation or second-generation HCV enzyme-linked immunoassay (ELISA) without supplemental or confirmatory assay, and 6- Studies failing to present evident data.

## 2.3. Study Selection and Data Extraction

All articles categorized as potentially relevant were reviewed separately by two of the authors (H. G-Z. and H. S.). In addition, blinding and task separation were applied in selection of articles and data extraction. For each eligible study, the following information was extracted: first author's name, year of publication, study period, study location, the method for the screening of HCV Ab, the method used for confirmation of HCV Ab, sample size, number of positive cases for HCV Ab, and the type of donors. Analysis was conducted according to the preferred reporting items for systematic reviews and meta-analysis (PRISMA) (9).

## 2.4. Data Analysis

In the current meta-analysis, the seroprevalence rate of HCV among blood donors from each country was computed by metan command. Statistical tests of heterogeneity among the studies were carried out using the Q test ( $P < 0.10$  indicates statistically significant heterogeneity) and I-squared statistics. According to the results of heterogeneity test, we used fixed- or random-effect models for determining the prevalence of HCV Ab among blood donors. We also used funnel plot to investigate publication bias. All data analyses were performed using STATA 10.

## 3. Results

### 3.1. Study Screening and Characteristics of the Included Studies

The study selection process is depicted in Figure 1. A total of 5662 studies potentially associated with the seroprevalence of HCV infection among blood donors in the Middle Eastern countries were identified through web search, of which 2,291 duplicates were excluded. After reviewing the abstracts and titles, 3,181 studies were eliminated based on the stated inclusion and exclusion criteria. After full-text screening, a total of 47 records were deemed as eligible papers published between 1995 and 2016. These 47 records were composed of 13 studies from Iran (4, 10-21), 8 studies from Turkey (22-29), 8 studies from Saudi Arabia (30-37), and 6 studies from Egypt (38-43). Two studies were available from each of the following countries: Syria (44, 45), Libya (46, 47), Yemen (48, 49), and Israel (50, 51). Only one study was available from each of the following countries: Lebanon (52), Oman (53), Cyprus (54), and UAE (55). The characteristics of included studies are summarized in Table 1.

### 3.2. Seroprevalence of HCV among Blood Donors in Middle Eastern Countries

Data on HCV seroprevalence as well as other indicators were abstracted from the 47 records as presented in Table 1. Information on HCV seroprevalence among blood donors was available from twelve countries.

As shown in Table 1, using random-effect model, the pooled seroprevalence of HCV among blood donors in the Middle Eastern countries from 1989 to 2014 was estimated 0.88% (95%CI = 0.79% - 0.97%). In addition, we used random-effect model to estimate pooled HCV seroprevalence rate for each of the Middle Eastern countries. The highest pooled HCV seroprevalence rate was related to Egypt (5.76% [95%CI = 3.30% - 8.22%]), Libya (1.56% [95%CI = 0.99% - 2.13%]), and Yemen (1.09% [95%CI = 0.69% - 1.50%]) while the lowest pooled seroprevalence rate was in Iran and Israel with the rates of 0.14% (95%CI = 0.12% - 0.17%) and 0.16% (95%CI = 0.06% - 0.25%), respectively. HCV seroprevalence among blood donors in countries with one study including Lebanon, Oman, Cyprus, and UAE was 0.40% (95%CI = 0.31% - 0.51%), 0.75% (95%CI = 0.64% - 0.88%), 0.45% (95%CI = 0.30% - 0.68%), and 0.11% (95%CI = 0.10% - 0.13%), respectively. Seroprevalence of HCV among blood donors by different countries is depicted in Figure 2.

For evaluating annual HCV seroprevalence rates and their trends, we only considered studies that reported HCV seroprevalence rate among blood donors annually. By exploring the trends of HCV seroprevalence rate in 6 countries, including Egypt (2007 - 2012), Iran (2003 - 2014), Lebanon (1997 - 2003), Libya (2008 - 2013), Oman (1995 -





Figure 1. Flowchart of Article Selection Based on PRISMA Statement

2001), and Saudi Arabia (1998 - 2014), a decline in HCV seroprevalence rate among blood donors over time was observed (Figure 3). Detailed information on trends is shown in Figure 3.

#### 4. Discussion

Blood transfusion and blood donation continue to play significant roles in the transmission of HCV in Middle Eastern countries (56). Several recent studies have provided good evidence on genotype distribution and epidemiology of HCV in the Middle East region (57-59). However, determination of HCV seroprevalence rates among blood donors in the Middle East region could be helpful to develop prevention and control strategies more effectively. The current systematic review was conducted to shed more light on the seroprevalence of HCV among blood donors in the Middle East region; a transcontinental region with diverse ethnic groups including Arabs, Turks,

Persians, Azeris, Kurds, etc. To determine HCV seroprevalence rate among blood donors, the estimated pooled seroprevalence rate was calculated for each of the Middle Eastern countries. The results of our study indicated that HCV seroprevalence rates among blood donors in the Middle East region is 0.88% that is higher than the seroprevalence rates in North-European blood donors (0.01% - 0.02%) (60). In addition, a recently published meta-analysis reported 0.65% HCV seroprevalence rate among blood donors in African countries, which is also lower than the rate found in our study (61). Based on our statistical analysis, there is a dramatic geographical difference between Middle Eastern countries in terms of HCV seroprevalence rate in blood donors. The highest seroprevalence was found in Egypt with the rate of 5.7% and the lowest seroprevalence was found in UAE and Iran with the rates of 0.11% and 0.14%, respectively. Our results are in good agreement with some studies from Egypt, including two investigations with large sample sizes among Egyptian blood donors con-




**Figure 2.** Seroprevalence of HCV among Blood Donors in Middle Eastern Countries

ducted by Eita et al. (62) and Rushdy et al. (63). In addition, based on a newly published systematic review from Egypt, HCV seroprevalence rates among blood donors ranged 5% - 25% (64). It should be noted that data from the aforementioned systematic review has to be treated with caution due to employing ELISA as a detection index and lack of confirmatory RIBA test. Moreover, the low sample size in some of the studies included in the aforementioned systematic review may affect the obtained results. The results of the present study are consistent with a number of reports from Iranian blood donors (16, 65). Interestingly, a comprehensive study by Kafi abad et al. among Iranian blood donors revealed 0.13% HCV seroprevalence rate that is in good agreement with our results (16). According to our results, Libya and Yemen have the second and third highest HCV seroprevalence rates in blood donors in the Middle East region. These findings are broadly consistent with other comprehensive epidemiologic studies on general populations in Libya and Yemen (66, 67). Seroprevalence of HCV in Libya and Yemen may increase in the next few years due to ongoing military conflicts and destruction of health-care infrastructures. The results of the present study demonstrated almost similar HCV seroprevalence rates in Saudi and Syrian blood donors. These

findings are also consistent with the findings of more recent studies among blood donors in Saudi Arabia and Syria (68). However, our results are not supported by Shobokshi et al. who found 1.1% HCV seroprevalence rate among Saudi blood donor (31). Moreover, military conflicts and refugee crisis in Syria may increase HCV seroprevalence rate in the next few years. In the present study, seroprevalence of HCV in blood donors from Turkey and Israel was low (0.27% and 0.16%, respectively) that might be explained by good infection control practices and organized system of blood screening and transfusion. Our results showed a fluctuating trend in annual HCV seroprevalence rate in the Middle East region. With the exception of Turkey, other countries included in this investigation had a decreasing HCV seroprevalence trend.

Recently, with introduction of HCV direct-acting antiviral agents, the treatment of hepatitis C has transformed greatly (69, 70). These new therapies resulting in more than 95% treatment success have made hepatitis C a curable infectious disease (71-73). With availability of efficient treatments for hepatitis C, there is great hope to eliminate this disease by 2030; however, prevention of hepatitis C transmission through blood and blood products has a key role in elimination programs, as well (69).

**Figure 3.** Annual HCV Seroprevalence Rates and Their Trends among Blood Donors

A. Oman; B. Libya; C. Lebanon; D. Turkey; E. Iran; F. Egypt; G. Saudi Arabia.

A number of limitations exist in the present investigation that should be noted. First, the sample size in some of the Middle Eastern countries was not adequate and the quantity of data varied between countries. Second, different types of kits were utilized for detection of HCV Ab, which would have affected the obtained results of the current systematic review. Third, HCV seroprevalence rates in blood donor populations of some countries might be underestimated due to selection bias. In addition, data deficiencies existed in some studies did not allow us to include them in the final analysis.

Taken together, the present systematic review provides

detailed and reliable data on the HCV seroprevalence rates among blood donors in Middle Eastern countries. The available data showed a dramatic geographical difference in the seroprevalence of HCV among Middle Eastern blood donors. The lowest seroprevalence rates were seen in Iranian and UAE blood donors, while the highest rate was observed in Egyptian blood donors. The trend of HCV infection among Middle Eastern blood donors was decreasing in recent years. It is suggesting recent safety measures implemented in Middle Eastern countries have been effective.

## Footnote

**Authors' Contribution:** Concept and design: Seyed Moayed Alavian, Heidar Sharafi; data acquisition: Heidar Sharafi, Hossein Ghaderi-Zefrehi, Alireza Farasat, Fatemeh Jahanpeyma; data analysis: Mohammad Gholami-Fesharaki; drafting the manuscript: Hossein Ghaderi-Zefrehi, Heidar Sharafi, Farzin Sadeghi, Alireza Farasat, Fatemeh Jahanpeyma; critical revising of the manuscript: Heidar Sharafi; final approval of the manuscript: Seyed Moayed Alavian.

## References

- Karim F, Nasar A, Alam I, Alam I, Hassan S, Gul R, et al. Incidence of Active HCV infection amongst Blood Donors of Mardan District, Pakistan. *Asian Pac J Cancer Prev*. 2016;**17**(1):235-8. [PubMed: 26838216].
- Makroo RN, Walia RS, Chowdhry M, Bhatia A, Hegde V, Rosamma NL. Seroprevalence of anti-HCV antibodies among blood donors of north India. *Indian J Med Res*. 2013;**138**:125-8. [PubMed: 24056566].
- Irani-Hakime N, Aoun J, Khoury S, Samaha HR, Tamim H, Almawi WY. Seroprevalence of hepatitis C infection among health care personnel in Beirut, Lebanon. *Am J Infect Control*. 2001;**29**(1):20-3. [PubMed: 11172314].
- Keshvari M, Sharafi H, Alavian SM, Mehrabadi H, Zolfaghari S. Prevalence and trends of transfusion-transmitted infections among blood donors in Tehran, Iran from 2008 to 2013. *Transfus Apher Sci*. 2015;**53**(1):38-47. doi: 10.1016/j.transci.2015.03.003. [PubMed: 25892591].
- Keshvari M, Hajibeigi B, Azarkeivan A, Keyvani H, Behnava B, Saiedi Hosseini SY, et al. Seroprevalence of human T-cell lymphotropic virus among Iranian adult thalassemic patients. *Transfus Med*. 2014;**24**(4):227-32. doi: 10.1111/tme.12133. [PubMed: 25124072].
- Thrift AP, El-Serag HB, Kanwal F. Global epidemiology and burden of HCV infection and HCV-related disease. *Nat Rev Gastroenterol Hepatol*. 2017;**14**(2):122-32. doi: 10.1038/nrgastro.2016.176. [PubMed: 27924080].
- Alavian SM, Rezaee-Zavareh MS. The Middle East and hepatitis C virus infection: does it need special attention?. *Lancet Infect Dis*. 2016;**16**(9):1006-7. doi: 10.1016/S1473-3099(16)30264-X. [PubMed: 27684342].
- Fallahian F, Najafi A. Epidemiology of hepatitis C in the Middle East. *Saudi J Kidney Dis Transpl*. 2011;**22**(1):1-9. [PubMed: 21196607].
- Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev*. 2015;**4**:1. doi: 10.1186/2046-4053-4-1. [PubMed: 25554246].
- Ghavanini AA, Sabri MR. Hepatitis B surface antigen and anti-hepatitis C antibodies among blood donors in the Islamic Republic of Iran. *East Mediterr Health J*. 2000;**6**(5-6):1114-6. [PubMed: 12197336].
- Alavian SM, Gholami B, Masarrat S. Hepatitis C risk factors in Iranian volunteer blood donors: a case-control study. *J Gastroenterol Hepatol*. 2002;**17**(10):1092-7. [PubMed: 12201870].
- Moniri R, Mosayebii Z, Mossavi GA. Seroprevalence of cytomegalovirus, hepatitis B, hepatitis C and human immunodeficiency virus antibodies among volunteer blood donors. *Iran J Public Health*. 2004;**33**(4):38-42.
- Pourshams A, Malekzadeh R, Monavvari A, Akbari MR, Mohamadkhani A, Yarahmadi S, et al. Prevalence and etiology of persistently elevated alanine aminotransferase levels in healthy Iranian blood donors. *J Gastroenterol Hepatol*. 2005;**20**(2):229-33. doi: 10.1111/j.1440-1746.2004.03511.x. [PubMed: 15683425].
- FariborzMansour G, MohammadSadegh F, Reyhaneh J, Farahnaz J, Afshin S, Ramin T. Prevalence of hepatitis B surface antigen and hepatitis C virus antibody and their risk factors among Guilan's volunteer blood donors (1998-2003). *Hepat Mon*. 2007;**2007**(4, Autumn):239-41.
- Hosseien K, Miri SM, Amini M, Abolghasemi H, Hajibeigi B, Alaeddini F, et al. Trends in seroprevalence of hepatitis B, hepatitis C, HIV, and syphilis infections in Iranian blood donors from 2003 to 2005. *Hepat Mon*. 2009;**2009**(1, Winter):24-8.
- Kafi-abad SA, Rezvan H, Abolghasemi H, Talebian A. Prevalence and trends of human immunodeficiency virus, hepatitis B virus, and hepatitis C virus among blood donors in Iran, 2004 through 2007. *Transfusion*. 2009;**49**(10):2214-20. doi: 10.1111/j.1537-2995.2009.02245.x. [PubMed: 19527477].
- Kasraian L. National disasters in Iran and blood donation: Bam earthquake experience. *Iran Red Crescent Med J*. 2010;**2010**(3):316-8.
- Sofian M, Aghakhani A, Izadi N, Banifazl M, Kalantar E, Eslamifard A, et al. Lack of occult hepatitis B virus infection among blood donors with isolated hepatitis B core antibody living in an HBV low prevalence region of Iran. *Int J Infect Dis*. 2010;**14**(4):e308-10. doi: 10.1016/j.ijid.2009.05.011. [PubMed: 19656713].
- Bozorgi SH, Ramezani H, Nooranipour M, Ahmadi M, Baghernejad A, Mostajeri A, et al. Risk factors of viral hepatitis: yet to explore. *Transfus Apher Sci*. 2012;**47**(2):145-9. doi: 10.1016/j.transci.2012.06.023. [PubMed: 22858443].
- Farshadpour F, Taherkhani R, Tajbakhsh S, Gholizadeh Tangestani M, Hajiani G, Sharifi N, et al. Prevalence and Trends of Transfusion-Transmissible Viral Infections among Blood Donors in South of Iran: An Eleven-Year Retrospective Study. *PLoS One*. 2016;**11**(6). e0157615. doi: 10.1371/journal.pone.0157615. [PubMed: 27309959].
- Kasraian L, Karimi MH. A study on confidential unit exclusion at Shiraz Blood Transfusion Center, Iran. *Asian J Transfus Sci*. 2016;**10**(2):132-5. doi: 10.4103/0973-6247.187939. [PubMed: 27605850].
- Sakarya S, Oncu S, Ozturk B, Oncu S. Effect of preventive applications on prevalence of hepatitis B virus and hepatitis C virus infections in West Turkey. *Saudi Med J*. 2004;**25**(8):1070-2. [PubMed: 15322600].
- Guroel E, Saban C, Oral O, Cigdem A, Armagan A. Trends in hepatitis B and hepatitis C virus among blood donors over 16 years in Turkey. *Eur J Epidemiol*. 2006;**21**(4):299-305. doi: 10.1007/s10654-006-0001-2. [PubMed: 16685581].
- Dilek I, Demir C, Bay A, Akdeniz H, Oner AF. Seropositivity rates of HBsAg, anti-HCV, anti-HIV and VDRL in blood donors in Eastern Turkey. *Turk J Haematol*. 2007;**24**(1):4-7. [PubMed: 27263477].
- Acar A, Kemahli S, Altunay H, Kosan E, Oncul O, Gorenek L, et al. HBV, HCV and HIV seroprevalence among blood donors in Istanbul, Turkey: how effective are the changes in the national blood transfusion policies?. *Braz J Infect Dis*. 2010;**14**(1):41-6. [PubMed: 20428653].
- Dayan S, Tekin A, Tekin R, Dal T, Hosoglu S, Yazgan UC, et al. HBsAg, anti-HCV, anti-HIV 1/2 and syphilis seroprevalence in healthy volunteer blood donors in southeastern Anatolia. *J Infect Dev Ctries*. 2013;**7**(9):665-9. doi: 10.3855/jidc.2835. [PubMed: 23491864].
- Kucukbayrak A, Cakmak S, Hakyemez IN, Tas T, Akdeniz H. Determining immunoassay cutoff value using Western blot results to predict hepatitis C infection in blood donors with low-titer anti-HCV reactivity. *Folia Microbiol (Praha)*. 2013;**58**(4):343-7. doi: 10.1007/s12223-012-0215-5. [PubMed: 23208738].
- Uzun B, Gungor S, Demirci M. Seroprevalence of transfusion transmissible infections among blood donors in western part of Turkey: a six-year study. *Transfus Apher Sci*. 2013;**49**(3):511-5. doi: 10.1016/j.transci.2013.02.039. [PubMed: 23491864].
- Yildiz SM, Candevir A, Kibar F, Karaboga G, Turhan FT, Kis C, et al. Hepatitis B, Hepatitis C, Human immunodeficiency virus and syphilis frequency among blood donors: A single center study. *Transfus Apher Sci*. 2015;**53**(3):308-14. doi: 10.1016/j.transci.2015.05.022. [PubMed: 26070837].
- Bervil SS, Andrews VJ, Sasich F. Second-generation anti-HCV screening in a Saudi Arabian donor population. *Vox Sang*. 1994;**66**(1):33-6. [PubMed: 8146980].

31. Shobokshi OA, Serebour FE, Al-Drees AZ, Mitwalli AH, Qahtani A, Skakni LI. Hepatitis C virus seroprevalence rate among Saudis. *Saudi Med J*. 2003;**24 Suppl 2**:S81-6. [PubMed: [12897907](#)].
32. Bashawri LA, Fawaz NA, Ahmad MS, Qadi AA, Almawi WY. Prevalence of seromarkers of HBV and HCV among blood donors in eastern Saudi Arabia, 1998-2001. *Clin Lab Haematol*. 2004;**26**(3):225-8. doi: [10.1111/j.1365-2257.2004.00601.x](#). [PubMed: [15163322](#)].
33. Mohammed Abdullah S. Prevalence of hepatitis B and C in donated blood from the jazan region of Saudi Arabia. *Malays J Med Sci*. 2013;**20**(2):41-6. [PubMed: [23983576](#)].
34. Kilany M. Seroprevalence of anti-Treponemapallidum antibodies (Syphilis) in blood donors in the southern area of Saudi Arabia. *Res J Pharm Biol Chem Sci*. 2015;**6**(1):549-56.
35. Elbjairami W, Al-Jedani H, Arsheed N, Elnagdi N, Abou Eisha H, Abdulwahab A, et al. Prevalence and trends of HBV, HCV, and HIV serological and NAT markers and profiles in Saudi blood donors. *Haematologica*; 2015.
36. AlMutairi HH, AlAhmari MM, Al-Zahrani BH, Abbas IS, Al Ghamdi JA, Raja A, et al. Prevalence of serological markers and nucleic acid for blood-borne viral infections in blood donors in Al-Baha, Saudi Arabia. *J Infect Dev Ctries*. 2016;**10**(6):619-25. doi: [10.3855/jidc.6666](#). [PubMed: [27367011](#)].
37. Bernvil SS, Andrews VJ, Kariem AA. Hepatitis C antibody prevalence in Saudi Arabian blood donor population. *Ann Saudi Med*. 1991;**11**(5):563-7. [PubMed: [17590794](#)].
38. El Damaty SI, Hassan SK, Mohamed MK, El Hosini M, Rekecewicz C, Fontanet A. Surveillance system for HCV infection: testing a model based on blood banks. *J Egypt Public Health Assoc*. 2007;**82**(5-6):451-71. [PubMed: [18706299](#)].
39. El-Zayadi AR, Ibrahim EH, Badran HM, Saeid A, Moneib NA, Shemis MA, et al. Anti-HBc screening in Egyptian blood donors reduces the risk of hepatitis B virus transmission. *Transfus Med*. 2008;**18**(1):55-61. doi: [10.1111/j.1365-3148.2007.00806.x](#). [PubMed: [18279193](#)].
40. Khattab MA, Eslam M, Sharwae MA, Hamdy L. Seroprevalence of hepatitis C and B among blood donors in Egypt: Minya Governorate, 2000-2008. *Am J Infect Control*. 2010;**38**(8):640-1. doi: [10.1016/j.ajic.2009.12.016](#). [PubMed: [20400204](#)].
41. Wasfi OA, Sadek NA. Prevalence of hepatitis B surface antigen and hepatitis C virus antibodies among blood donors in Alexandria, Egypt. *East Mediterr Health J*. 2011;**17**(3):238-42. [PubMed: [21735965](#)].
42. Hussein E. Blood donor recruitment strategies and their impact on blood safety in Egypt. *Transfus Apher Sci*. 2014;**50**(1):63-7. doi: [10.1016/j.transci.2013.11.005](#). [PubMed: [24325889](#)].
43. Abdel Messih IY, Ismail MA, Saad AA, Azer MR. The degree of safety of family replacement donors versus voluntary non-remunerated donors in an Egyptian population: a comparative study. *Blood Transfus*. 2014;**12**(2):159-65. doi: [10.2450/2012.0115-12](#). [PubMed: [23245714](#)].
44. Othman BM, Monem FS. Prevalence of hepatitis C virus antibodies among intravenous drug abusers and prostitutes in Damascus, Syria. *Saudi Med J*. 2002;**23**(4):393-5. [PubMed: [11953762](#)].
45. Muselmani W, Habbal W, Monem F. Significance of screening antibodies to hepatitis B virus core antigen among Syrian blood donors. *Transfus Med*. 2013;**23**(4):265-8. doi: [10.1111/tme.12043](#). [PubMed: [23621787](#)].
46. Daw MA, Elkaber MA, Drah AM, Werfalli MM, Mihat AA, Siala IM. Prevalence of hepatitis C virus antibodies among different populations of relative and attributable risk. *Saudi Med J*. 2002;**23**(11):356-60. [PubMed: [12506296](#)].
47. Daw MA, Shabash A, El-Bouzedi A, Dau AA, Habas M, Libyan Study Group of H, et al. Modelling the prevalence of hepatitis C virus amongst blood donors in Libya: An investigation of providing a preventive strategy. *World J Virol*. 2016;**5**(1):14-22. doi: [10.5501/wjv.v5.i1.14](#). [PubMed: [26870670](#)].
48. Harmankaya O, Cetin B, Obek A, Seber E. Low prevalence of hepatitis C virus infection in hemodialysis units: effect of isolation?. *Ren Fail*. 2002;**24**(5):639-44. [PubMed: [12380910](#)].
49. Al-Waleedi AA, Khader YS. Prevalence of hepatitis B and C infections and associated factors among blood donors in Aden City, Yemen. *East Mediterr Health J*. 2012;**18**(6):624-9. [PubMed: [22888620](#)].
50. Cengiz K, Gunaydiotan M, Bedir A, Pekbay A. Low prevalence of hepatitis G virus antibodies in glomerular diseases. *Nephron*. 1998;**79**(4):472-3. [PubMed: [9689165](#)].
51. Yarom N, Dagon N, Shinar E, Gorsky M. Association between hepatitis C virus infection and oral lichen planus in Israeli patients. *Isr Med Assoc J*. 2007;**9**(5):370-2. [PubMed: [17591375](#)].
52. Irani-Hakime N, Musharrafieh U, Samaha H, Almawi WY. Prevalence of antibodies against hepatitis B virus and hepatitis C virus among blood donors in Lebanon, 1997-2003. *Am J Infect Control*. 2006;**34**(4):241-3. doi: [10.1016/j.ajic.2005.06.009](#). [PubMed: [16679184](#)].
53. Al Dhahry SH, Nograles JC, Rajapakse SM, Al Toqi FS, Kaminski GZ. Laboratory diagnosis of viral hepatitis C: The Sultan Qaboos University Hospital experience. *J Sci Res Med Sci*. 2003;**5**(1-2):15-20. [PubMed: [24019730](#)].
54. Altindis M, Yilmaz S, Dikengil T, Acemoglu H, Hosoglu S. Seroprevalence and genotyping of hepatitis B, hepatitis C and HIV among healthy population and Turkish soldiers in Northern Cyprus. *World J Gastroenterol*. 2006;**12**(42):6792-6. [PubMed: [17106927](#)].
55. Al Shaer L, AbdulRahman M, John TJ, AlHashimi A. Trends in prevalence, incidence, and residual risk of major transfusion-transmissible viral infections in United Arab Emirates blood donors: impact of individual-donation nucleic acid testing, 2004 through 2009. *Transfusion*. 2012;**52**(11):2300-9. doi: [10.1111/j.1537-2995.2012.03740.x](#). [PubMed: [22691239](#)].
56. Mohamoud YA, Riome S, Abu-Raddad LJ. Epidemiology of hepatitis C virus in the Arabian Gulf countries: Systematic review and meta-analysis of prevalence. *Int J Infect Dis*. 2016;**46**:116-25. doi: [10.1016/j.ijid.2016.03.012](#). [PubMed: [26996460](#)].
57. Ghaderi-Zefrehi H, Gholami-Fesharaki M, Sharafi H, Sadeghi F, Alavian SM. The Distribution of Hepatitis C Virus Genotypes in Middle Eastern Countries: A Systematic Review and Meta-Analysis. *Hepat Mon*. 2016;**16**(9). e40357. doi: [10.5812/hepatmon.40357](#). [PubMed: [27826320](#)].
58. Sadeghi F, Salehi-Vaziri M, Almasi-Hashiani A, Gholami-Fesharaki M, Pakzad R, Alavian SM. Prevalence of Hepatitis C Virus Genotypes Among Patients in Countries of the Eastern Mediterranean Regional Office of WHO (EMRO): A Systematic Review and Meta-Analysis. *Hepat Mon*. 2016;**16**(4). e35558. doi: [10.5812/hepatmon.35558](#). [PubMed: [27274353](#)].
59. Hesamizadeh K, Alavian SM, Najafi Tireh Shabankareh A, Sharafi H. Molecular Tracing of Hepatitis C Virus Genotype 1 Isolates in Iran: A NS5B Phylogenetic Analysis with Systematic Review. *Hepat Mon*. 2016;**16**(12). e42938. doi: [10.5812/hepatmon.42938](#). [PubMed: [28123445](#)].
60. Wasley A, Alter MJ. Epidemiology of hepatitis C: geographic differences and temporal trends. *Semin Liver Dis*. 2000;**20**(1):1-16. [PubMed: [10895428](#)].
61. Riou J, Ait Ahmed M, Blake A, Vozlinsky S, Brichler S, Eholie S, et al. Hepatitis C virus seroprevalence in adults in Africa: a systematic review and meta-analysis. *J Viral Hepat*. 2016;**23**(4):244-55. doi: [10.1111/jvh.12481](#). [PubMed: [26477881](#)].
62. Eita N. Prevalence of HCV and HBV infections among blood donors in Dakahlia, Egypt. *Vox Sanguinis*. 2009;**96**:106-7.
63. Rushdy O, Moftah F, Zakareya S. Transmitted transfused viral infections among blood donors during years 2006 and 2007 in suez canal area, Egypt. *Vox Sanguinis*. 2009;**96**:86-7.
64. Mohamoud YA, Mumtaz GR, Riome S, Miller D, Abu-Raddad LJ. The epidemiology of hepatitis C virus in Egypt: a systematic review and data synthesis. *BMC Infect Dis*. 2013;**13**:288. doi: [10.1186/1471-2334-13-288](#). [PubMed: [23799878](#)].
65. Khodabandehloo M, Roshani D, Sayehmiri K. Prevalence and trend of hepatitis C virus infection among blood donors in Iran: A systematic review and meta-analysis. *J Res Med Sci*. 2013;**18**(8):674-82. [PubMed: [23799878](#)].

- 24379843].
66. Fadlalla FA, Mohamoud YA, Mumtaz GR, Abu-Raddad LJ. The epidemiology of hepatitis C virus in the Maghreb region: systematic review and meta-analyses. *PLoS One*. 2015;**10**(3). e0121873. doi: [10.1371/journal.pone.0121873](https://doi.org/10.1371/journal.pone.0121873). [PubMed: [25803848](https://pubmed.ncbi.nlm.nih.gov/25803848/)].
  67. Chaabna K, Kouyoumjian SP, Abu-Raddad LJ. Hepatitis C Virus Epidemiology in Djibouti, Somalia, Sudan, and Yemen: Systematic Review and Meta-Analysis. *PLoS One*. 2016;**11**(2). e0149966. doi: [10.1371/journal.pone.0149966](https://doi.org/10.1371/journal.pone.0149966). [PubMed: [26900839](https://pubmed.ncbi.nlm.nih.gov/26900839/)].
  68. Sievert W, Altraif I, Razavi HA, Abdo A, Ahmed EA, Alomair A, et al. A systematic review of hepatitis C virus epidemiology in Asia, Australia and Egypt. *Liver Int*. 2011;**31** Suppl 2:61-80. doi: [10.1111/j.1478-3231.2011.02540.x](https://doi.org/10.1111/j.1478-3231.2011.02540.x). [PubMed: [21651703](https://pubmed.ncbi.nlm.nih.gov/21651703/)].
  69. Hesamizadeh K, Sharafi H, Rezaee-Zavareh MS, Behnava B, Alavian SM. Next Steps Toward Eradication of Hepatitis C in the Era of Direct Acting Antivirals. *Hepat Mon*. 2016;**16**(4). e37089. doi: [10.5812/hepatmon.37089](https://doi.org/10.5812/hepatmon.37089). [PubMed: [27275164](https://pubmed.ncbi.nlm.nih.gov/27275164/)].
  70. Dolatimehr F, Karimi-Sari H, Rezaee-Zavareh MS, Alavian SM, Behnava B, Gholami-Fesharaki M, et al. Combination of sofosbuvir, pegylated-interferon and ribavirin for treatment of hepatitis C virus genotype 1 infection: a systematic review and meta-analysis. *Daru*. 2017;**25**(1):11. doi: [10.1186/s40199-017-0177-x](https://doi.org/10.1186/s40199-017-0177-x). [PubMed: [28427463](https://pubmed.ncbi.nlm.nih.gov/28427463/)].
  71. Alavian SM, Hajarizadeh B, Bagheri Lankarani K, Sharafi H, Ebrahimi Daryani N, Merat S, et al. Recommendations for the Clinical Management of Hepatitis C in Iran: A Consensus-Based National Guideline. *Hepat Mon*. 2016;**16**(8). e40959. doi: [10.5812/hepatmon.guideline](https://doi.org/10.5812/hepatmon.guideline). [PubMed: [27799966](https://pubmed.ncbi.nlm.nih.gov/27799966/)].
  72. Rezaee-Zavareh MS, Hesamizadeh K, Behnava B, Alavian SM, Gholami-Fesharaki M, Sharafi H. Combination of Ledipasvir and Sofosbuvir for Treatment of Hepatitis C Virus Genotype 1 Infection: Systematic Review and Meta-Analysis. *Ann Hepatol*. 2017;**16**(2):188-97. doi: [10.5604/16652681.1231562](https://doi.org/10.5604/16652681.1231562). [PubMed: [28233739](https://pubmed.ncbi.nlm.nih.gov/28233739/)].
  73. Sharafi H, Nikbin M, Alavian SH, Behnava B, Alavian SM. Efficacy and Safety of Generic Sofosbuvir/Ledipasvir Fixed-Dose Combination in Iranian Patients with Chronic Hepatitis C Virus Infection. *Hepat Mon*. 2017;**17**(6). doi: [10.5812/hepatmon.12216](https://doi.org/10.5812/hepatmon.12216).



**Table 1.** Characteristics of Included Studies from Middle Eastern Countries

Country	City, Province or Region	Reference Number	Screening Test	Confirmatory Test	Year of Publish	Year of Study	Sample Size, n	HCV Ab Rate, %	Donor Type
Cyprus	Nicosia and Kyrenia	(54)	EIA $\geq$ 3	NA	2006	2000	5057	0.45	NA
<b>Total<sup>a</sup></b>	0.45% (0.30% - 0.68%)								
Egypt	Cairo	(38)	EIA $\geq$ 3	EIA $\geq$ 3	2007	1999 - 2000	2146	8.90	Replacement donor
	Cairo	(39)	EIA $\geq$ 3	EIA $\geq$ 3	2008	2005	760	5.00	Voluntary donor
	Minya governorate	(40)	EIA $\geq$ 3	NA	2010	2000 - 2008	21772	9.03	Voluntary donor
	Alexandria	(41)	EIA $\geq$ 3	EIA $\geq$ 3	2011	2007 - 2008	3420	3.48	Voluntary donor
	Cairo	(42)	EIA $\geq$ 3	NA	2014	2006 - 2012	308762	4.34	Voluntary or replacement donor
<b>Total<sup>a</sup></b>	5.76% (3.30% - 8.22%)								
Iran	Shiraz	(10)	EIA < 3	RIBA	2000	1998	7879	0.60	Voluntary donor
	Tehran	(11)	EIA $\geq$ 3	RIBA	2002	1996 - 1998	319375	0.09	Voluntary donor
	Kashan	(12)	EIA $\geq$ 3	NA	2004	2001 - 2002	600	0.50	Voluntary donor
	Tehran	(13)	EIA < 3	RIBA	2005	2001 - 2002	1959	0.46	Voluntary donor
	Guilan	(14)	EIA $\geq$ 3	RIBA	2007	1998 - 2003	221508	0.32	Voluntary donor
	Tehran	(15)	EIA $\geq$ 3	RIBA	2009	2003 - 2005 (2003) <sup>b</sup>	329516	0.14	Voluntary donor
	Iran	(16)	EIA $\geq$ 3	RIBA	2009	2004 - 2007	6499851	0.13	Voluntary donor
	Shiraz	(17)	EIA $\geq$ 3	RIBA	2010	2003	1933	0.52	Voluntary donor
	Arak	(18)	EIA $\geq$ 3	NA	2010	2008	531	0.19	Voluntary donor
	Qazvin	(19)	EIA $\geq$ 3	RIBA	2012	2009	20591	0.17	NA
	Tehran	(4)	EIA $\geq$ 3	RIBA	2015	2008 - 2013	1796090	0.10	Voluntary donor
	Bushehr, Borazjan and other cities	(20)	EIA $\geq$ 3	RIBA	2016	2004 - 2014 (2008 - 2014) <sup>b</sup>	219552	0.06	Voluntary donor
Shiraz	(21)	EIA $\geq$ 3	RIBA	2016	2012 - 2013	101245	0.05	NA	
<b>Total<sup>a</sup></b>	0.14% (0.12% - 0.17%)								
Israel	Tel HaShomer	(50)	EIA < 3	RIBA	1998	1994 - 1996	193806	0.20	Voluntary donor
	Tel HaShomer	(51)	EIA $\geq$ 3	RIBA	2007	2003	225452	0.11	Voluntary donor
<b>Total<sup>a</sup></b>	0.16% (0.06% - 0.25%)								
Lebanon	Beirut	(52)	EIA $\geq$ 3	EIA $\geq$ 3	2006	1997 - 2003	16084	0.40	NA
<b>Total<sup>a</sup></b>	0.4% (0.31% - 0.51%)								
Libya	Tripoli	(46)	EIA $\geq$ 3	NA	2002	1999 - 2001	1200	1.17	NA
	Tripoli	(47)	EIA $\geq$ 3	NA	2016	2008 - 2013	1008214	1.78	Voluntary donor
<b>Total<sup>a</sup></b>	1.56% (0.99% - 2.13%)								
Oman	Muscat	(53)	EIA2&3	RIBA	2003	1991 - 2001 (1995 - 2001) <sup>b</sup>	2131	0.75	NA
<b>Total<sup>a</sup></b>	0.75% (0.64% - 0.88%)								
Saudi	Riyadh	(37)	EIA $\geq$ 3	NA	1991	1991	10646	1.01	NA
	Riyadh	(30)	EIA < 3	RIBA	1994	1992 - 1993	18540	0.44	NA
	Bisha, El-Baha, Najran, Jizan, Aseer, Hafer El-Baten, Al-Hessa and El-Sharkia, Al-Qassim, Riyadh, Jeddah, Makkah, Medinah, Taif, El-Qunfuda, Tabuk, Al-Goofe, El-Quriate, Hail, El-Shamalia	(31)	EIA $\geq$ 3	RIBA	2003	1998 - 2002	557813	1.13	NA
	Eastern Saudi Arabia	(32)	EIA $\geq$ 3	NA	2004	1998 - 2001	13443	1.44	Voluntary, replacement or employment/licensing donor

	Jazan	(33)	EIA $\geq$ 3	EIA $\geq$ 3	2013	2004 - 2009	29949	0.42	Voluntary or replacement donor
	Aseer Region	(34)	EIA $\geq$ 3	EIA $\geq$ 3	2015	2012 - 2013	7267	0.07	Voluntary donor
	Makkah	(35)	EIA $\geq$ 3	NA	2015	2011 - 2014	22963	0.44	Voluntary or replacement donor
	Al-Baha	(36)	EIA $\geq$ 3	RIBA	2016	2009 - 2011	2807	0.04	NA
<b>Total<sup>a</sup></b>	0.62% (0.23% - 1.01%)								
<b>Syria</b>	Damascus	(44)	EIA $\geq$ 3	EIA $\geq$ 3	2002	2002	2100	0.95	NA
	Damascus	(45)	EIA $\geq$ 3	NA	2013	2011	1939	0.41	NA
<b>Total<sup>a</sup></b>	0.66% (0.13% - 1.19%)								
<b>Turkey</b>	Aydin	(22)	EIA	RIBA	2004	1993 - 2002	37866	0.19	Voluntary donor
	Denizli, Istanbul, Izmir, Adana, Bursa, Samsun, Eskisehir, Erzurum, Diyarbakir, Gaziantep, Sivas, Zonguldak, Trabzon, Antalya, Konya, Duzce, Balikesir, Sanliurfa, Kayseri, Isparta and Zeynep Kamil	(23)	EIA $\geq$ 3	RIBA	2006	1989-2004	6240130	0.38	Voluntary or family donor or military personnel
	Van	(24)	EIA $\geq$ 3	NA	2007	1995 - 2003	39002	0.17	NA
	Istanbul	(25)	EIA $\geq$ 3	RIBA	2010	2007	72695	0.07	NA
	Southeastern Anatolia	(26)	EIA $\geq$ 3	NA	2013	2000 - 2010 (2005 - 2010) <sup>b</sup>	155690	0.67	Voluntary donor
	Ankara	(27)	EIA $\geq$ 3	RIBA	2013	2004 - 2010	15540	0.22	Voluntary donor
	Izmir	(28)	EIA $\geq$ 3	EIA $\geq$ 3	2013	2004 - 2010 (2005 - 2010) <sup>b</sup>	73213	0.41	Voluntary or replacement donor
	Adana	(29)	EIA $\geq$ 3	RIBA	2015	2007 - 2009	62461	0.05	Voluntary or replacement donor
	<b>Total<sup>a</sup></b>	0.27% (0.13% - 0.41%)							
<b>Yemen</b>	Hajjah	(48)	EIA $\geq$ 3	NA	2002	1999	2434	1.07	Replacement donor
	Aden	(49)	EIA $\geq$ 3	NA	2012	2007	469	1.28	Replacement donor
<b>Total<sup>a</sup></b>	1.09% (0.69% - 1.50%)								
<b>UAE</b>	Dubai	(55)	ELFA	RIBA	2012	2004 - 2009	169781	0.11	NA
<b>Total<sup>a</sup></b>	0.11% (0.10% - 0.13%)								
<b>Pooled<sup>c</sup></b>	0.88% (0.79% - 0.97%)								

Abbreviations: EIA, enzyme immunoassay; ELFA, enzyme-linked fluorescence assay; NA, not available; RIBA, recombinant immunoblot assay.  
<sup>a</sup>Total HCV seroprevalence in the country and its 95% confidence interval.

<sup>b</sup>In the studies with temporal and spatial overlap, the time span of study with the included data in this meta-analysis.

<sup>c</sup>Pooled HCV seroprevalence in the Middle Eastern countries and its 95% confidence interval.