

The Sensitivity of the PCR Method for Detection of *Coxiella burnetii* in the Milk Samples

Mohammad Kargar^{1,*}; Afsaneh Rashidi¹; Abbass Doosti²; Akram Najafi³; Sadegh Ghorbani-Dalini⁴

¹Department of Microbiology, Jahrom Branch, Islamic Azad University, Jahrom, IR Iran

²Biotechnology Research Center, Shahrekord Branch, Islamic Azad University, Shahrekord, IR Iran

³Persian Gulf Marine Biotechnology Medicine Research Center, Bushehr University of Medical Sciences, Bushehr, IR Iran

⁴Department of Microbiology, Jahrom Branch, Young Researcher's Club, Islamic Azad University, Jahrom, IR Iran

*Corresponding author: Mohammad Kargar, Department of Microbiology, Jahrom Branch, Islamic Azad University, Jahrom, IR Iran. E-mail: mkargar@jia.ac.ir

Received: January 13, 2014; Accepted: April 11, 2014

Background: *Coxiella burnetii* is an obligate intracellular bacterium that causes the zoonotic disease Q fever with a worldwide distribution. Also *C. burnetii* is classified as a bioterrorism agent. In order to management, prevention and control of Q fever the fast and accurate detection of *C. burnetii* is necessary. However, the isolation of this strain is very difficult and dangerous.

Objectives: The aim of the study was to evaluate the sensitivity of PCR using different primers for the detection of *C. burnetii* in milk samples.

Materials and Methods: In this cross-sectional study 70 bovine bulk milk samples were collected randomly from dairy herds in Jahrom, Iran in 2010. All the samples were analyzed for the presence of *C. burnetii* by PCR targeting 3 different genes (Trans, OMP, Coc). The PCR products were examined by electrophoresis using an agarose gel.

Results: The frequency of *C. burnetii* in the evaluated samples using Trans-PCR, OMP-PCR and Coc-PCR were 17.14%, 10% and 10%, respectively.

Conclusions: The results of this study show that Trans-PCR is highly sensitive and useful for the direct detection of *C. burnetii* in milk samples. This technique is a one-step and fast process in comparison to the other assays.

Keywords: Q fever; *Coxiella burnetii*; Polymerase Chain Reaction (PCR); Bovine milk

1. Background

Coxiella burnetii is an obligate intracellular bacterium that causes the zoonotic disease Q fever with a worldwide distribution. It has a wide range of hosts including mammals such as ruminants, dogs, cats, non-mammal primates, wild rodents, small mammals, big game, and non-mammals such as reptiles, amphibians, birds, fish and ticks [1, 2].

Cattle, sheep and goats are the main sources of infection in humans [3, 4]. Infected animals excrete *C. burnetii* into the environment via birth products such as the placenta. This bacterium is very stable in different environments. It is also highly infectious and one to ten organisms can cause Q fever in humans [5]. Also, *C. burnetii* can be present in milk, urine, feces, vaginal mucus and semen. In milk, it can be secreted for 8 days in ewes and up to 13 months in cattle [6]. The consumption of contaminated raw milk does not seem to represent an efficient route of disease transmission, however bulk milk samples is an important specimen for epidemiological survey on dairy herds [7]. In order to management, prevention, control and treatment of Q fever in animal and human, early and

accurate detection of *C. burnetii* is very necessary. Previous studies on the prevalence of *C. burnetii* in dairy cows were based mainly on serologic tests that detect antibodies that could have been introduced months earlier [8]. Isolation of *C. burnetii* is very difficult and dangerous. Recently, PCR has been used to detect *C. burnetii*. PCR is a safe, sensitive and specific method for the detection of *C. burnetii* in different samples [9]. Several target genes that are used for specific *C. burnetii* identification include: the superoxide dismutase (Sod B) gene, com1 encoding a 27 kDa Outer membrane protein, the heat shock operon encoding two heat shock proteins (htpA and htpB), isocitrate dehydrogenase (icd), the macrophage infectivity potentiator protein (cbmip) and a transposon-like repetitive region of the *C. burnetii* genome (Trans) [10].

2. Objectives

Since the clinically healthy cattle are the main source of *C. burnetii* infection in Iran [11], in the present study, we evaluated the sensitivity of PCR with 3 different primers for detection of *C. burnetii* in bovine bulk milk samples.

3. Materials and Methods

In this cross-sectional study 70 bovine bulk milk samples were collected randomly from dairy herds in Jahrom city in the Southern of Iran in 2010. The samples were immediately transported to the laboratory and were tested. One milliter of raw milk was centrifuged. This procedure was performed to isolate the bacterial cells in pellet of the milk samples. After removing the cream and milk layers [12], DNA was extracted from the pellet by a genomic DNA purification kit (Cinna Gen Co., Iran) according to the manufacturer's protocol. DNA samples were stored at -20°C until they were used. In this study, we used 8 genomic primers targeting 3 different genes: A) Trans₁ and Trans₂ were designed based on the transposing-like repetitive region of the *C. burnetii* genome [13]. The length of the genome target for amplification was expected to be 687 bp. B) OMP₁, OMP₂, OMP₃ and OMP₄ were designed from the nucleotide sequence of the com1 gene encoding a 27 kDa outer Membrane Protein (OMP) as previously described [14]. The expected amplification product of the target sequence with OMP₁, OMP₂ was 501 bp long and with OMP₃, OMP₄ was 438 bp long. C) The new primers Coc-F and Coc-R were designed based on the 16S rRNA gene in the present study. The length of the predicted product was 242 bp.

All oligonucleotide primers were obtained from a commercial source (Cinna Gen Co., Iran). The sequence of the primers is shown in Table 1. The Trans-PCR thermal program was carried out according to the method described in [9]. The amplification was performed in a total volume of 25 µL containing 2 µL of DNA sample, 1.5 mM MgCl₂, 0.2 mM (each) dNTPs, 0.2 µM primer Trans₁, 0.2 µM primer Trans₂ and 1 U/reaction of Smar Taq DNA polymerase (Cinna Gen Co., Iran). The thermal program was carried out under the following conditions: five cycles of 94°C for 30 second, 66 - 61°C (the temperature was decreased by 1°C between consecutive steps) for 1 min, 72°C for 1 min and then 40 cycles of 94°C for 30 s, 61°C for 30 s and 72°C for 1 min. For the Nested PCR with primers OMP₁ - OMP₂ and

OMP₃ - OMP₄, the first amplification was performed in a total volume of 25 µL, containing 2.5 µL of DNA sample, 1.5 mM MgCl₂, 0.2 mM (each) dNTPs, 1 µL primer OMP₁, 1 µL primer OMP₂, and 2.5 U/reaction of Smar Taq DNA polymerase (Cinna Gen Co., Iran). The PCR assay was done at 94°C for 4 min and then for 30 cycles of 94°C for 1 min, 56°C for 1 min, 72°C for 1 min and a final extension at 72°C for 5 min in a DNA thermal cycler. In the second amplification, the reaction mixture was the same as that in the first amplification, except for primers and DNA templates. In this amplification, primers OMP₃ OMP₄ were used and the first amplification product was used as the DNA template. The PCR assay was performed at 95°C for 4 min and then for 30 cycles of 94°C for 1 min, 57°C for 1 min, 72°C for 1 min and a final extension at 72°C for 5 min. For the semi nested PCR with primers Coc-F and Coc-R, the amplification was performed in a total volume of 25 µL containing 2 µL of DNA sample, 1.5 mM MgCl₂, 0.2 mM (each) dNTPs, 0.2 µM primer Coc-F, 0.2 µM primer Coc-R and 1 U/reaction of Smar Taq DNA polymerase (Cinna Gen, Iran).

The thermal program was carried out under the following conditions: 95°C for 5 min and then for 32 cycles of 94°C for 1 min, 58°C for 1 min, 72°C for 1 min and a final extension at 72°C for 5 min. All of PCR reactions were performed in a DNA thermal cycler (Techne). In this study, we used positive and negative controls in each PCR run. *C. burnetii* DNA (serial Number: 3154; Genekam Biotechnology AG, Duisburg, Germany) was used as the positive control and negative controls were reaction mixtures without a DNA template. Sterile distilled water was used instead of a DNA template. DNA samples of *C. burnetii* and 5 other bacteria were used in the semi-nested PCR assay in order to evaluate the specificity of new primers Coc-F and Coc-R. The bacteria used in this test were, *Staphylococcus aureus*, *Escherichia coli*, *Brucella abortus*, *Brucella melitensis*, and *Listeria monocytogenes*. The PCR-amplification products (OMP₁ - OMP₂: 501 bp; OMP₃ - OMP₄: 438 bp; trans₁ - trans₂: 687; Coc_F - Coc_R: 242 bp) were examined by electrophoresis in a 1.5% agarose gel, visualized under UV and photographed by gel documentation (U GENIUS-SYSGENE).

Table 1. The PCR Primers Used for the Detection of *Coxiella burnetii* Genes

Target Gene	Primers	Primers Sequence	Size (bp)
com1	OMP ₁	AGT AGA AGC ATC CCA AGC ATT G	501
	OMP ₂	T GAA GCG CAA CAA GAA GAA CAC	501
	OMP ₃	GC CTG CTA GCT GTA ACG ATT G	438
	OMP ₄	TTG GAA GTT ATC ACG CAG TTG	438
IS1111	Trans ₁	TAT GTA TCC ACC GTA GCC AGT C	687
	Trans ₂	CCC AAC AAC ACC TCC TTA TTC	
16S rRNA			242
	Coc-f	GTA ATA TCC TTG GGC GTT GAC G	
	Coc-r	ATC TAC GCA TTT CAC CGC TAC AC	

4. Results


4.1. Specificity of Semi-Nested PCR

In the semi-nested PCR assay with new primers Coc-F and Coc-R only one specific band was observed with the expected size (242 bp) of *C. burnetii*.

4.2. PCR


In total, 7 of 70 tested samples were positive with primers OMP₁, OMP₂, OMP₃ and OMP₄. The primers OMP₁ - OMP₂ and OMP₃ - OMP₄ amplified the predicted products of the 501 bp DNA in the first amplification and the 438 bp DNA in the second amplification of PCR (Figure 1). Also 10% of samples were positive with primers Coc_F - Coc_R and showed the 242 bp PCR product on agarose gel (Figure 2). While 12 positive samples were observed with primers Trans₁ - Trans₂, in amplification with these primers the bands appeared at approximately 687 bp, which was in line with the expected length for detection of *C. burnetii* (Figure 3). The results of the different methods are shown in Table 2.

Figure 1. Electrophoresis of PCR Product of 438 bp Fragment of the ComI Gene in *Coxiella burnetii*




M) 100 bp DNA ladder. 1) Negative control. 2- 5) Positive samples of *Coxiella burnetii*. 6) Positive control.

Figure 2. Electrophoresis of PCR Product of 242 bp Fragment of the 16S rRNA Gene in *Coxiella burnetii*



M) 100 bp DNA ladder. 1) Negative control. 2 - 4) Positive samples of *Coxiella burnetii*. 5) Positive control.

Figure 3. Electrophoresis of PCR Product of 687 bp Fragment of the IS1111 Gene in *Coxiella burnetii*



M) 100 bp DNA ladder. 1) Negative control. 2 - 6) Positive samples of *Coxiella burnetii*. 7) Positive control.

Table 2. Sensitivity and Specificity of the Different Methods for Detection of *C. burnetii*

Methods	Numbers of Steps	Numbers of Positive Samples, %	Sensitivity	Specificity
Trans-PCR	1	12 (17.14)	More than other assays	-
OMP-PCR	2	7 (10)	-	-
Coc-PCR	2	7 (10)	-	Acceptable ^a

^a In this study the specificity of new primers Coc-f and Coc-r was evaluated and only one specific band was observed with the expected size (242 bp) of *C. burnetii*.

5. Discussion

In this study we evaluated the sensitivity of PCR with 3 different primers for detection of *C. burnetii* in bovine bulk milk samples. The Trans-PCR showed more positive samples (17.14%) while the frequency of *C. burnetii* by OMP-PCR and Coc-PCR in the tested samples was 10%. This suggests that PCR with primers Trans₁ and Trans₂ are highly sensitive and useful for the detection of *C. burnetii*.

Previous studies on the prevalence of *C. burnetii* in dairy bovine were based mainly on serologic tests that detect antibodies that could have been introduced months earlier [8]. Also *C. burnetii* is classified as a select agent and a CDC (Centers for Disease Control and Prevention) category B bioterrorism agent, with current research on the agent requiring specialized high-containment biosafety level-3 facilities [15]. Recently, PCR have been used to detection of *C. burnetii*. PCR is usually used for the diagnosis of Q fever in ruminants in research and clinical works [9]. Although the prevalence of *C. burnetii* is likely to be low in milk, PCR can be used for the detection of *C. burnetii* in this matrix [13]. Several target genes are used for detection of *C. burnetii* by PCR. Most of studies have focused on detection and determination of the rate of *C. burnetii* by PCR, while a little information is available about the comparison of different primers.

The purpose of the present study was to evaluate the sensitivity of PCR with 8 different primers targeting 3 genes for the detection of *C. burnetii* in bulk milk samples. These primers were designed based on different target genes. Com1 and 16S rRNA are single copy genes while IS1111 is present at multiple copy numbers (7 to 110 copies), depending on the strains of *C. burnetii* [16]. So it was expected that some of the Trans-positive samples would be negative using the com1 and 16S rRNA assay.

In this test 7 out of 70 milk samples were positive for *C. burnetii* targeting com1 and 16S rRNA. The number of positive samples with Trans-PCR was larger than by any other assay. The use of Trans-primers for the amplification of IS1111 allows the sensitivity of the assay to be increased and this is because of the presence of several copies in the *Coxiella* genome. Also, Trans-PCR was run in one step whilst nested PCR and semi-nested PCR with other primers were run in two steps. Run of PCR in two steps is time consuming and will increase risk of contamination between two steps. Due to running the PCR in one step and the larger number of positive samples, the Trans-PCR can be more sensitive, reliable, and an easier and faster method for the detection of *C. burnetii* in bulk milk samples. These results are similar to a study in France, which Berri et al. showed Trans-PCR to be very highly specific and sensitive for the direct detection of *C. burnetii* in genital swabs, milk and fecal samples from ewes. Also they pointed that the high degree of efficacy of the trans-PCR can be attributed to the fact that the targeted region exists in at least 19 copies in the *C. burnetii* Nine Mile, phase I, genome, which gives the trans-PCR a level of sensitivity 100

times higher than that of the PCR assay [9]. In a similar study, Kim et al. in the USA showed the higher sensitivity of Trans-PCR in detection of *C. burnetii*. Also they reported that the trans-PCR assay detects *C. burnetii* in samples immediately, unlike serologic assays that detect antibodies that could have been introduced months earlier [8]. Vaidya et al. reported that the PCR assay with primers targeting IS1111, the repetitive, transposon-like element (Trans-PCR), is very specific and sensitive for the detection of *C. burnetii* in clinical samples [17]. In this research the Trans-PCR showed 17.14% positive samples while the frequency of *C. burnetii* by OMP-PCR and Coc-PCR in the tested samples was 10%. Since bulk milk samples is an important specimen for epidemiological survey on dairy herds, we evaluated the sensitivity of PCR with 3 different primers for detection of *C. burnetii* in bovine bulk milk samples. The results of this study suggest that the IS1111 assay is reliably detecting *C. burnetii* genomic DNA in milk samples and PCR with primers Trans₁ and Trans₂ are highly sensitive and useful for the detection of *C. burnetii*.

The results of this study are limited to the PCR-based methods for detection of *C. burnetii* in the bulk milk samples, so we cannot compare the specificity and sensitivity of PCR with other methods such as ELISA. This study suggests that in order to obtain reliable results, the large numbers of samples should be analyzed in subsequent studies. Also it is better to compare different methods for detection of *C. burnetii*.

Acknowledgements

The authors are grateful to the Islamic Azad University, Jahrom branch for their executive support of this project. This study was a part of MSc thesis with the number of 8735290.

Authors' Contributions

All authors had equal role in design, work, statistical analysis, and manuscript writing.

Funding/Support

Islamic Azad University, Jahrom.

References

1. Parker NR, Barralet JH, Bell AM. Q fever. *Lancet*. 2006;**367**(9511):679-88.
2. Fretz R, Schaeren W, Tanner M, Baumgartner A. Screening of various foodstuffs for occurrence of *Coxiella burnetii* in Switzerland. *Int J Food Microbiol*. 2007;**116**(3):414-8.
3. Psaroulaki A, Hadjichristodoulou C, Loukaides F, Soteriades E, Konstantinidis A, Papastergiou P, et al. Epidemiological study of Q fever in humans, ruminant animals, and ticks in Cyprus using a geographical information system. *Eur J Clin Microbiol Infect Dis*. 2006;**25**(9):576-86.
4. Angelakis E, Raoult D. Q Fever. *Vet Microbiol*. 2010;**140**(3-4):297-309.
5. Kersh GJ, Wolfe TM, Fitzpatrick KA, Candee AJ, Oliver LD, Patterson NE, et al. Presence of *Coxiella burnetii* DNA in the environment of the United States, 2006 to 2008. *Appl Environ Microbiol*. 2010;**76**(13):4469-75.

6. Roest HI, Tilburg JJ, van der Hoek W, Vellema P, van Zijderveld FG, Klaassen CH, et al. The Q fever epidemic in The Netherlands: history, onset, response and reflection. *Epidemiol Infect.* 2011;**139**(1):1-12.
7. Vicari N, Faccini S, Ricchi M, Garbarino C, Decastelli L, Boldini M, et al. Occurrence of *Coxiella burnetii* in bulk tank milk from northwestern Italy. *Vet Rec.* 2013;**172**(26):687.
8. Kim SG, Kim EH, Lafferty CJ, Dubovi E. *Coxiella burnetii* in bulk tank milk samples, United States. *Emerg Infect Dis.* 2005;**11**(4):619-21.
9. Berri M, Laroucau K, Rodolakis A. The detection of *Coxiella burnetii* from ovine genital swabs, milk and fecal samples by the use of a single touchdown polymerase chain reaction. *Vet Microbiol.* 2000;**72**(3-4):285-93.
10. World Organization for Animal Health. *Manual of diagnostic tests and vaccines for terrestrial animals 2013.* 2012. Available from: <http://www.oie.int/international-standard-setting/terrestrial-manual>.
11. Rahimi E, Doosti A, Ameri M, Kabiri E, Sharifian B. Detection of *Coxiella burnetii* by nested PCR in bulk milk samples from dairy bovine, ovine, and caprine herds in Iran. *Zoonoses Public Health.* 2010;**57**(7-8):38-41.
12. Berri M, Arricau-Bouvery N, Rodolakis A. PCR-based detection of *coxiella burnetii* from clinical samples. In: Sachse K, Frey J editors. *PCR detection of microbial pathogens (Methods in Molecular Biology).* 1st ed. New Jersey: Humana Press; 2002. pp. 153-61.
13. Hoover TA, Vodkin MH, Williams JC. A *Coxiella burnetii* repeated DNA element resembling a bacterial insertion sequence. *J Bacteriol.* 1992;**174**(17):5540-8.
14. Zhang GQ, Nguyen SV, To H, Ogawa M, Hotta A, Yamaguchi T, et al. Clinical evaluation of a new PCR assay for detection of *Coxiella burnetii* in human serum samples. *J Clin Microbiol.* 1998;**36**(1):77-80.
15. Loftis AD, Priestley RA, Massung RF. Detection of *Coxiella burnetii* in commercially available raw milk from the United States. *Foodborne Pathog Dis.* 2010;**7**(12):1453-6.
16. Klee SR, Tyczka J, Ellerbrok H, Franz T, Linke S, Baljer G, et al. Highly sensitive real-time PCR for specific detection and quantification of *Coxiella burnetii*. *BMC Microbiol.* 2006;**6**:2.
17. Vaidya VM, Malik SV, Kaur S, Kumar S, Barbuddhe SB. Comparison of PCR, immunofluorescence assay, and pathogen isolation for diagnosis of q fever in humans with spontaneous abortions. *J Clin Microbiol.* 2008;**46**(6):2038-44.

Archive of SID