



Antibacterial and Antibiofilm Potential of Sea Anemone (*Stichodactyla haddoni*) Isolated *Vibrio* (*V. parahaemolyticus* and *V. alginolyticus*), and *Pseudoalteromonas* (*P. gelatinilytica* and *P. piscicida*) Against *Staphylococcus aureus* and *Pseudomonas aeruginosa*

Neda Fazeli ¹, Akram Sadat Naeemi ^{1,*}, Seyed Amir Hossein Jalali ² and Hojjatollah Zamani ¹

¹Department of Biology, Faculty of Sciences, University of Guilan, Rasht, Iran

²Department of Natural Resources, Isfahan University of Technology, Isfahan, Iran

*Corresponding author: Department of Biology, Faculty of Sciences, University of Guilan, Rasht, Iran. Email: naeemi.marine@gmail.com

Received 2020 August 31; Revised 2021 January 23; Accepted 2021 January 30.

Abstract

Background: *Staphylococcus aureus* and *Pseudomonas aeruginosa* are important human bacterial pathogens, which are resistant to several antibiotics. One of the main causes of their resistance is the ability of biofilm formation.

Objectives: The present study aimed to evaluate the antibacterial and antibiofilm activity of the extracts of *Vibrio parahaemolyticus*, *V. alginolyticus*, *Pseudoalteromonas gelatinilytica*, and *Pseudoalteromonas piscicida* isolated from sea anemone (*Stichodactyla haddoni*) against *S. aureus* and *P. aeruginosa*.

Methods: Four isolated bacteria were identified using biochemical and molecular identification methods, and their extracts were obtained by mixing the cell-free supernatants from their old broth culture using ethyl acetate and methanol as the solvents. The agar well-diffusion and micro-dilution methods were also applied to determine the antibacterial activity, minimum bactericidal concentration (MBC), and minimum inhibitory concentration (MIC) of the extracts. The ability of the extracts to inhibit biofilm formation and disrupt the preformed biofilm of the pathogens was attained through crystal violet staining in 96-well microtiter plates. To determine the nature of the extracts, they were exposed to protease enzyme, and the antibiofilm activity was compared with the untreated extracts.

Results: The extracts of the four isolated bacteria inhibited bacterial growth and biofilm formation and disrupted the preformed biofilm of *S. aureus* (MIC = BIC = 600 µg/mL) and *P. aeruginosa* (MIC = BIC = 300 µg/mL). In addition, the active compounds of the extracts with antibiofilm activities were mainly proteases.

Conclusions: According to the results, *V. parahaemolyticus*, *V. alginolyticus*, *P. gelatinilytica*, and *P. piscicida* had antibacterial and antibiofilm potential against *S. aureus* and *P. aeruginosa*, and their extract could also be further analyzed as an alternative to antibiotics.

Keywords: Sea Anemone-Isolated Bacteria, MIC, Biofilm, *Staphylococcus aureus*, *Pseudomonas aeruginosa*

1. Background

Staphylococcus aureus is an opportunistic pathogen and the causative agent of the most common healthcare-related infections; the extent of these infections could range from a simple skin lesion to an acute infection (1). *S. aureus*-related infections are often recurrent and a leading cause of mortality and morbidity in medical facilities. Failure in the treatment of these infections is mainly due to resistance to multiple antibiotics and the diverse range of biofilm-induced virulence factors (2).

Pseudomonas aeruginosa is a Gram-negative opportunistic pathogen that causes various infections in the

community, especially in patients with immunodeficiency, cancer, AIDS, cystic fibrosis, and burns. The pathogenicity of *P. aeruginosa* depends on the production of several virulence factors, including pili, phallus, non-pili adhesins, exotoxins, protease, and biofilm. One of the therapeutic complications of this bacterium is its antibiotic resistance to common antibiotic therapies, which in turn leads to biofilm production (3). The pathogenesis, survival, and colonization of the cells in the biofilm of *P. aeruginosa* rely on cell communications (4). In addition, three different polysaccharides (mainly alginate) are responsible for the high viscoplasticity of *P. aeruginosa* biofilm structure and

stability (5).

To date, no treatments have effectively targeted microbial biofilms due to the inherent resistance of biofilms to antibiotics (6). Therefore, there is an urgent need for new drugs that are not only active against planktonic bacteria, but also effective against drug-resistant biofilms. Among the potential organisms that are able to produce natural compounds, microorganisms (particularly marine bacteria) are considered as prolific producers of antimicrobials (7). Marine benthic fauna have soft and unprotected bodies, live a sessile lifestyles, and are home to diverse microorganisms. The chemical compounds produced by symbiotic bacteria play a pivotal role in the host immune system (8). Several studies have also demonstrated the antimicrobial potential of these chemical compounds against human pathogens (8-10).

2. Objectives

The present study aimed to evaluate the antibacterial and antibiofilm activities of the extracts derived from sea anemone (*Stichodactyla haddoni*)-isolated *Vibrio* and *Pseudoalteromonas* against *S. aureus* and *P. aeruginosa*.

3. Methods

3.1. Isolation and Cultivation of SIB

The procedure was performed in accordance with the method described by (11). Initially, two grams of tissue was separated from sea anemone, washed three times with sterile sea water, homogenized, and serially diluted. Afterwards, 100 microliters of each diluted sample was spread on Marine agar (MA; Difco), and the plates were incubated for five days at the temperature of 25°C. Colonies were cultured multiple times in order to obtain a pure culture. After the initial biochemical tests, *Vibrio* and *Pseudoalteromonas* were selected for further analyses.

3.2. Bacterial Strains and Extract Preparation

Four strains of *Pseudoalteromonas* and *Vibrio* were identified at the genus level via biochemical assays (5), followed by the species level by the PCR amplification of the 16S rRNA gene (12). The properties of these bacteria are presented in Table 1.

The bacterial extract of each strain was obtained using the method proposed in (9). Briefly, a fresh culture of each purified isolate was grown in Marine broth in a shaker incubator at 180 rpm and the temperature of 25°C for five

days. The cell-free supernatant (CFS) was obtained by centrifugation followed by filtration (0.2 μm). Following that, the CFS was mixed with an equal volume of methanol and ethyl acetate (Merck, Germany) and shaken at 220 rpm for one hour. The solvent was removed under reduced pressure at the temperature of 25°C to collect the crude extracts (13), and the test was carried out in triplicate.

3.3. Pathogenic Strains

In this study, we used *S. aureus* (ATCC 25923) and *P. aeruginosa* (PTCC1430), which were provided by the Persian Gulf Marine Biotechnology Research Center in Bushehr, Iran. Both strains were grown at the temperature of 37°C in Luria-Bertani medium (LB; Sigma Aldrich) and Müller-Hinton agar (MHA; Difco).

3.4. Antimicrobial Activity

To evaluate the antibacterial activity of the extracts, the well-diffusion method was used in accordance with the protocol of the Clinical and Laboratory Standards Institute (9, 14). Briefly, diluted overnight cultures of *S. aureus* were prepared (OD600 = 0.1) and spread on the surface of agar plates. After the excess moisture was absorbed, 50 microliters of each extract was loaded into the wells, and the wells containing tetracycline and medium were used as the positive and negative controls, respectively. The process was repeated for *P. aeruginosa*, the plates were preserved at the temperature of 37°C for 24 hours. The test was carried out in triplicate.

3.5. Minimum Inhibitory Concentration and Minimum Bactericidal Concentration

To determine minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of each extract, the broth microdilution method was applied using 96-well microtiter plates (MTPs) in accordance with the protocol in (15) with slight modifications. We prepared an overnight culture of each pathogen (20 μL; OD600 = 0.1), 110 microliters of MHB, and 90 microliters of MHB supplemented with different concentrations of the extracts and the final concentrations of 75, 150, 300, 600, and 1,200 μg.mL⁻¹. In addition, tetracycline and medium were used as controls. The plates were preserved at the temperature of 37°C for 24 hours. The lowest concentration of each extract in which the extract inhibited visible growth was considered as the MIC. At the next stage, 100 microliters of the samples were collected from the wells with no turbidity and spread on a plate containing MHA.

Table 1. Status of Four Examined Sea Anemone Isolated Bacteria

Bacterial Extracts	Isolated Bacteria	GenBank Accession Number	Oxidase	Catalase	Triple Sugar Iron	Methyl Red	Voges-Proskauer	Simmons Citrate
E2	<i>Vibrio parahaemolyticus</i>	MN563125	+	+	K/A	+	-	+
E12	<i>Pseudoalteromonas gelatinilytica</i>	MN563128	+	+	A/A	+	+	+
E14	<i>Vibrio alginolyticus</i>	MN563124	+	+	K/A	-	-	+
E53	<i>Pseudoalteromonas piscicida</i>	MN563129	+	+	A/A	-	-	+

After 24 hours of incubation at the temperature of 37°C, the lowest concentration of the extract in which no colony appeared was considered as the MBC. The test was carried out in triplicate.

3.6. Growth Curve Analysis

Growth curve analysis was applied to confirm the antibacterial ability of the bacterial extracts against the two examined pathogens in accordance with the protocol in (16). In brief, the fresh culture of each pathogen (OD600 = 0.1), 110 microliters of MHB, and 90 microliters of each extract were dissolved in MHB at MIC and poured into the 96-well MTPs. Absorption values (OD600) were recorded for up to 24 hours at two-hour intervals and compared with the controls. The test was carried out in triplicate.

3.7. Antibiofilm Activity and Biofilm Inhibitory Concentration

For antibiofilm analysis, we used the method described in (17). An overnight culture of each pathogen (OD600 = 0.1), 110 microliters of the LB broth, and 90 microliters of the bacterial extract were prepared at the concentrations of MIC and sub-MICs. Notably, tetracycline and medium were used as controls. After incubation at the temperature of 37°C for 48 hours, the planktonic bacteria were removed by inverting the plate. Following that, the wells were gently washed three times with sterile phosphate-buffered saline (PBS), dried, and stained with 0.1% crystal violet (CV). At the next stage, the plate was inverted to remove the excess CV, rinsed with distilled water, and air-dried. In addition, 20% glacial acetic acid was poured into each well to dissolve the CV. The percentage of the biofilm inhibition of each extract was reported for each pathogen based on the absorbance of the wells at 570 nanometers and using the following formula:

$$\text{Percentage of Inhibition} = (\text{Control OD}_{570 \text{ nm}} - \text{Test OD}_{570 \text{ nm}}) / \text{Control OD}_{570 \text{ nm}} \times 100$$

The biofilm inhibitory concentration (BIC) was obtained as the lowest concentration indicating the significant reduction of the well absorbance compared to the controls. The test was carried out in triplicate.

3.8. Disruption of Preformed Biofilm

The ability of the extracts to disrupt the preformed biofilm was assessed based on the protocol in (18). An overnight culture of each pathogen (OD600 = 0.1) was inoculated into each well of the 96-well MTPs, supplemented with 200 microliters of the LB broth, and allowed to form biofilm at the temperature of 37°C. After 48 hours, the plates were inverted to remove the medium, rinsed with PBS, and dried at room temperature. Afterwards, the preformed biofilm in the wells were inoculated with 200 microliters of the bacterial extracts at their BICs. After incubation at the temperature of 37°C for 24 hours, the medium was discarded, and biofilm formation in each well was evaluated based on the CV staining using the previously described protocol. The wells containing the medium and sodium metaperiodate (40 mM) were considered as the negative and positive controls, respectively. The test was carried out in triplicate.

3.9. Active Compounds of the Extracts

To determine the nature of the extracts, they were exposed to proteinase K (Sigma-Aldrich) as a chemical treatment using the method proposed in (19). Proteinase K (1 mg/mL) was added to each of the 96 well plates containing the extracts (BIC) and incubated for one hour at the temperature of 37°C. Afterwards, the antibiofilm effects of the extracts were evaluated against both pathogens using the 96 MTPs and CV staining and compared to the controls. The samples that were not exposed to proteinase K were used as controls, and the test was carried out in triplicate.

3.10. Statistical Analysis

Data analysis was performed in SPSS version 16 (SPSS, Chicago, Illinois, USA). The tests were carried out in triplicate, and the values were expressed as mean and standard deviation (SD) of the replicates. One-way analysis of variance (ANOVA) and Duncan's test were applied for the comparison of the tests. In addition, the differences between the tests and controls were evaluated using Dunnett's test, and *t*-test was applied to compare the biofilm inhibition between the proteinase treatment and controls. Regarding the differences between the mean values, the significance level was considered less than 0.05.

4. Results

4.1. Antimicrobial Activity and Determination of MIC and MBC

Our findings regarding the antimicrobial activity of the four bacterial extracts indicated their growth inhibitory potential against *S. aureus* and *P. aeruginosa* (Table 2). Furthermore, the extracts could effectively inhibit the growth of *S. aureus* at the concentration of 600 µg/mL (MIC) and kill the bacteria at the concentration of 1,200 µg/mL (MBC) (Figure 1). The extracts also exhibited antibacterial activity against *P. aeruginosa* growth at the MIC of 300 µg/mL and MBC of 600 µg/mL, while the second extract showed significantly higher antibacterial activity against both pathogens at the lowest concentration ($P < 0.05$). On the other hand, the results of the growth curve analysis confirmed that at the determined MICs, the extracts could prevent the growth of both pathogens (Figure 2).

Table 2. Antibacterial Activity of Four Bacterial Extracts against *Staphylococcus aureus* and *Pseudomonas aeruginosa*

Bacterial Extracts	Inhibition Zone Against <i>S. aureus</i> , mm	Inhibition Zone Against <i>P. aeruginosa</i> , mm
E2	12.00 ± 1.40	14.20 ± 0.80
E12	9.20 ± 1.50	11.10 ± 0.80
E14	6.20 ± 0.70	9.30 ± 1.10
E53	8.10 ± 1.10	11.80 ± 1.40

4.2. Antibiofilm Activity and Disruption of the Preformed Biofilm

The calculated BIC of the extracts for *S. aureus* and *P. aeruginosa* was 600 and 300 µg/mL, respectively (Figure 3). According to the obtained results, the second extract exhibited significantly higher antibiofilm activity ($P < 0.05$)

against both pathogens. In addition, the four examined extracts disrupted the preformed biofilms of *S. aureus* and *P. aeruginosa* (Figure 4), while the second extract had the highest activity against the *S. aureus* biofilm ($P < 0.05$).

4.3. Active Compounds of the Extracts

The evaluation of the nature of the bacterial extracts indicated that the antibiofilm activity of the extracts against both pathogens decreased significantly after exposure to the protease ($P < 0.05$) (Figure 5). In fact, the test demonstrated that the four extracts had proteolysis activity, which played a key role in their antibiofilm activity.

5. Discussion

We initially prepared an extract from four examined strains of *Pseudoalteromonas* and *Vibrio*, which had metabolites that could inhibit the growth of *S. aureus* and *P. aeruginosa*. *Pseudoalteromonas* and *Vibrio* are abundant species in the marine habitat, and multiple strains have been isolated using commercially available complex media with the concentration of ions mimicking seawater (20). The genus *Pseudoalteromonas* currently consists of 41 species, 16 of which are known as the producers of antimicrobials. Various identified products have been reviewed in (21), and the compounds are reported to be alkaloids, polyketides, and peptides. In addition, *Pseudoalteromonas* have also been reported to produce digestive enzymes, which are capable of killing the surrounding bacteria through penetration into their cell wall (22). These bacteria are mostly detected in healthy animals and plants (23).

Vibrionaceae includes 126 species, and only six are reported to produce antimicrobial compounds (24). In the current research, *V. alginolyticus* and *V. parahaemolyticus* were observed to have a high antibacterial potential against *S. aureus* and *P. aeruginosa*. Although these strains are known as pathogens, they produce several bioactive compounds (e.g., indole, tetrodotoxin, and organic acids), which have anti-growth activity against various pathogens (25-27). Consistent with our findings, previous studies have confirmed the antibacterial potential of *Vibrio* and/or *Pseudoalteromonas* in sea slug (20), sea anemone (*S. haddoni*) (28), sea anemone (*Anemonia sulcata* and *Actinia equina*) (8), and coral (13) against various human pathogens.

The four examined isolates inhibited the biofilm formation of *S. aureus* and *P. aeruginosa*. Biofilm formation depends on several factors and their correlations; such exam-




Figure 1. Antibacterial activity of four selected SIB extracts at varying concentrations ($\mu\text{g.mL}^{-1}$) against the growth of *S. aureus* (S.A.) and *P. aeruginosa* (P.A.). Different letters shows significant difference (*, $P < 0.05$).




Figure 2. Growth curve of *S. aureus* (S.A) and *P. aeruginosa* (P.A) in the absence and presence of the selected extracts at MIC.

ples are growth, adhesions, extracellular matrix-binding proteins, biofilm architecture, and cell communication (quorum sensing [QS]) (29-31). Therefore, antibiofilm agents are expected to affect at least one of these factors. Since the four sea anemone extracts in the current research exhibited antibacterial activity, it was inferred that they

also were able to prevent the biofilm formation of the pathogens by inhibiting their growth. However, further investigations are required to clarify the other influential factors in biofilm formation. Our findings in this regard are in line with previous studies indicating the antibiofilm activity of the bacteria isolated from marine corals (9,19,32) and





Figure 3. Antibiofilm activity of four selected SIB extracts at varying concentrations ($\mu\text{g.mL}^{-1}$) against the biofilm formation of *S. aureus* (S.A) and *P. aeruginosa* (P.A). Different letters shows significant difference (*, $P < 0.05$).




Figure 4. Biofilm disruption activity of four selected SIB extracts at varying concentrations ($\mu\text{g.mL}^{-1}$) against the preformed biofilm of *S. aureus* (S.A) and *P. aeruginosa* (P.A). Different letters shows significant difference (*, $P < 0.05$).

sea slugs (20) against various pathogens.

The extracts could also disrupt the preformed biofilm of *S. aureus* and *P. aeruginosa* at their BICs. To date, data have been scarce regarding the disruptive effects of marine bacteria or their products on the preformed biofilm of pathogens. Similar to the blockage of biofilm formation, the dispersion of the preformed biofilm could be due to the secretion of matrix-degrading enzymes (33), cellular mortality (34), biofilm-degrading polysaccharides, organic acids (35), and compounds with biosurfactant and bioemulsifier activities (36) produced by marine bacteria.

We also evaluated the nature of the active compounds of the extracts, and a significant reduction was observed

in their antibiofilm activity following the enzymatic treatment, in which proteases were destroyed in the extracts. Proteins and peptides play a critical role in the biofilm structure (37). Therefore, the presence of proteolytic activity in an extract could be effective in the destruction of the biofilm network. Our findings in this regard demonstrated that the major active components of the extracts were proteases, which are essentially involved in the lysis of the proteins within the biofilm network. Consistent with our findings, previous studies have shown that proteases are the active compounds of bacterial extracts (19, 38). We could not identify all the active compounds of the extracts. However, it was observed that the antibiofilm potential of the extracts mainly relied on their proteolysis properties. Other effective compounds with fewer roles (e.g., anti-QS compounds) may also be detected, which require further investigation.

5.1. Conclusions

According to the results, the extracts derived from *Pseudoalteromonas* and *Vibrio* isolated from sea anemone (*S. haddoni*) had the potential to effectively control the growth and biofilm formation of *S. aureus* and *P. aeruginosa*. Therefore, the extracts should be explored further for the control of biofilm-associated infections.

Footnotes

Authors' Contribution: Conceptualization: Akram Sadat Naeemi and Seyed Amir Hossein Jalali. Methodology: Seyed Amir Hossein Jalali and Hojjatollah Zamani. Writing Original draft: Neda Fazeli. Investigation: Neda Fazeli. Supervision: Akram Sadat Naeemi and Seyed Amir Hossein Jalali. Validation: Akram Sadat Naeemi, Seyed Amir Hossein Jalali, and Hojjatollah Zamani. Reviewing and editing: Akram Sadat Naeemi, Seyed Amir Hossein Jalali, and Hojjatollah Zamani.

Conflict of Interests: There is no conflict of interest.

Ethical Approval: All applicable international and national guidelines for the care and use of animals were followed.

Funding/Support: There is no funding or support.




Figure 5. Effects of proteinase K treatment on the anti biofilm activity of four selected SIB extracts at MIC against the biofilm of *S. aureus* (S.A) and *P. aeruginosa* (P.A). Different letters shows significant difference (*, P < 0.05).

References

1. Dryden MS. Skin and soft tissue infection: microbiology and epidemiology. *Int J Antimicrob Agents*. 2009;34 Suppl 1:S2-7. doi: [10.1016/S0924-8579\(09\)70541-2](https://doi.org/10.1016/S0924-8579(09)70541-2). [PubMed: 19560670].
2. Figueiredo AMS, Ferreira FA, Beltrame CO, Cortes MF. The role of biofilms in persistent infections and factors involved in

- ica-independent biofilm development and gene regulation in *Staphylococcus aureus*. *Crit Rev Microbiol*. 2017;43(5):602-20. doi: [10.1080/1040841X.2017.1282941](https://doi.org/10.1080/1040841X.2017.1282941). [PubMed: 28581360].
3. Nikbin VS, Aslani MM, Sharafi Z, Hashemipour M, Shahcheraghi F, Ebrahimipour GH. Molecular identification and detection of virulence genes among *Pseudomonas aeruginosa* isolated from dif

- ferent infectious origins. *Iran J Microbiol.* 2012;**4**(3):118-23. [PubMed: 23066485]. [PubMed Central: PMC3465536].
4. Barr HL, Halliday N, Camara M, Barrett DA, Williams P, Forrester DL, et al. *Pseudomonas aeruginosa* quorum sensing molecules correlate with clinical status in cystic fibrosis. *Eur Respir J.* 2015;**46**(4):1046-54. doi: 10.1183/09031936.00225214. [PubMed: 26022946]. [PubMed Central: PMC4589431].
 5. Stempel N, Neidig A, Nusser M, Geffers R, Vieillard J, Lesouhaitier O, et al. Human host defense peptide LL-37 stimulates virulence factor production and adaptive resistance in *Pseudomonas aeruginosa*. *PLoS One.* 2013;**8**(12). e82240. doi: 10.1371/journal.pone.0082240. [PubMed: 24349231]. [PubMed Central: PMC3862677].
 6. Franci G, Falanga A, Galdiero S, Palomba L, Rai M, Morelli G, et al. Silver nanoparticles as potential antibacterial agents. *Molecules.* 2015;**20**(5):8856-74. doi: 10.3390/molecules20058856. [PubMed: 25993417]. [PubMed Central: PMC6272636].
 7. Debbab A, Aly AH, Lin WH, Proksch P. Bioactive compounds from marine bacteria and fungi. *Microb Biotechnol.* 2010;**3**(5):544-63. doi: 10.1111/j.1751-7915.2010.00179.x. [PubMed: 21255352]. [PubMed Central: PMC3815768].
 8. Leon-Palmero E, Joglar V, Alvarez PA, Martin-Platero A, Llamas I, Reche I. Diversity and antimicrobial potential in sea anemone and holothurian microbiomes. *PLoS One.* 2018;**13**(5). e0196178. doi: 10.1371/journal.pone.0196178. [PubMed: 29742123]. [PubMed Central: PMC5942802].
 9. Bakkiyaraj D, Pandian SK. In vitro and in vivo antibiofilm activity of a coral associated actinomycete against drug resistant *Staphylococcus aureus* biofilms. *Biofouling.* 2010;**26**(6):711-7. doi: 10.1080/08927014.2010.511200. [PubMed: 20706890].
 10. Balasubramanian S, Othman EM, Kampik D, Stopper H, Hentschel U, Ziebuhr W, et al. Marine Sponge-Derived Streptomyces sp. SBT343 Extract Inhibits Staphylococcal Biofilm Formation. *Front Microbiol.* 2017;**8**:236. doi: 10.3389/fmicb.2017.00236. [PubMed: 28261188]. [PubMed Central: PMC5311426].
 11. Chen YH, Kuo J, Sung PJ, Chang YC, Lu MC, Wong TY, et al. Isolation of marine bacteria with antimicrobial activities from cultured and field-collected soft corals. *World J Microbiol Biotechnol.* 2012;**28**(12):3269-79. doi: 10.1007/s11274-012-1138-7. [PubMed: 22872580].
 12. Kieser T, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA. *Practical streptomycetes genetics*. 291. John Innes Foundation Norwich; 2000.
 13. Nithyanand P, Pandian SK. Phylogenetic characterization of culturable bacterial diversity associated with the mucus and tissue of the coral *Acropora digitifera* from the Gulf of Mannar. *FEMS Microbiol Ecol.* 2009;**69**(3):384-94. doi: 10.1111/j.1574-6941.2009.00723.x. [PubMed: 19619231].
 14. CLSI. *Performance standards for antimicrobial susceptibility testing; twenty-fifth informational supplement*. Wayne, PA: Clinical and Laboratory Standards Institute; 2015. Report No.: M100-S25.
 15. Sambanthamoorthy K, Feng X, Patel R, Patel S, Paravavita C. Antimicrobial and antibiofilm potential of biosurfactants isolated from lactobacilli against multi-drug-resistant pathogens. *BMC Microbiol.* 2014;**14**:197. doi: 10.1186/1471-2180-14-197. [PubMed: 25124936]. [PubMed Central: PMC4236506].
 16. Medina-Martinez MS, Truchado P, Castro-Ibanez I, Allende A. Antimicrobial activity of hydroxytyrosol: a current controversy. *Biosci Biotechnol Biochem.* 2016;**80**(4):801-10. doi: 10.1080/09168451.2015.1116924. [PubMed: 26679750].
 17. Stepanovic S, Vukovic D, Hola V, Di Bonaventura G, Djukic S, Cirkovic I, et al. Quantification of biofilm in microtiter plates: overview of testing conditions and practical recommendations for assessment of biofilm production by staphylococci. *APMIS.* 2007;**115**(8):891-9. doi: 10.1111/j.1600-0463.2007.apm_630.x. [PubMed: 17696944].
 18. Dusane DH, Rajput JK, Kumar AR, Nancharaiyah YV, Venugopalan VP, Zinjarde SS. Disruption of fungal and bacterial biofilms by lauroyl glucose. *Lett Appl Microbiol.* 2008;**47**(5):374-9. doi: 10.1111/j.1472-765X.2008.02440.x. [PubMed: 19146524].
 19. Papa R, Selan L, Parrilli E, Tilotta M, Sannino F, Feller G, et al. Anti-Biofilm Activities from Marine Cold Adapted Bacteria Against Staphylococci and *Pseudomonas aeruginosa*. *Front Microbiol.* 2015;**6**:1333. doi: 10.3389/fmicb.2015.01333. [PubMed: 26696962]. [PubMed Central: PMC4677098].
 20. Bohringer N, Fisch KM, Schillo D, Bara R, Hertzler C, Grein F, et al. Antimicrobial Potential of Bacteria Associated with Marine Sea Slugs from North Sulawesi, Indonesia. *Front Microbiol.* 2017;**8**:1092. doi: 10.3389/fmicb.2017.01092. [PubMed: 28659904]. [PubMed Central: PMC5469899].
 21. Offret C, Desriac F, Le Chevalier P, Mounier J, Jegou C, Fleury Y. Spotlight on Antimicrobial Metabolites from the Marine Bacteria *Pseudoalteromonas*: Chemodiversity and Ecological Significance. *Mar Drugs.* 2016;**14**(7). doi: 10.3390/md14070129. [PubMed: 27399731]. [PubMed Central: PMC4962019].
 22. Richards GP, Watson MA, Needleman DS, Uknalis J, Boyd EF, Fay JP. Mechanisms for *Pseudoalteromonas piscicida*-Induced Killing of *Vibrios* and Other Bacterial Pathogens. *Appl Environ Microbiol.* 2017;**83**(11). doi: 10.1128/AEM.00175-17. [PubMed: 28363962]. [PubMed Central: PMC5440704].
 23. Choudhury JD, Pramanik A, Webster NS, Llewellyn LE, Gachhui R, Mukherjee J. The Pathogen of the Great Barrier Reef Sponge *Rhopaloeides odorabile* Is a New Strain of *Pseudoalteromonas agarivorans* Containing Abundant and Diverse Virulence-Related Genes. *Mar Biotechnol (NY).* 2015;**17**(4):463-78. doi: 10.1007/s10126-015-9627-y. [PubMed: 25837832].
 24. Mansson M, Gram L, Larsen TO. Production of bioactive secondary metabolites by marine vibronaceae. *Mar Drugs.* 2011;**9**(9):1440-68. doi: 10.3390/md9091440. [PubMed: 22131950]. [PubMed Central: PMC3225927].
 25. Fotso Fondja Yao CB, Al Zereini W, Fotso S, Anke H, Laatsch H. Aqabamycins A-G: novel nitro maleimides from a marine *Vibrio* species: II. Structure elucidation. *J Antibiot (Tokyo).* 2010;**63**(6):303-8. doi: 10.1038/ja.2010.35. [PubMed: 20431615].
 26. Nursyam H. Antibacterial Activity of Metabolites Products of *Vibrio Alginolyticus* Isolated from Sponge *Haliciona* sp. Against *Staphylococcus Aureus*. *Ital J Food Saf.* 2017;**6**(1):6237. doi: 10.4081/ijfs.2017.6237. [PubMed: 28299291]. [PubMed Central: PMC5337784].
 27. Veluri R, Oka I, Wagner-Dobler I, Laatsch H. New indole alkaloids from the North Sea bacterium *Vibrio parahaemolyticus* Bio249. *J Nat Prod.* 2003;**66**(11):1520-3. doi: 10.1021/np030288g. [PubMed: 14640534].
 28. Williams GP, Babu S, Ravikumar S, Kathiresan K, Prathap SA, Chinnappara S, et al. Antimicrobial activity of tissue and associated bacteria from benthic sea anemone *Stichodactyla haddoni* against microbial pathogens. *J Environ Biol.* 2007;**28**(4):789-93. [PubMed: 18405113].
 29. Raffa RB, Iannuzzo JR, Levine DR, Saeid KK, Schwartz RC, Susic NT, et al. Bacterial communication ("quorum sensing") via ligands and receptors: a novel pharmacologic target for the design of antibiotic drugs. *J Pharmacol Exp Ther.* 2005;**312**(2):417-23. doi: 10.1124/jpet.104.075150. [PubMed: 15528454].
 30. Stewart PS, Costerton JW. Antibiotic resistance of bacteria in biofilms. *Lancet.* 2001;**358**(9276):135-8. doi: 10.1016/s0140-6736(01)05321-1. [PubMed: 11463434].
 31. Swiatlo E, Champlin FR, Holman SC, Wilson WW, Watt JM. Contribution of choline-binding proteins to cell surface properties

- of *Streptococcus pneumoniae*. *Infect Immun*. 2002;**70**(1):412-5. doi: [10.1128/iai.70.1.412-415.2002](https://doi.org/10.1128/iai.70.1.412-415.2002). [PubMed: [11748210](https://pubmed.ncbi.nlm.nih.gov/11748210/)]. [PubMed Central: [PMC127640](https://pubmed.ncbi.nlm.nih.gov/PMC127640/)].
32. Rypien KL, Ward JR, Azam F. Antagonistic interactions among coral-associated bacteria. *Environ Microbiol*. 2010;**12**(1):28-39. doi: [10.1111/j.1462-2920.2009.02027.x](https://doi.org/10.1111/j.1462-2920.2009.02027.x). [PubMed: [19691500](https://pubmed.ncbi.nlm.nih.gov/19691500/)].
33. Saggu SK, Jha G, Mishra PC. Enzymatic Degradation of Biofilm by Metalloprotease From *Microbacterium* sp. SKS10. *Front Bioeng Biotechnol*. 2019;**7**:192. doi: [10.3389/fbioe.2019.00192](https://doi.org/10.3389/fbioe.2019.00192). [PubMed: [31448272](https://pubmed.ncbi.nlm.nih.gov/31448272/)]. [PubMed Central: [PMC6692633](https://pubmed.ncbi.nlm.nih.gov/PMC6692633/)].
34. Karatan E, Watnick P. Signals, regulatory networks, and materials that build and break bacterial biofilms. *Microbiol Mol Biol Rev*. 2009;**73**(2):310-47. doi: [10.1128/MMBR.00041-08](https://doi.org/10.1128/MMBR.00041-08). [PubMed: [19487730](https://pubmed.ncbi.nlm.nih.gov/19487730/)]. [PubMed Central: [PMC2698413](https://pubmed.ncbi.nlm.nih.gov/PMC2698413/)].
35. Yin Q, Liang J, Zhang W, Zhang L, Hu ZL, Zhang Y, et al. Butenolide, a Marine-Derived Broad-Spectrum Antibiofilm Agent Against Both Gram-Positive and Gram-Negative Pathogenic Bacteria. *Mar Biotechnol (NY)*. 2019;**21**(1):88-98. doi: [10.1007/s10126-018-9861-1](https://doi.org/10.1007/s10126-018-9861-1). [PubMed: [30612218](https://pubmed.ncbi.nlm.nih.gov/30612218/)]. [PubMed Central: [PMC6394721](https://pubmed.ncbi.nlm.nih.gov/PMC6394721/)].
36. Kiran GS, Sabarathnam B, Selvin J. Biofilm disruption potential of a glycolipid biosurfactant from marine *Brevibacterium casei*. *FEMS Immunol Med Microbiol*. 2010;**59**(3):432-8. doi: [10.1111/j.1574-695X.2010.00698.x](https://doi.org/10.1111/j.1574-695X.2010.00698.x). [PubMed: [20528933](https://pubmed.ncbi.nlm.nih.gov/20528933/)].
37. Pandey A, Naik M, Dubey SK. Hemolysin, Protease, and EPS Producing Pathogenic *Aeromonas hydrophila* Strain An4 Shows Antibacterial Activity against Marine Bacterial Fish Pathogens. *J Marine Biol*. 2010;**2010**:1-9. doi: [10.1155/2010/563205](https://doi.org/10.1155/2010/563205).
38. Wu S, Liu G, Zhang D, Li C, Sun C. Purification and biochemical characterization of an alkaline protease from marine bacteria *Pseudoalteromonas* sp. 129-1. *J Basic Microbiol*. 2015;**55**(12):1427-34. doi: [10.1002/jobm.201500327](https://doi.org/10.1002/jobm.201500327). [PubMed: [26213994](https://pubmed.ncbi.nlm.nih.gov/26213994/)].