Iranian Journal of Fuzzy Systems Vol. 4, No. 2, (2007) pp. 15-29

OPTIMIZATION OF LINEAR OBJECTIVE FUNCTION SUBJECT TO FUZZY RELATION INEQUALITIES CONSTRAINTS WITH MAX-AVERAGE COMPOSITION

E. SHIVANIAN, E. KHORRAM AND A. GHODOUSIAN

ABSTRACT. In this paper, the finitely many constraints of a fuzzy relation inequalities problem are studied and the linear objective function on the region defined by a fuzzy max-average operator is optimized. A new simplification technique which accelerates the resolution of the problem by removing the components having no effect on the solution process is given together with an algorithm and a numerical example to illustrate the steps of the problem resolution process.

1. Introduction

Fuzzy set theory was first introduced in 1965 by Zadeh [30]. The operations proposed by him are specified as the membership function of intersection and union of two fuzzy sets, and that of complement of the normalized fuzzy set. Since the resolution of fuzzy relation equations was proposed by Sanchez [25], fuzzy relation equations (FRE), fuzzy relation inequalities (FRI) and the problems related to them have been studied by many researchers (for instance, see Han et al [14], Hu [16], Loetamonphong and Fang [19, 20], Pedrycz [23], Wang [27, 28], Zhang et al [31] and Guo, Xia [11],and also[3, 6, 9, 10, 12, 15, 18, 22, 24, 29]). The max-min composition is commonly used when a system requires conservative solutions in the sense that the goodness of one value can not compensate for the badness of another [20]. Recent results in the literature, however, show that the min operator is not always the best choice for the intersection operation, but, in fact, the max-product composition provides results better or equivalent to the max-min composition in some applications [27].

The fundamental result for fuzzy relation equations with max-product composition goes back to Pedrycz [23] and recent study in this regard can be found in Bourk and Fisher [2], who extended the study of an inverse solution of a system of fuzzy relation equations with max-product composition and provided theoretical results for determining the complete solution sets as well as the conditions for the existence of resolutions. Their results show that such

Received: September 2006; Revised: January 2007; Accepted: April 2007

Key words and phrases: Linear objective function optimization, Fuzzy relation equations, Fuzzy relation inequalities.

complete solution sets can be characterized by one maximum solution and a number of minimal solutions. The monograph by Di Nola, Sessa, Pedrycz and Sanchez [7] contains a thorough discussion of this class of equations. The optimization problem with max-product and max-min compositions was first considered by Loetamonphong and Fang [19] and Fang and Li [8].

The optimization problem of max-min and max-product can be separated into two sub problems by separating the nonnegative and negative coefficients in the objective function. The sub- problem formed by the negative coefficients can be solved easily by obtaining the maximum solution of the feasible solutions set. On the other hand, the sub-problem formed by the nonnegative coefficients can be converted into a 0-1 integer programming problem. The associated 0-1 integer programming problem was solved by Fang and Li [8] using the branch and bound technique. Lu and Fang [21] proposed a genetic algorithm to solve nonlinear optimization problem with max-min composition and Guu and Wu [13] improved Fang and Li's method by providing an upper bound for the branch-and-bound procedure. For the optimization problem with max-product composition, Loetamonphong and Fang solved the corresponding 0-1 integer problem by reducing its size and by employing the branch and bound method. On other hand, Loetamonphong, Fang, and Young [20] extended the study of the problem of max-min composition to the solution with multi-objective functions. The application of (FRE) and (FRI) can be seen in many areas, for instance, fuzzy control, fuzzy decision making, fuzzy symptom, diagnosis, and especially fuzzy medical diagnosis [1,4,26].

In this paper, we generalize the linear optimization problem of the FRE with max-average operator [17] by considering the FRI constraints. This problem can be formulated as follows:

$$\min c' x$$
s.t. $A \bullet x \ge d^1$

$$B \bullet x \le d^2$$

$$x \in [0,1]^n$$
(1)

Where $A = (a_{ij})_{m \times n}$, $a_{ij} \in [0,1]$, $B = (b_{ij})_{l \times n}$, $b_{ij} \in [0,1]$, are fuzzy matrices, $d^1 = (d_i^1)_{m \times 1} \in [0,1]^m$, $d^2 = (d_i^2)_{l \times 1} \in [0,1]^l$, $x = (x_j)_{n \times 1} \in [0,1]^n$ are fuzzy vectors and, $c = (c_j)_{n \times 1} \in \mathbb{R}^n$, and "•" denotes the fuzzy max-average operator. Problem (1) can be rewritten as follows:

Optimization of Linear Objective Function Subject to Fuzzy Relation Inequalities Constraints 17 with Max-average Composition

$$\min c^{t} x$$
s.t. $a_{i} \bullet x \ge d_{i}^{1}, \quad i \in I^{1} = \{1, 2, ..., m\}$

$$b_{i} \bullet x \le d_{i}^{2} \quad i \in I^{2} = \{m + 1, m + 2, ..., m + 1\}$$

$$0 \le x_{j} \le 1, \quad j \in J = \{1, 2, ..., n\}$$
(2)

Where a_i and b_i are *i*'th row of the matrices *A* and *B*, respectively and the constraints are expressed by the max-average operator definition as;

$$a_{i} \bullet x = \max_{j \in J} \{ \frac{a_{ij} + x_{j}}{2} \} \ge d_{i}^{1} \qquad \forall i \in I^{1}$$

$$b_{i} \bullet x = \max_{j \in J} \{ \frac{b_{ij} + x_{j}}{2} \} \le d_{i}^{2} \qquad \forall i \in I^{2}$$
(3)

In section 2, the feasible solutions set of the problem (2) and its properties are studied. Moreover, necessary and sufficient conditions are given for the feasibility of the problem (2). In section 3, some simplification operations are presented to accelerate the resolution process. Finally, in section 4 an algorithm is introduced to solve the problem and a numerical example is given to illustrate the algorithm. Section 5 provides a conclusion.

2. The Characteristics of the Feasible Solution Set

Definition 2.1. Let

$${}^{i}S(A,d^{1}) = \{x \in [0,1]^{n} : a_{i} \bullet x \ge d_{i}^{1}\} \text{ for each } i \in I^{1},$$

$${}^{i}S(B,d^{2}) = \{x \in [0,1]^{n} : b_{i} \bullet x \le d_{i}^{2}\} \text{ for each } i \in I^{2},$$

$$S(A,d^{1}) = \bigcap_{i \in I^{1}} S(A,d^{1}) = \{x \in [0,1]^{n} : A \bullet x \ge d^{1}\},$$

$$S(B,d^{2}) = \bigcap_{i \in I^{2}} S(B,d^{2}) = \{x \in [0,1]^{n} : B \bullet x \le d^{2}\} \text{ and}$$

$$S(A,B,d^{1},d^{2}) = S(A,d^{1}) \cap S(B,d^{2}) = \{x \in [0,1]^{n} : A \bullet x \ge d^{1}, B \bullet x \le d^{2}\}.$$

Corollary 2.2. According to (3), we have $x \in {}^{i}S(A, d^{1})$ for each $i \in I^{1}$ iff there exists $j \in J$ such that $x_{j} \ge 2d_{i}^{1} - a_{ij}$. Similarly, $x \in {}^{i}S(B, d^{2})$ for each $i \in I^{2}$ iff $x_{j} \le 2d_{i}^{2} - b_{ij}$, $\forall j \in J$.

Lemma 2.3.

(a) $S(A,d^1) \neq \emptyset$ iff for each $i \in I^1$ there exists $j \in J$ such that $2d_i^1 - a_{ij} \leq 1$. (b) If $S(A,d^1) \neq \emptyset$ then $\overline{1} = [1,1,...,1]_{1 \times n}^t$ is the single maximum solution of $S(A,d^1)$.

www.SID.ir

Proof.

18

(a) Suppose $x \in S(A, d^1)$. Hence by Definition 2.1, $\forall i \in I^1$, $x \in {}^iS(A, d^1)$ and thus, by Corollary 2.2, for each $i \in I^1$, and some $j_i \in J$, we have $x_{j_i} \ge 2d_i^1 - a_{ij_i}$. Moreover, since $x \in [0,1]^n$, $x \in S(A, d^1)$, we have $2d_i^1 - a_{ij} \le 1 \forall i \in I^1$. Conversely, suppose there exists $j_i \in J$ such that $2d_i^1 - a_{ij} \le 1, \forall i \in I^1$.

Set $x = \overline{1} = [1,1,...,1]_{1 \times n}^t$. Since $x \in [0,1]^n$ and $x_{j_i} = 1 \ge 2d_i^1 - a_{ij_i}$, $\forall i \in I^1$, hence, by Corollary 2.2, $x \in S(A, d^1)$, $\forall i \in I^1$, and as a result, $x \in S(A, d^1)$.

(b) Proof of this part follows easily from part (a) and Corollary 2.2.

Lemma 2.4.

(a) $S(B,d^2) \neq \emptyset$ iff $2d_i^2 - b_{ij} \ge 0$, $\forall i \in I^2$ and $\forall j \in J$. (b) If $S(B,d^2) \neq \emptyset$ then $\overline{0} = [0,0,...,0]_{l_{ND}}^t$ is the single minimum solution of $S(B,d^2)$.

Proof. It is similar to the proof of the Lemma 2.3 by using the Corollary 2.2.

Theorem 2.5. (necessary condition) if $S(A, B, d^1, d^2) \neq \emptyset$ then (a) $\forall i \in I^1 \exists j \in J$ Such that $2d_i^1 - a_{ij} \leq 1$. (b) $2d_i^2 - b_{ij} \geq 0$, $\forall i \in I^2$ and $\forall j \in J$.

Proof. This theorem follows from Lemmas 2.3, 2.4 and Definition 2.1.

Definition 2.6. Set $\bar{x} = (\bar{x}_j)_{n \ge 1}$, where $\bar{x}_j = \max\{\min\{1, \min_{1 \le i < l} \{2d_i^2 - b_{ij}\}\}, 0\}$.

Lemma 2.7. If $S(B,d^2) \neq \emptyset$ then x is the single maximum solution of $S(B,d^2)$.

Proof. See the proof of the Theorem 2 in [17].

Corollary 2.8. If $S(B,d^2) \neq \emptyset$ then $S(B,d^2) = [\overline{0},\overline{x}]$.

Proof. We recall that, by Lemmas 2.3 and 2.4, $\overline{0}$ and \overline{x} are, respectively, the single minimum and the maximum solutions. Now, let $x \in [\overline{0}, \overline{x}]$. Then $x \in [0,1]^n$ and $x \le \overline{x}$. Thus, $b_i \bullet x \le b_i \bullet \overline{x} \le d_i^2$, $\forall i \in I^2$ that implies $x \in S(B, d^2)$. Conversely, let $x \in S(B, d^2)$. By Lemma 2.4 and Definition 2.1, $\overline{0} \le x$ and $x \in S(B, d^2)$, $\forall i \in I^2$. Next, by Corollary 2.2 we have $x_j \le 2d_i^2 - b_{ij}$, $\forall i \in I^2$ and $\forall j \in J$. Hence, $x_i \le \overline{x}_j$, $\forall j \in J$ i.e. $x \le \overline{x}$. It follows that $x \in [\overline{0}, \overline{x}]$.

Definition 2.9. Let $J_i = \{j \in J : 2d_i^1 - a_{ij} \le 1\}$, $\forall i \in I^1$. For each $j \in J_i$, we define $i_{x(j)} = (i_{x(j)_k})_{n \times 1}$ such that

www.SID.ir

Optimization of Linear Objective Function Subject to Fuzzy Relation Inequalities Constraints 19 with Max-average Composition

$$i_{x(j)_{k}} = \begin{cases} \max\{2d_{i}^{1} - a_{ij}, 0\} & k = j \\ 0 & k \neq j \end{cases}$$

Lemma 2.10. Assume $i \in I^1$ is a fixed number.

(a) For each $j \in J_i$, the vectors $i_{x(i)}$ are the minimal solutions of $S(A, d^1)$.

(b) If there exists $j \in J_i$ such that $2d_i^1 - a_{ij} \leq 0$ then $\overline{0}$ is the single minimum solution of $S(A, d^1)$.

Proof.

(a) Suppose $j \in J_i$ and $i \in I^1$, since $i_{x(j)_j} = \max\{2d_i^1 - a_{ij}, 0\}$ then, by Corollary 2.2, $i_{x(j)} \in S(A, d^1)$, $\forall i \in I^1$.

Now suppose that there exists $x \in S(A, d^1)$ such that $x < i_{x(j)}$. Then $x_j < 2d_i^1 - a_{ij}$ and, for $k \in J - \{j\}$, $x_k = 0$. Hence $x_j < 2d_i^1 - a_{ij}$, $\forall j \in J$ and so, by Corollary 2.2, $x \notin S(A, d^1)$, which is a contradiction.

(b) The proof of this part is similar to (a).

Corollary 2.11. If
$${}^{i}S(A,d^{1}) \neq \emptyset$$
 then ${}^{i}S(A,d^{1}) = \bigcup_{i=1}^{i} [i_{x(i)},\overline{1}]$, where $i \in I^{1}$.

Proof. We recall that ${}^{i}S(A, d^{1}) \neq \emptyset$ means that the vector $\overline{1}$ is the maximum solution and the vectors $i_{x(j)}$, $\forall j \in J_i$ are the minimal solutions in ${}^{i}S(A, d^{1})$ as a result of Lemmas 2.3, 2.4, 2.7 and 2.10, respectively. Now, let $x \in \bigcup_{i \in J_i} [i_{x(j)}, \overline{1}]$.

Hence, for some $j \in J_i$, $x \in [i_{x(i)}, \overline{1}]$.

Also, $x \in [0,1]^n$ and, by Definition 2.9, $x_j \ge i_{x(j)_j} = 2d_i^1 - a_{ij}$. Hence, by Corollary 2.2, $x \in S(A, d^1)$.

Conversely, let $x \in {}^{i}S(A, d^{1})$. Then, by Corollary 2.2, there exists $j' \in J$ such that $x_{j'} \ge 2d_i^{1} - a_{ij'}$. Since $x \in [0,1]^n$, so $2d_i^{1} - a_{ij'} \le 1$ and hence $j' \in J_i$. Therefore $i_{x(j')} \le x \le \overline{1}$, which implies that $x \in \bigcup_{i \in J_i} [i_{x(j)}, \overline{1}]$.

Definition 2.12. Suppose that $e = (e(1), e(2), \dots e(m)) \in J_1 \times J_2 \times \dots \times J_m$ such that $e(i) = j \in J_i$. $x(e)_j = \max\{\max_{i \in I_j^e} \{2d_i^1 - a_{ij}\}, 0\}$ if $I_j^e \neq \emptyset$ and $x(e)_j = 0$ if $I_j^e = \emptyset$, where $I_i^e = \{i \in I^1 : e(i) = j\}$. E. Shivanian, E. Khorram and A. Ghodousian

Corollary 2.13.

(a) If $2d_i^1 - a_{ij} \le 0$ for some $j \in J_i$, then we can remove the *i* th row of the matrix *A*. (b) If $j \notin J_i$, $\forall i \in I^1$ then we can omit the *j* th column of the matrix *A* for the purpose of finding x(e).

We recall that in part (a), by Definition 2.12 and part (b) of Lemma 2.10, the *i*th row of the matrix A has no effect in the calculation of the vectors x(e) for any $e \in J_1 = J_1 \times J_2 \times ... \times J_m$. Also, in part (b), before calculating the vectors x(e), $\forall e \in I_j$, we can remove the *j*th column of the matrix A using Definition 2.12.

Lemma 2.14. Let $S(A, d^{1}) \neq \emptyset$, then $S(A, d^{1}) = \bigcup_{X(e)} [x(e), \overline{1}]$, where

 $X(e) = \{x(e) : e \in I_J\}.$

Proof. If $S(A, d^1) \neq \emptyset$ then, ${}^iS(A, d^1) \neq \emptyset$, $\forall i \in I^1$. Hence, by Corollary 2.11 and Definitions 2.1, 2.6, 2.9 and 2.12, we have

$$S(A, d^{1}) = \bigcap_{i \in I^{1}} S(A, d^{1}) = \bigcap_{i \in I^{1}} \bigcup_{j \in J_{i}} [i_{x(j)}, \bar{1}]] = \bigcap_{i \in I^{1}} \bigcup_{e(i) \in J_{i}} [i_{x(e(i))}, \bar{1}]] = \bigcup_{e \in J_{I}} [\prod_{i \in I^{1}} [i_{x(e(i))}, \bar{1}]] = \bigcup_{e \in J_{I}} [x(e), \bar{1}] = \bigcup_{x(e)} [x(e), \bar{1}]$$

From Lemma 2.14 it is obvious that

$$S(A,d^{1}) = \bigcup_{X_{0}(e)} [x(e),\overline{1}] \text{ and } X_{0}(e) = S_{0}(A,d^{1}),$$

where $X_0(e)$ and $S_0(A, d^1)$ are the set of minimal solutions of X(e) and $S(A, d^1)$, respectively.

Theorem 2.15. If $S(A, B, d^1, d^2) \neq \emptyset$ then $S(A, B, d^1, d^2) = \bigcup_{X_0(e)} [x(e), \overline{x}]$.

Proof. Similar to the proof of the Lemma 2.14.

Corollary 2.16. (*Necessary and Sufficient Conditions*) Assume that $S(B,d^2) \neq \emptyset$. Then $S(A, B, d^1, d^2) \neq \emptyset$ iff $\overline{x} \in S(A, d^1)$ or, equivalently, $S(A, B, d^1, d^2) \neq \emptyset$ iff there exists $e \in J_1$ such that $x(e) \leq \overline{x}$.

Proof. the proof follows easily from Definition 2.1, Theorem 2.15 and Lemma 2.7.

3. Simplification Operations and the Resolution Algorithm

21

In order to solve the problem (2), it is initially converted into the two following sub-problems

 $\min c^{+'} x \qquad \min c^{-'} x$ s.t. $A \bullet x \ge d^1$ (4a) s.t. $A \bullet x \ge d^1$ (4b) $B \bullet x \le d^2$ $B \bullet x \le d^2$ $x \in [0,1]^n$ $x \in [0,1]^n$

Where, $c_{j}^{+} = \max(0, c_{j})$ and $c_{j}^{-} = \min(0, c_{j})$.

It is clear that x is an optimal solution of (4b) and (4a) achieves its optimal points at some $x(e) \in X_0(e)$. Once $x(e_0)$ optimizes (4a), we set $x^* = (x_i^*)_{n \ge 1}$ such that

$$x_{j}^{*} = \begin{cases} \overline{x}_{j} & , & c_{j} \leq 0 \\ x(e_{0})_{j} & , & c_{j} > 0 \end{cases}$$

Now the following lemma gives us an optimal point of the problem (2).

Lemma 3.1. x^* is an optimal solution of the problem (2).

Proof. See Theorem 2.1 in [11].

In order to calculate x^* , it is enough to find \overline{x} and $x(e_0)$. Although \overline{x} is easily obtained by Definition 2.6, $x(e_0)$ is not so, because, $X_0(e)$ is obtained by the pairwise comparison of X(e) members. Hence, having a complete set of $X_0(e)$ is time-consuming, especially when X(e) has several members. Simplification operations can hasten the resolution of the problem (4a). With the intention of simplification the vectors $e \in J_1$ are removed when x(e) is not optimal for (4a). One such operation is given in Corollary 2.13. Other operations are obtained by the following theorems.

Definition 3.2. Let $\overline{J}_i = \{j \in J_i : 2d_i^1 - a_{ij} \leq \overline{x}_j\}, \forall i \in I^1 \text{ where } \overline{x} \text{ is as in Definition 2.6.}$

Theorem 3.3. Let $S(B, d^2) \neq \emptyset$. Then $S(A, B, d^1, d^2) \neq \emptyset$ iff $\overline{J}_i \neq \emptyset, \forall i \in I^1$.

Proof. Suppose $S(A, B, d^1, d^2) \neq \emptyset$. Therefore, by Theorem 2.15, $\overline{x} \in S(A, B, d^1, d^2)$ and so we have $\overline{x} \in {}^iS(A, d^1)$, $\forall i \in I^1$. Thus, by Corollary 2.2, for each $i \in I^1$, there exists $j \in J$ such that $\overline{x}_j \ge 2d_i^1 - a_{ii}$. Consequently, $\overline{J}_i \neq \emptyset$, $\forall i \in I^1$. Conversely, suppose that $\forall i \in I^1$, $\overline{J}_i \neq \emptyset$. Then $\forall i \in I^1$ there exists $j \in J$ such that $\overline{x}_j \ge 2d_i^1 - a_{ij}$ and, by Corollary 2.2, it follows that $\overline{x} \in S(A, d^1)$, $\forall i \in I^1$. As a result, $\overline{x} \in S(A, d^1)$. This fact, together with Lemma 2.7, implies $\overline{x} \in S(A, B, d^1, d^2)$ and therefore, $S(A, B, d^1, d^2) \neq \emptyset$.

Theorem 3.4. Let $S(A, B, d^1, d^2) \neq \emptyset$, then $S(A, B, d^1, d^2) = \bigcup_{\overline{X}(e)} [x(e), \overline{x}]$, where $\overline{X}(e) = \{x(e) : e \in \overline{J}_1 = \overline{J}_1 \times \overline{J}_2 \times ... \times \overline{J}_m\}.$

Proof. By Theorem 2.15, it is sufficient to show that $x(e) \notin S(A, B, d^1, d^2)$ when $e \notin \overline{J}_I$. Suppose $e \notin \overline{J}_I$. Then there exist $i' \in I^1$ and $j' \in J_{i'}$ such that e(i') = j' and $2d_{i'}^1 - a_{ij'} > \overline{x}_{j'}$. Hence $i' \in I_{j'}^e$ and, by Definition 2.12, $x(e)_{j'} = \max\{\max_{i \in I_{j'}^e} \{2d_i^1 - a_{ij'}\}, 0\} \ge 2d_{i'}^1 - a_{ij'} > \overline{x}_j$. Therefore the inequality $x(e) \le \overline{x}$ does not hold and, as a consequence of Theorem 2.15, we obtain $x(e) \notin S(A, B, d^1, d^2)$.

We remark that as a result of Definition 3.2, $\overline{J}_i \subseteq J_i$, $\forall i \in I^1$. In other words, $\overline{X}(e) \subseteq X(e)$. Also, by Theorem 3.4, $S_0(A, B, d^1, d^2) \subseteq \overline{X}(e)$. Thus, the region of search can be reduced to the smaller set $S_0(A, B, d^1, d^2)$.

Definition 3.5. Let $J_i^* = \{ j \in \overline{J}_i : c_i^- \neq 0 \}, \forall i \in I^1.$

Theorem 3.6. Suppose $x(e_0)$ is an optimal solution of (4a) and $J_{i'}^* \neq \emptyset$ for some $i' \in I^1$, then there exists x(e') such that $e'(i') \in J_{i'}^*$ and x(e') is the optimal solution of (4a).

Proof. Suppose $J_{i'}^* \neq \emptyset$ for some $i' \in I^1$ and $e_0(i') = j'$. Define $e' \in \overline{J}_I$ such that $e'(i') = k \in J_{i'}^*$ and $e'(i) = e_0(i)$ for each $i \in I^1$ and $i \neq i'$. By means of Definition 2.12, we have

 $x(e_0)_{j'} = \max\{\max_{i \in I_{j'}^{e_0}} \{2d_i^1 - a_{ij'}\}, 0\} \ge \max\{\max_{\substack{i \in I_{j'}^{e_0}\\i \neq i'}} \{2d_i^1 - a_{ij'}\}, 0\} = x(e')_{j'} \text{ and }$

 $x(e_0)_j = x(e')_j$ for each $j \in J$ and $j \neq j',k$. Therefore, since $c_k^+ = 0$,

$$c^{+^{t}}x(e_{0}) = c^{+}_{j'}x(e_{0})_{j'} + \sum_{\substack{j \in J \\ j \neq j'}} c^{+}_{j}x(e_{0})_{j} \ge c^{+}_{j'}x(e')_{j'} + \sum_{\substack{j \in J \\ j \neq j'}} c^{+}_{j}x(e')_{j} = c^{+^{t}}x(e')$$

Hence x(e') is an optimal solution of (4a) and the proof is complete.

23

Corollary 3.7. If $J_i^* \neq \emptyset$ for some $i \in I^1$ then by omitting the *i* th row we reach a reduced problem for which each optimal solution is an optimal solution for the previous (main) problem.

Proof. Results from Theorem 3.6 and the fact that $c_i^+ = 0$ for each $j \in J_i^*$.

Definition 3.8. Let $j_1, j_2 \in J$, $c_{j_1} > 0$ and $c_{j_2} > 0$. We say that j_2 dominates j_1 iff

- (a) $j_1 \in \overline{J}_i$ implies $j_2 \in \overline{J}_i$, $\forall i \in I^1$.
- (b) For each $i \in I^1$ such that $j_1 \in \overline{J}_i$, $c_{j_1}(2d_i^1 a_{ij_1}) \ge c_{j_2}(2d_i^1 a_{ij_2})$.

Theorem 3.9. Suppose $x(e_0)$ is optimal in (4a) and, for $j_1, j_2 \in J$, j_2 dominates j_1 . Then there exists x(e') such that $I_{j_1}^{e'} = \emptyset$ and x(e') is an optimal solution of (4a).

Proof. Define $e' = (e'(i))_{m \times 1}$ such that

$$e'(i) = \begin{cases} e_0(i) & i \notin I_{j_1}^{e_0} \\ j_2 & i \in I_{j_1}^{e_0} \end{cases}$$

It is obvious that $I_{j_1}^{e'} = \emptyset$ and so $x(e')_{j_1} = 0$. Also, $x(e_0)_j = x(e')_j$ for each $j \in J$ and $j \neq j_1, j_2, x(e')_{j_2} = 2d_{i_0}^1 - a_{i_0j_2}$. If $i_0 \notin I_{j_1}^{e_0}$ then:

$$x(e_0)_{j_2} = x(e')_{j_2} = 2d_{i_0}^1 - a_{i_0j_2},$$

So we have:

$$c^{+'}x(e_0) = c^{+}_{j_1}x(e_0)_{j_1} + \sum_{\substack{j \in J \\ j \neq j_1}} c^{+}_jx(e_0)_j \ge \sum_{\substack{j \in J \\ j \neq j_1}} c^{+}_jx(e')_j = c^{+'}x(e')$$

and the result follows.

Now suppose $i_0 \in I_{j_1}^{e_0}$. We show that $c^{+t}x(e_0) \ge c^{+t}x(e')$. Let $x(e_0)_{j_2} = 2d_i^1 - a_{ij_2}$. Then, by part (a) of the Corollary 2.13 and Definition 3.8, $c_{j_2}^+x(e_0)_{j_2} \ge 0$ and since

$$c^{+'}x(e_0) = c_{j_1}^+x(e_0)_{j_1} + c_{j_2}^+x(e_0)_{j_2} + \sum_{j \neq j_1, j_2} c_j^+x(e_0)_j$$

and

$$c^{+'}x(e') = c^{+}_{j_2}x(e')_{j_2} + \sum_{j \neq j_1, j_2} c^{+}_{j}x(e')_{j_2}$$

it is sufficient to show that $c_{j_1}^+ x(e_0)_{j_1} \ge c_{j_2}^+ x(e')_{j_2}$. By Definition 2.12, set $x(e_0)_{j_1} = 2d_{i'}^1 - a_{i'_1}$.

Since j_2 dominates j_1 , we have $c_{j_1}^+ (2d_{i'}^1 - a_{i'j_1}) \ge c_{j_2}^+ (2d_{i_0}^1 - a_{i_0j_2})$. i.e. $c_{j_1}^+ x(e_0)_{j_1} \ge c_{j_2}^+ x(e')_{j_2}$ once $i_0 = i'$. Otherwise, suppose $i_0 \ne i'$. Since $i_0 \in I_{j_1}^{e_0}$ and j_2 dominates j_1 , hence

$$c_{j_1}^+ (2d_{i_0}^1 - a_{i_0j_1}) \ge c_{j_2}^+ (2d_{i_0}^1 - a_{i_0j_2})$$

Also, by Definition 2.12 we have $x(e_0)_{j_1} = \max\{\max_{i \in I_{j_1}^{e_0}} \{2d_i^1 - a_{ij_1}\}, 0\} = 2d_{i'}^1 - a_{ij_1}$ which implies that

$$2d_{i'}^1 - a_{i'j_1} \ge 2d_i^1 - a_{ij_1}$$
 , $\forall i \in I_{j_1}^{e_0}$

Therefore

24

$$c_{j_1}^+(2d_{i'}^1-a_{ij_1}) \ge c_{j_1}^+(2d_{i_0}^1-a_{i_0j_1}) \ge c_{j_2}^+(2d_{i_0}^1-a_{i_0j_2})$$

It follows that $c_{j_1}^+ x(e_0)_{j_1} \ge c_{j_2}^+ x(e')_{j_2}$. Hence $c^{+'} x(e_0) \ge c^{+'} x(e')$ and the proof is complete.

Corollary 3.10. If j_2 dominates j_1 for each $j_1, j_2 \in J$, then, by omitting the j_1 th column we reach a reduced problem for which each optimal solution is an optimal solution for the previous (main) problem.

4. An Algorithm for Finding an Optimal Solution and Example

Definition 4.1. Consider problem (1). We call $\overline{A} = (\overline{a}_{ij})_{m \times n}$ the characteristic matrix of the matrix A, where $\overline{a}_{ij} = 2d_i^1 - a_{ij}$ for each $i \in I^1$ and $j \in J$. The characteristic matrix $\overline{B} = (\overline{b}_{ij})_{l \times n}$ is similarly defined.

Algorithm: Given problem (2),

- 1- Find the matrices \overline{A} and \overline{B} by Definition 4.1.
- 2- Calculate x from B by Definition 2.6.
- 3- If there exists $i \in I^1$ such that $\overline{a}_{ij} > \overline{x}_j$, $\forall j \in J$ then stop. Problem 2 is infeasible (see Theorem 2.5).
- 4- If there exists $i \in I^2$ and $j \in J$ such that $\overline{b}_{ij} < 0$ then stop. Problem 2 is infeasible (see Theorem 2.5).

25

- 5- If there exists $i \in I^1$ and $j \in J$ such that $\overline{a_{ij}} \leq 0$ then remove the *i*th row of the matrix \overline{A} (see the part (a) of the Corollary 2.13).
- 6- If there exists $j' \in J$ such that $\overline{a}_{ij'} > \overline{x}_{j'}$, $\forall i \in I^1$ then remove the j' th column of the matrix \overline{A} (see Theorem 3.4) and set $x(e_0)_{j'} = 0$ if $c_{j'} > 0$.
- 7- For each $i \in I^1$, if $J_i^* \neq \emptyset$ then remove the *i*th row of the matrix *A* (see Corollary 3.7).
- 8- If $c_i < 0$, remove column $j \in J$ from \overline{A} .
- 9- If j_2 dominates j_1 then remove the column j_1 from \overline{A} , $\forall j_1, j_2 \in J$ (see Corollary 3.10) and set $x(e_0)_{j_1} = 0$.
- 10- Let $J_i^{new} = \{j \in \overline{J}_i : \overline{a}_{ij} \leq \overline{x}_j\}$ and $J_I^{new} = J_1^{new} \times J_2^{new} \times ... \times J_m^{new}$. First obtain the vectors x(e), $\forall e \in J_I^{new}$ of \overline{A} using Definition 2.12, and then $x(e_0)$ by pairwise comparison between the vectors x(e).
- 11- Find x^* by Lemma 3.1.

Numerical Example:

Consider the problem below:

min $2x_1 - x_2 + 3x_3 + 2.5x_4 - x_5 - 6x_6 + 3x_7 + 2x_8$

0	0.16	0.37	0.95	0.17	0.07	0.77	0.14]	0.4	
0.08	0.51	0.26	0.1	0.3	0.4	0.3	0.35		0.5	
0.99	0.59	0.28	0.34	0.34	0.74	0.19	0.21		0.57	7
0.83	0.75	0.25	0.35	0.2	0.5	0.2	0.2	$ \bullet x \ge$	0.6	
0.73	0.84	0.94	0.44	0.54	0.84	0.99	0.44		0.72	2
0.37	0.7	0.55	0.4	0.2	0.2	0.73	0.24		0.6	
[0	0.94	0.69	0.49	0.5	0.51	0.87	0.43	Γ	0.7	
0.02	2 1	0.82	0.59	0.89	0.76	0.74	0.52		0.76	
0.1	0.73	0.4	0.37	0.45	0.7	0.69	0.2		0.6	
0.1	0.1	0.3	0.2	0.25	0.49	0.58	0.01	L	0.5	
$x \in [$	$[0,1]^n$									

Step 1. The matrices, \overline{A} , \overline{B} are as follows:

	0]	.8	0.64	0.43	-0.15	5 0.63	3 0.73	3 0.0	3 0.6	56]
	0.	92	0.49	0.74	0.9	0.7	0.6	0.7	0.6	55
<u> </u>	0.	15	0.55	0.86	0.8	0.8	0.4	0.9	5 0.9	93
A –	0.	37	0.45	0.95	0.85	1	0.7	1	1	
	0.	71	0.6	0.5	1	0.9	0.6	0.4	5 1	
	0.	83	0.5	0.65	0.8	1	1	0.4′	7 0.9	96]
		1.4	0.46	0.71	0.91	0.9	0.89	0.53	0.97	
		1.5	0.52	0.7	0.93	0.63	0.76	0.78	1	
D		1.1	0.47	0.8	0.83	0.75	0.5	0.51	1	
		0.9	0.9	0.7	0.8	0.75	0.51	0.42	0.99	

Step 2. $\bar{x} = \begin{bmatrix} 0.9 & 0.46 & 0.7 & 0.8 & 0.63 & 0.5 & 0.42 & 0.97 \end{bmatrix}$

Step 3. There is no $i \in I^1$ such that $\overline{a}_{ij} > \overline{x}_j$, $\forall j \in J$ therefore we can go to step 4.

Step 4. There is no $i \in I^2$ and $j \in J$ such that $\overline{b}_{ij} < 0$ hence we can perform step5.

Step 5. Since $\overline{a}_{14} < 0$, hence the first row of the matrix \overline{A} is removed and \overline{A} becomes

	0.92	0.49	0.74	0.9	0.7	0.6	0.7	0.65
	0.15	0.55	0.86	0.8	0.8	0.4	0.95	0.93
$\overline{A} =$	0.37	0.45	0.95	0.85	1	0.7	1	1
	0.71	0.6	0.5	1	0.9	0.6	0.45	1
	0.83	0.5	0.65	0.8	1	1	0.47	0.96

Step 6. The fifth and seventh columns are removed and we have $x(e_0)_7 = 0$. Also, matrix \overline{A} becomes

Optimization of Linear Objective Function Subject to Fuzzy Relation Inequalities Constraints 27 with Max-average Composition

	0.92	0.49	0.74	0.9	0.6	0.65
	0.15	0.55	0.86	0.8	0.4	0.93
$\overline{A} =$	0.37	0.45	0.95	0.85	0.7	1
	0.71	0.6	0.5	1	0.6	1
	0.83	0.5	0.65	0.8	1	0.96

Step 7. Since $J_3^* \neq \emptyset, J_4^* \neq \emptyset$, we can delete the second and third row. We get

	0.92	0.49	0.74	0.9	0.6	0.65
$\overline{A} =$	0.71	0.6	0.5	1	0.6	1
	0.83	0.5	0.65	0.8	1	0.96

Step 8. Since $c_2, c_6 < 0$ then, we can remove the second and fifth columns and we get

$$\overline{A} = \begin{bmatrix} 0.92 & 0.74 & 0.9 & 0.65 \\ 0.71 & 0.5 & 1 & 1 \\ 0.83 & 0.5 & 0.8 & 0.96 \end{bmatrix}$$

Step 9. In the above matrix, the first and fourth columns dominate the second and third respectively. After removing the second and third columns, matrix \overline{A} becomes

$$\overline{A} = \begin{bmatrix} 0.92 & 0.65 \\ 0.71 & 1 \\ 0.83 & 0.96 \end{bmatrix}$$

And we have $x(e_0)_3 = x(e_0)_4 = 0$.

Step 11. Since $x(e_1)$ optimizes the problem with objective function $c^{+'}x$ then $x(e_1) = x(e_0)$ and also $x^* = (0.83, 0.46, 0, 0, 0.63, 0.5, 0, 0.65)$.

www.SID.ir

5. Conclusion

In this paper we studied the linear optimization problem with fuzzy relational inequalities constraints defined by max-average operator. First, we discuss the feasibility region characterization and then, introduce a new simplification technique to solve the usual difficulty of finding the minimal solutions that optimize the problem with objective function $c^{+t}x$. An algorithm together with simplification operations to accelerate the problem resolution is also presented together with an illustrative numerical example.

Acknowledgments. The authors are grateful to referees for their valuable comments and constructive suggestions.

References

- [1] K. -P. Adlassnig, *Fuzzy set theory in medical diagnosis*, IEEE Trans. Systems Man Cybernet., **16** (1986), 260-265.
- [2] M. M. Brouke and D. G. Fisher, *Solution algorithms for fuzzy relation equations with max-product composition*, Fuzzy Sets and Systems, **94** (1998), 61-69.
- [3] E. Czogala and W. Pedrycz, *Control problems in fuzzy systems*, Fuzzy Sets and Systems, 7 (1982), 257-273.
- [4] E. Czogala and W. Predrycz, On identification in fuzzy systems and its applications in control problem, Fuzzy Sets and Systems, 6, 73-83.
- [5] E. Czogala, J. Drewniak and W. Pedrycz, *Fuzzy relation equations on a finite set*, Fuzzy Sets and Systems, 7 (1982), 89-101.
- [6] A. Di Nola, *Relational equations in totally ordered lattices and their complete resolution*, J. Math. Appl., **107** (1985), 148-155.
- [7] A. Di Nola, S. Sessa, W. Pedrycz and E. Sanchez, *Fuzzy relational equations and their applications in knowledge engineering*, Dordrecht: Kluwer Academic Press, 1989.
- [8] S. -C. Fang and G. Li, Solving fuzzy relations equations with a linear objective function, Fuzzy Sets and Systems, 103 (1999), 107-13.
- [9] S. -C. Fang and S. Puthenpura, *Linear optimization and extensions: theory and algorithm*, Prentice-Hall, Englewood Cliffs, NJ, 1993.
- [10] S. Z. Guo, P. Z. Wang, A. Di Nola and S. Sessa, Further contributions to the study of finite fuzzyrelation equations, Fuzzy Sets and Systems, 26 (1988), 93-104.
- [11] F. -F. Guo and Z. -Q. Xia, An algorithm for solving optimization Problems with one linear objective function and finitely many constraints of fuzzy relation inequalities, Fuzzy Optimization and Decision Making, 5 (2006), 33-47.
- [12] M. M. Gupta and J. Qi, *Design of fuzzy logic controllers based on generalized t-operators*, Fuzzy Sets and Systems, **40** (1991), 473-486.
- [13] M. Guu and Y. K. Wu, *Minimizing a linear objective function with fuzzy relation equation constraints*, Fuzzy Optimization and Decision Making, **12** (2002), 1568-4539.
- [14] S. S. Z. Han, A. H. Song, and T. Sekiguchi, *Fuzzy inequality relation system identification via sign matrix method*, Proceeding of 1995 IEEE International Conference, 3 (1995), 1375-1382.

29

- [15] M. Higashi and G. J. Klir, *Resolution of finite fuzzy relation equations*, Fuzzy Sets and Systems, 13 (1984), 65-82.
- [16] C. F. Hu, Generalized Variational inequalities with fuzzy relation, Journal of Computationaland Applied Mathematics, 146 (1998), 198-203.
- [17] E. Khorram and A.Ghodousian, *Linear objective function optimization with fuzzy relation constraints regarding max-av composition*, Applied Mathematics and Computation, **173** (2006), 827-886.
- [18] G. Li and S. -C. Fang, *Resolution of finite fuzzy resolution equations*, Report No. 322, North Carolina State University, Raleigh, NC, May 1996.
- [19] J. Loetamonphong and S. -C. Fang, Optimization of fuzzy relation equations with maxproduct composition, Fuzzy Sets and Systems, 118 (2001), 509-517.
- [20] J. Loetamonphong, S. -C. Fang and R.E. Young, *Multi-objective optimization problems with fuzzy relation equation constraints*, Fuzzy Sets and Systems, **127** (2002), 141-164.
- [21] J. Lu and S. -C. Fang, Solving nonlinear optimization problems with fuzzy relation equation constraints, Fuzzy Sets and Systems, **119** (2001), 1-20.
- [22] W. Pedrycz, On Generalized fuzzy relational equations and their applications, Journal of Mathematical Analysis and Applications, 107 (1985), 520-536.
- [23] W. Pedrycz, Proceeding in relational structures: fuzzy relational equations, Fuzzy Sets and Systems, 40 (1991), 77-106.
- [24] M. Prevot, Algorithm for the solution of fuzzy relations, Fuzzy Sets and Systems, 5 (1985), 319-322.
- [25] E. Sanchez, Resolution of composite fuzzy relation equations, Inform. Control, 30 (1976), 38-48.
- [26] W. B. Vasantha Kandasamy and F. Smarandache, *Fuzzy relational maps and neutrosophic relational maps*, Hexis Church Rock 2004 (chapter two).
- [27] P. Z. Wang, How many lower solutions of finite fuzzy relation equations, Fuzzy Mathematics (Chinese), 4 (1984), 67-73.
- [28] P. Z. Wang, Lattecized linear programming and fuzzy relaion inequalies, Journal of Mathematical Analysis and Applications, 159 (1991), 72-87.
- [29] W. L. Winston, *Introduction to mathematical programming: application and algorithms*, Duxbury Press, Belmont, CA, 1995.
- [30] L. A. Zadeh, Fuzzy sets, Inform. Control, 8 (1965), 338-353.
- [31] H. T. Zhang, H. M. Dong and R. H. Ren, *Programming problem with fuzzy relation inequality constraints*, Journal of Liaoning Noramal University, **3** (2003), 231-233.

ELYAS SHIVANIAN*, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, AMIRKABIR UNIVERSITY OF TECHNOLOGY, TEHRAN 15914, IRAN

E-mail address: eshivanian@gmail.com

ESMAILE KHORRAM, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, AMIRKABIR UNIVERSITY OF TECHNOLOGY, TEHRAN 15914, IRAN

E-mail address: eskor@aut.ac.ir

Amin Ghodousian, Faculty of Mathematics and Computer Science, Amirkabir University of Technology, Tehran 15914, Iran

* CORRESPONDING AUTHOR