

UNIVERSAL TRIPLE I METHOD FOR FUZZY REASONING AND FUZZY CONTROLLER

Y. M. TANG AND F. J. REN

ABSTRACT. As a generalization of the triple I method, the universal triple I method is investigated from the viewpoints of both fuzzy reasoning and fuzzy controller. The universal triple I principle is put forward, which improves the previous triple I principle. Then, unified form of universal triple I method is established based on the (0,1)-implication or R-implication. Moreover, the reversibility property of universal triple I method is analyzed from expansion, reduction and other type operators, which demonstrate that its reversibility property seems fine, especially for the case employing the (0,1)-implication. Lastly, we analyze the response ability of fuzzy controllers based on universal triple I method, then the practicability of triple I method is improved.

1. Introduction

Fuzzy reasoning plays a significant role in fuzzy control, artificial intelligence, affective computing, image processing and complex system (see [1, 7, 17, 18, 26, 28]). It is well-known that the most basic problem of fuzzy reasoning is fuzzy modus ponens (FMP) as follows:

FMP: For a given rule “If x is A then y is B ” and input “ x is A^* ”,
to compute B^* (output), (1)

where $A, A^* \in F(X)$, $B, B^* \in F(Y)$ ($F(X)$, $F(Y)$ respectively denote the set of all fuzzy subsets on X and Y). As for the FMP problem (1), the broadly used method in fuzzy control is the famous CRI (Compositional Rule of Inference) method proposed by Zadeh (see [4, 8, 20, 37]). The CRI solution is as follows:

$$B^*(y) = \sup_{x \in X} \{A^*(x) \wedge (A(x) \rightarrow B(y))\} \quad (y \in Y) \quad (2)$$

where \rightarrow is an implication. In 1999, Wang pointed out that the CRI method had some blemishes (see [33]). To improve the CRI method Wang put forward the triple

Received: December 2011; Revised: June 2012 and August 2012; Accepted: December 2012

Key words and phrases: Fuzzy reasoning, Fuzzy controller, Interpolation mechanism, Reversibility property, CRI method.

This work was supported by the National High-Tech Research & Development Program of China (863 Program, No. 2012AA011103), the National Natural Science Foundation of China (No. 61203077; No. 61070124; No. 61105076; No. 60672120; No. 41076120; No. 61271381), the Science and Technological Fund of Anhui Province for Outstanding Youth (No. 10040606Y09), the China Postdoctoral Science Foundation (No. 2012M521218), and the Fundamental Research Funds for the Central Universities (No. 2010HGXJ0017; No. 2012HGXJC0011; No. 2012HGCX0001; No. 2012HGBZ0639).

I method (see [33]), whose idea is to seek the optimal $B^* \in F(Y)$ such that

$$(A(x) \rightarrow B(y)) \rightarrow (A^*(x) \rightarrow B^*(y)) \quad (3)$$

takes the maximum for any $x \in X, y \in Y$. Following that, lots of scholars carried through a series of researches related to the triple I method (see [5, 14, 34, 38]), demonstrating that the triple I method possesses many advantages such as its logic basis, excellent reversibility property, and the property of pointwise optimization (see [11, 24, 27, 36]), which is excellent from the viewpoint of logic.

It is pointed out in [10,12] that many usual fuzzy controllers (including the one based on the CRI method) can be regarded as an interpolation method. Following that, the interpolation mechanism (or more general terminology called “response ability”) of fuzzy controllers based on the triple I method or CRI method, have attracted rapidly growing interests (see [6, 14, 15, 25, 29]). Li et al. discussed the response ability of 23 fuzzy controllers based on the CRI method, where got the fact that 12 fuzzy controllers can be used (see [14]). In [6], Hou et al. analyzed the response ability of 51 fuzzy controllers based on the triple I method, and achieved only 2 usable fuzzy controllers. Li et al. discussed the response ability of 4 fuzzy controllers based on the triple I method, and obtained 2 usable fuzzy controllers (see [15]). We drew the conclusion that 2 fuzzy controllers are practicable in 11 fuzzy controllers via the triple I method, while 4 fuzzy controllers are usable in 11 ones via the CRI method, which are constructed by the same 11 implications (see [29]). As a result, there are very few usable fuzzy controllers based on the triple I method, implying that the response ability and practicability of the triple I method are imperfect (from the viewpoint of fuzzy controllers).

To solve this problem, an important way is to improve the triple I method. Li pointed out in [11] the fact that the CRI method is a special case of the triple I method only if three implications in (3) are different. In detail, the CRI method can be regarded as the triple I method where (3) is changed into

$$(A(x) \rightarrow B(y)) \rightarrow_2 (A^*(x) \rightarrow_2 B^*(y)),$$

where \rightarrow_2 takes the Mamdani operator I_M .

Enlightened by this idea, we can let the latter two implications be the same and the first one unlimited, that is, generalize (3) to:

$$(A(x) \rightarrow_1 B(y)) \rightarrow_2 (A^*(x) \rightarrow_2 B^*(y)), \quad (4)$$

where \rightarrow_1 and \rightarrow_2 (respectively called the first implication and second implication in the sequel) can take different implications, and the triple I method derived from (4) is called the differently implicational universal triple I method of (1, 2, 2) type (universal triple I method for short). In [30, 31], we have already proposed and discussed the universal triple I method with some preliminary results.

In the theory of fuzzy reasoning, there are no acknowledged standards to judge whether a fuzzy reasoning method is excellent; but the reversibility property of fuzzy reasoning method is a basic demand (embodying the compatibility), which is recognized by a lot of scholars (see [5, 23, 24, 27]). The reversibility property of universal tripe I method is not researched. Therefore, one aim of this paper is to

analyze the reversibility property of universal triple I method. Following that, we shall inspect the universal triple I method from the viewpoint of fuzzy controllers, i.e., the response ability of fuzzy controllers based on universal triple I methods.

The rest of this paper is organized as follows. Section 2 is the preliminaries. In sections 3 and 4, the universal triple I method is investigated from the viewpoint of fuzzy reasoning. The solutions of universal triple I method are discussed, then based on them, the reversibility property of universal triple I method is analyzed. In section 5, the universal triple I method is investigated from the viewpoint of fuzzy controllers. The response ability of single-input single-output (SISO) fuzzy controllers and double-input single-output (DISO) fuzzy controllers based on the universal triple I method are respectively researched. Section 6 provides several discussions of the universal triple I method. Section 7 concludes this paper.

2. Preliminaries

Definition 2.1. An implication on $[0, 1]$ is a function $I : [0, 1]^2 \rightarrow [0, 1]$ satisfying (C1) $I(0, 0) = I(0, 1) = I(1, 1) = 1, I(1, 0) = 0$. $I(a, b)$ is also written as $a \rightarrow b$ ($a, b \in [0, 1]$).

Definition 2.2. A function $T : [0, 1]^2 \rightarrow [0, 1]$ is called a t-norm if T is associative commutative, increasing and satisfies $T(1, a) = a$ ($a \in [0, 1]$).

Definition 2.3. A function $S : [0, 1]^2 \rightarrow [0, 1]$ is called a t-conorm if S is associative commutative, increasing and satisfies $S(0, a) = a$ ($a \in [0, 1]$).

Definition 2.4. A decreasing function $N : [0, 1] \rightarrow [0, 1]$ is called a strong negation, if $N(0) = 1, N(1) = 0$ and $N(N(a)) = a$ ($a \in [0, 1]$).

Definition 2.5. [18] An implication \rightarrow is said to be an R-implication if there exist a left-continuous t-norm T such that (where \vee denotes supremum)

$$a \rightarrow b = \vee \{x \in [0, 1] \mid T(a, x) \leq b\}, \quad a, b \in [0, 1]. \quad (5)$$

Definition 2.6. Let T, \rightarrow be two $[0, 1]^2 \rightarrow [0, 1]$ functions, (T, \rightarrow) is called a residual pair or, T and \rightarrow are residual to each other, if the residual condition holds, i.e., (iff denotes "if and only if")

$$T(a, b) \leq c \text{ iff } b \leq a \rightarrow c, \quad a, b, c \in [0, 1]. \quad (6)$$

Proposition 2.7. [3] Let T be a t-norm, then the following are equivalent: (i) T is left-continuous; (ii) T and \rightarrow form a residual pair, where \rightarrow is from (5).

Definition 2.8. An implication \rightarrow is said to be an S-implication if there exist a t-conorm S and a strong negation N such that

$$a \rightarrow b = S(N(a), b), \quad a, b \in [0, 1].$$

Definition 2.9. An implication \rightarrow is called a QL-implication if there exist a t-norm T , a t-conorm S and a strong negation N such that

$$a \rightarrow b = S(N(a), T(a, b)), \quad a, b \in [0, 1].$$

Definition 2.10. [32] Let \rightarrow be a function $I : [0, 1]^2 \rightarrow [0, 1]$.

(i) \rightarrow is called an expansion type operator, if $a \rightarrow b \geq b$ holds for all $a \in [0, 1], b \in [0, 1]$.

(ii) \rightarrow is called a reduction type operator, if $a \rightarrow b \leq b$ holds for all $a \in [0, 1], b \in [0, 1]$.

(iii) \rightarrow is called an other type operator if it is neither expansion type nor reduction type.

Proposition 2.11. [32] If \rightarrow is an R -implication or S -implication, then \rightarrow is an expansion type operator.

It follows from Theorem 3.1 of [30] that Proposition 2.12 can be proved.

Proposition 2.12. Let \rightarrow be an implication satisfying

(C2) $a \rightarrow b$ is non-decreasing w.r.t. b ($a, b \in [0, 1]$),

(C3) $a \rightarrow b$ is right-continuous w.r.t. b ($a \in [0, 1], b \in [0, 1]$),

(C4) $a \leq b$ iff $a \rightarrow b = 1$ ($a, b \in [0, 1]$),

and define $T : [0, 1]^2 \rightarrow [0, 1]$ as follows:

$$T(a, b) = \bigwedge \{x \in [0, 1] \mid b \leq a \rightarrow x\}, \quad a, b \in [0, 1],$$

then (T, \rightarrow) is a residual pair, and (5) holds, where \bigwedge denotes infimum.

Definition 2.13. Let \rightarrow be an implication satisfying (C2), (C3) and (C4), and its residual function T satisfy $T(1, b) = b$, $T(0, b) = 0$ ($b \in [0, 1]$), then \rightarrow is called a (0,1)-implication.

It is easy to prove Lemma 2.14 (from residual condition (6)).

Lemma 2.14. Let \rightarrow be an implication satisfying (C2), (C3) and (C4), and T the function residual to \rightarrow , then ($a, b, c \in [0, 1]$) (i) if $b \leq c$, then $T(a, b) \leq T(a, c)$; (ii) $T(a, b) \leq a$; (iii) $T(0, b) = 0$; (iv) $T(a, a \rightarrow b) \leq b$.

It follows from Lemma 2.14 that $T(0, b) = 0$ can be deleted in Definition 2.13.

Proposition 2.15. If \rightarrow is an R -implication, then \rightarrow is a (0,1)-implication.

Proof. There uniquely exists T which is the residual function w.r.t. \rightarrow , and T is a left-continuous t-norm, thus we have $T(1, b) = b$, $T(0, b) = 0$ ($b \in [0, 1]$). It follows from Proposition 4.3 in [30] and Proposition 1 in [36] that \rightarrow satisfies (C2), (C3) and (C4). Thus it is easy to get that \rightarrow is a (0,1)-implication. \square

Definition 2.16. [34] Let T, \rightarrow be two $[0, 1]^2 \rightarrow [0, 1]$ functions. $([0, 1], T, \rightarrow)$ is called a residuated lattice, if the following conditions hold:

(i) $([0, 1], T)$ is a commutative semigroup with unit 1.

(ii) (T, \rightarrow) is a residual pair in which T is non-decreasing w.r.t. every variable, and \rightarrow is an implication where $a \rightarrow b$ is non-increasing w.r.t. a and non-decreasing w.r.t. b ($a, b \in [0, 1]$);

Lemma 2.17. [34] *If $([0, 1], T, \rightarrow)$ is a residuated lattice, then*

- (i) $\wedge\{a \rightarrow x_i | i \in I\} = a \rightarrow \wedge\{x_i | i \in I\}$ holds for any $a, x_i \in [0, 1]$ ($I \neq \emptyset$);
- (ii) \rightarrow satisfies (C1) and (C4).

Lemma 2.18. [30] *If \rightarrow satisfies $\wedge\{a \rightarrow x_i | i \in I\} = a \rightarrow \wedge\{x_i | i \in I\}$ where $a, x_i \in [0, 1]$, $I \neq \emptyset$, then \rightarrow satisfies (C2) and (C3).*

Proposition 2.19. *If $([0, 1], T, \rightarrow)$ is a residuated lattice, then \rightarrow is a $(0, 1)$ -implication.*

Proof. Since $([0, 1], T, \rightarrow)$ is a residuated lattice, (T, \rightarrow) is a residual pair, and it follows from Lemma 2.17 that the implication \rightarrow satisfies (C4), and $\wedge\{a \rightarrow x_i | i \in I\} = a \rightarrow \wedge\{x_i | i \in I\}$ holds for any $a, x_i \in [0, 1]$ ($I \neq \emptyset$). By Lemma 2.18, we get that \rightarrow satisfies (C2) and (C3). Meanwhile, since $([0, 1], T)$ is a commutative semigroup with unit 1, we have $T(a, b) = T(b, a)$, $T(1, b) = b$ ($a, b \in [0, 1]$). Then considering (T, \rightarrow) is a residual pair, we achieve that (T, \rightarrow) is a residual pair, thus \rightarrow is a $(0, 1)$ -implication. \square

Here we mainly consider 9 familiar operators. They are Lukasiewicz implication I_L , Fodor implication I_{FD} (see [2], which is also called I_0 implication, see [22, 33]), Gödel implication I_G , Goguen implication I_{Go} , revised Reichenbach implication I_{RR} (see [14, 21]), Zadeh implication I_Z , Kleene-Dienes implication I_{KD} , Mamdani operator I_M and Larsen operator I_{La} as the following.

$$\begin{aligned} I_L(a, b) &= \begin{cases} 1, & a \leq b \\ 1 - a + b, & a > b \end{cases}, & I_{FD}(a, b) &= \begin{cases} 1, & a \leq b \\ (1 - a) \vee b, & a > b \end{cases}, \\ I_G(a, b) &= \begin{cases} 1, & a \leq b \\ b, & a > b \end{cases}, & I_{Go}(a, b) &= \begin{cases} 1, & a \leq b \\ b/a, & a > b \end{cases}, \\ I_{RR}(a, b) &= \begin{cases} 1, & a \leq b \\ 1 - a + ab, & a > b \end{cases}, & I_Z(a, b) &= (1 - a) \vee (a \wedge b), \\ I_{KD}(a, b) &= (1 - a) \vee b, & I_M(a, b) &= a \wedge b, \\ I_{La}(a, b) &= a \times b. \end{aligned}$$

Here $I_L, I_{FD}, I_G, I_{Go}, I_{RR}$ satisfies (C1), (C2), (C3) and (C4); I_Z, I_{KD} satisfies (C1), (C2) and (C3); I_M, I_{La} satisfies (C2), (C3).

It is noted that I_M, I_{La} do not satisfy (C1), but they are also recognized by some authors (see e.g. [6, 11, 14, 32]). Moreover it is pointed out in [19] that I_M, I_{La} are referred to collectively as engineering implications. Therefore, for convenience, I_M, I_{La} can also be regarded as special implications.

It is easy to prove Propositions 2.20 and 2.21.

Proposition 2.20. *The operations residual to $I_L, I_{FD}, I_G, I_{Go}, I_{RR}$ are respectively:*

$$\begin{aligned} T_L(a, b) &= \begin{cases} a + b - 1, & a + b > 1 \\ 0, & a + b \leq 1 \end{cases}, & T_{FD}(a, b) &= \begin{cases} a \wedge b, & a + b > 1 \\ 0, & a + b \leq 1 \end{cases}, \\ T_G(a, b) &= a \wedge b, & T_{Go}(a, b) &= a \times b, \\ T_{RR}(a, b) &= \begin{cases} [(a + b - 1)/a] \wedge a, & a + b > 1 \\ 0, & a + b \leq 1 \end{cases}. \end{aligned}$$

Proposition 2.21. *In these implications, we have the following results:*

- (i) $I_L, I_{FD}, I_G, I_{Go}, I_{RR}$ are $(0,1)$ -implications.
- (ii) I_L, I_{FD}, I_G, I_{Go} are R -implications; I_L, I_{FD}, I_{KD} are S -implications; I_L, I_{FD}, I_{KD}, I_Z are QL -implications.
- (iii) $I_L, I_{FD}, I_G, I_{Go}, I_{RR}, I_{KD}$ are expansion type operators; I_M, I_{La} are reduction type operators; and I_Z is an other type operator.

Definition 2.22. Let Z be any nonempty set and $F(Z)$ the set of all fuzzy subsets on Z , define partial order relation \leq_F on $F(Z)$ (according to pointwise order) as: $A \leq_F B$ iff $A(z_0) \leq B(z_0)$ for any $z_0 \in Z$, where $A, B \in F(Z)$.

Lemma 2.23. $[35] < F(Z), \leq_F >$ is a complete lattice.

3. Solutions of Universal Triple I Method

For the FMP problem (1), from the point of view of universal triple I method, we can obtain the following principle:

Universal triple I principle: The conclusion B^* (in $< F(Y), \leq_F >$) of the FMP problem (1) is the smallest fuzzy set which makes (4) get its maximum for any $x \in X, y \in Y$.

Such principle improves the previous triple I principle for FMP in [33] or [36], since (4) is a generalization of (3) and the former can provide bigger choosing space.

Definition 3.1. Suppose that $A, A^* \in F(X)$, $B \in F(Y)$, if B^* (in $< F(Y), \leq_F >$) makes (4) get its maximum for any $x \in X, y \in Y$, then B^* is called a universal triple I solution.

Definition 3.2. Suppose that $A, A^* \in F(X)$, $B \in F(Y)$, and that nonempty set \mathbb{E} is the set of all universal triple I solutions, and finally that D^* (in $< F(Y), \leq_F >$) is the infimum of \mathbb{E} . Then D^* is called an Inf-quasi-solution. And, if D^* is the minimum of \mathbb{E} , then D^* is also called a Min-solution.

From Lemma 2.23, $< F(Y), \leq_F >$ is a complete lattice. Thus the Inf-quasi-solution (i.e., the infimum of \mathbb{E}) uniquely exists since the non-empty set $\mathbb{E} \subset F(Y)$.

Proposition 3.3. Suppose that \rightarrow_2 is an implication satisfying (C2), and that D_1 is a universal triple I solution, and finally that $D_1 \leq_F D_2$ (in which $D_1, D_2 \in < F(Y), \leq_F >$). Then D_2 is a universal triple I solution.

Proof. Since D_1 is a universal triple I solution, it follows that $(A(x) \rightarrow_1 B(y)) \rightarrow_2 (A^*(x) \rightarrow_2 D_1(y))$ takes its maximum for any $x \in X, y \in Y$. Because $D_1 \leq_F D_2$ and \rightarrow_2 satisfies (C2), we get that $A^*(x) \rightarrow_2 D_1(y) \leq A^*(x) \rightarrow_2 D_2(y)$ and

$$(A(x) \rightarrow_1 B(y)) \rightarrow_2 (A^*(x) \rightarrow_2 D_1(y)) \leq (A(x) \rightarrow_1 B(y)) \rightarrow_2 (A^*(x) \rightarrow_2 D_2(y))$$

hold for any $x \in X, y \in Y$. Therefore D_2 is also a universal triple I solution. \square

Remark 3.4. Suppose that \rightarrow_2 satisfies (C2). For (4), once there exists a universal triple I solution B^* , then every fuzzy set D which is larger than B^* ($D \in F(Y)$), will be a solution (it is easy to know from Proposition 3.3). This means that there

are many solutions, including $B^*(y) \equiv 1$ ($y \in Y$). This last is a special solution, for which (4) always takes its maximum no matter what major premise $A \rightarrow_1 B$ and minor premise A^* are adopted. Therefore, when the optimal universal triple I solution exists, it should be the smallest one; in other words, it should be the infimum of all solutions (i.e. the infimum of \mathbb{E}).

Theorem 3.5. *Suppose that \rightarrow_2 is a $(0,1)$ -implication, and that T the function residual to \rightarrow_2 , then the Min-solution can be computed as follows:*

$$B^*(y) = \sup_{x \in X} \{T(A^*(x), (A(x) \rightarrow_1 B(y)))\}, \quad y \in Y. \quad (7)$$

Proof. To begin with, we shall show that B^* (given by (7)) is a universal triple I solution, that is, the following formula holds for any $x \in X, y \in Y$:

$$(A(x) \rightarrow_1 B(y)) \rightarrow_2 (A^*(x) \rightarrow_2 B^*(y)) = 1. \quad (8)$$

In fact, it follows from (7) that $T(A^*(x), (A(x) \rightarrow_1 B(y))) \leq B^*(y)$, $x \in X, y \in Y$. Since (T, \rightarrow_2) is a residual pair, we get that $A(x) \rightarrow_1 B(y) \leq A^*(x) \rightarrow_2 B^*(y)$ holds for any $x \in X, y \in Y$. Thus (8) holds (considering \rightarrow_2 satisfies (C4)).

Furthermore we shall prove that B^* is the minimum of all universal triple I solutions. Suppose that D is any universal triple I solution, thus $(A(x) \rightarrow_1 B(y)) \rightarrow_2 (A^*(x) \rightarrow_2 D(y)) = 1$ holds for any $x \in X, y \in Y$. So $A(x) \rightarrow_1 B(y) \leq A^*(x) \rightarrow_2 D(y)$ holds for any $x \in X, y \in Y$ (noting that \rightarrow_2 satisfies (C4)). Then, from the fact that (T, \rightarrow_2) is a residual pair, we obtain $T(A^*(x), (A(x) \rightarrow_1 B(y))) \leq D(y)$ holds for any $x \in X, y \in Y$. Therefore, $D(y)$ is an upper bound of $\{T(A^*(x), (A(x) \rightarrow_1 B(y))) \mid x \in X\}$, $y \in Y$. Thus it follows from (7) that $B^* \leq_F D$. Thus, we obtain that B^* is the minimum of all universal triple I solutions.

Therefore B^* expressed as (7) is the Min-solution (from Definition 3.2). \square

We can get Proposition 3.6 from Theorem 3.5 and Proposition 2.15.

Proposition 3.6. *Suppose that \rightarrow_2 is an R-implication, and that T the function residual to \rightarrow_2 , then the Min-solution can be computed as (7).*

Lemma 3.7. [30] (i) *If \rightarrow_2 is I_M , then the Min-solution is $B^*(y) = \sup_{x \in X} \{A^*(x) \wedge (A(x) \rightarrow_1 B(y))\}$, $y \in Y$.*

(ii) *If \rightarrow_2 is I_Z , then the Min-solution is $B^*(y) = \sup_{x \in E_y} \{A^*(x) \wedge (A(x) \rightarrow_1 B(y))\}$, $y \in Y$, where $E_y = \{x \in X \mid (1 - A^*(x)) \vee 0.5 < A(x) \rightarrow_1 B(y)\}$.*

(iii) *If $\rightarrow_2 \in \{I_{KD}, I_{La}\}$, then the Min-solution is $B^*(y) = 1$ if $y \in E$, $B^*(y) = 0$ if $y \in Y - E$, where $E = \{y \in Y \mid \sup_{x \in X} \{(A(x) \rightarrow_1 B(y)) \times A^*(x)\} > 0\}$.*

Remark 3.8. When $\rightarrow_1 = \rightarrow_2$ in (4), the universal triple I method degenerates into the triple I method. From Lemma 3.7, it is easy to know that when \rightarrow_2 takes I_M , the universal triple I method degenerates into the CRI method.

4. Reversibility Property of Some Universal Triple I Methods

As pointed out in [36], the most fundamental deduction rule in logic is the classical modus ponens (meaning that if $A \rightarrow B$ and A are given, then B follows), thus it is natural to hope that the FMP conclusion B^* of (1) should be equal to

B if $A^* = A$, which is also called that such fuzzy reasoning method possesses the reversibility property (see Definition 4.1).

Definition 4.1. For a method (of fuzzy reasoning) to solve the FMP problem, if this method satisfies the classical modus ponens whenever condition P holds, then this method is said to be reversible under condition P , or P -reversible.

Here 3 conditions are considered as follows: (i) P_1 means that A is a normal fuzzy set (i.e., there exists $a \in X$ such that $A(a) = 1$). (ii) P_2 means that P_1 and $\{A(x)|x \in X\} \supset \{B(y)|y \in Y\}$ hold. (iii) P_3 means that P_1 and $B(y) \in (0, 5, 1] \cup \{0\}$ (for any $y \in Y$) hold.

When Wang proposed the triple I method, he pointed out the fact that the triple I method has the reversibility property while the CRI method has not (see [33]), which is an important evidence illustrating that the triple I method has stronger logicity. Following that, the reversibility property of triple I method becomes a hot research topic (see [24, 27, 34, 36]).

Similarly, we also need to analyze the reversibility property of universal triple I method, which is not discussed in [30]. Since the universal triple I method is decided by the first implication and second implication, we shall research aiming at some familiar implications (i.e., $\rightarrow_1, \rightarrow_2 \in \{I_{FD}, I_L, I_{Go}, I_G, I_{RR}, I_{KD}, I_M, I_{La}, I_Z\}$).

For convenience, for any fuzzy set $A(x)$, denote $A'(x) \triangleq 1 - A(x)$.

4.1. Reversibility Property for Expansion Type Operators. We shall investigate the case that \rightarrow_2 takes an expansion type operator (in detail, $I_{FD}, I_L, I_{Go}, I_G, I_{RR}, I_{KD}$).

Lemma 4.2. Let \rightarrow be a $(0,1)$ -implication, then \rightarrow satisfies the following condition: (C5) $1 \rightarrow b \geq b$, ($b \in [0, 1]$).

Theorem 4.3. If \rightarrow_2 is a $(0,1)$ -implication, and \rightarrow_1 satisfies (C5) and (C6) $a \rightarrow_1 b \leq a \rightarrow_2 b$ whenever $a > b$ ($a, b \in [0, 1]$), then the universal triple I method is P_1 -reversible.

Proof. Since \rightarrow_2 is a $(0,1)$ -implication, there uniquely exists the function T which is residual to \rightarrow_2 and $T(1, b) = b$ ($b \in [0, 1]$). When $A^* = A$, it follows from Theorem 3.5 that the Min-solution is $B^*(y) = \sup_{x \in X} \{T(A(x), (A(x) \rightarrow_1 B(y)))\}$, $y \in Y$.

Taking into account that P_1 holds, there exists $a \in X$ such that $A(a) = 1$. Since \rightarrow_2 is a $(0,1)$ -implication and \rightarrow_1 satisfies (C5), we obtain

$$B^*(y) \geq T(A(a), (A(a) \rightarrow_1 B(y))) = T(1, (1 \rightarrow_1 B(y))) = 1 \rightarrow_1 B(y) \geq B(y),$$

i.e. $B^*(y) \geq B(y)$.

We shall show $B^*(y) \leq B(y)$. For any $x \in X$, it follows Lemma 2.14(iv) that $T(A(x), (A(x) \rightarrow_2 B(y))) \leq B(y)$. Since \rightarrow_1 satisfies (C6), we get $a \rightarrow_1 b \leq a \rightarrow_2 b$ if $a > b$, and $a \rightarrow_2 b = 1 \geq a \rightarrow_1 b$ if $a \leq b$ (noting that \rightarrow_2 satisfies (C4)). Thus $a \rightarrow_1 b \leq a \rightarrow_2 b$ holds for any $a, b \in [0, 1]$. So, we get by Lemma 2.14(i) that $T(A(x), (A(x) \rightarrow_1 B(y))) \leq T(A(x), (A(x) \rightarrow_2 B(y))) \leq B(y)$. Then we have $B^*(y) \leq B(y)$.

Therefore $B^* = B$. □

From Proposition 2.15, we can further get Theorem 4.4 and Proposition 4.5.

Theorem 4.4. *If \rightarrow_2 is an R-implication, and \rightarrow_1 satisfies (C5) and (C6), then the universal triple I method is P_1 -reversible.*

Proposition 4.5. (i) *If \rightarrow_2 takes I_{FD} , and \rightarrow_1 satisfies (C5) and (C7) $a \rightarrow_1 b \leq (1-a) \vee b$ whenever $a > b$, then the universal triple I method is P_1 -reversible, especially for the case of $\rightarrow_1 \in \{I_{FD}, I_G, I_{KD}, I_M, I_{La}, I_Z\}$.*

(ii) *If \rightarrow_2 takes I_L , and \rightarrow_1 satisfies (C5) and (C8) $a \rightarrow_1 b \leq 1-a+b$ whenever $a > b$ ($a, b \in [0, 1]$), then the universal triple I method is P_1 -reversible, especially for the case of $\rightarrow_1 \in \{I_{FD}, I_L, I_{Go}, I_G, I_{RR}, I_{KD}, I_M, I_{La}, I_Z\}$.*

(iii) *If \rightarrow_2 takes I_{Go} , and \rightarrow_1 satisfies (C5) and (C9) $a \rightarrow_1 b \leq b/a$ whenever $a > b$, then the universal triple I method is P_1 -reversible, especially for the case of $\rightarrow_1 \in \{I_{Go}, I_G, I_M, I_{La}\}$.*

(iv) *If \rightarrow_2 takes I_G , and \rightarrow_1 satisfies (C5) and (C10) $a \rightarrow_1 b \leq b$ whenever $a > b$ ($a, b \in [0, 1]$), then the universal triple I method is P_1 -reversible, especially for the case of $\rightarrow_1 \in \{I_G, I_M, I_{La}\}$.*

(v) *If \rightarrow_2 takes I_{RR} , and \rightarrow_1 satisfies (C5) and (C11) $a \rightarrow_1 b \leq 1-a+ab$ whenever $a > b$ ($a, b \in [0, 1]$), then the universal triple I method is P_1 -reversible, especially for the case of $\rightarrow_1 \in \{I_{FD}, I_G, I_{RR}, I_{KD}, I_M, I_{La}, I_Z\}$.*

Remark 4.6. If \rightarrow_2 takes I_{FD} , and \rightarrow_1 satisfies (C5) but does not satisfy (C7), then the universal triple I method can not ensure to be P_1 -reversible. In fact, when $A(x) > B(y) > 0$, considering \rightarrow_1 does not satisfy (C7), there may exist $x_0 \in E_y$ such that $A(x_0) > B(y) > 0$ and $A(x_0) \rightarrow_1 B(y) > A(x_0) \vee B(y)$, then $A(x_0) \wedge (A(x_0) \rightarrow_1 B(y)) > B(y)$, thus $B^*(y) > B(y)$. Especially, if \rightarrow_2 takes I_{FD} , $\rightarrow_1 \in \{I_L, I_{Go}, I_{RR}\}$, then we have the same result. When $\rightarrow_2 \in \{I_L, I_{Go}, I_{RR}, I_G\}$, we can get similar analysis.

If \rightarrow_2 takes I_{KD} , then the universal triple I method can not ensure to be P -reversible where $P \in \{P_1, P_2, P_3\}$. In fact, when $A^* = A$, it follows from Lemma 3.7 that we can get the Min-solution $B^*(y)$. When $0 < B(y) < 1$, then $B^*(y) \neq B(y)$ holds (noting that $B^*(y) \in \{0, 1\}$).

Corollary 4.7. *Let \rightarrow_2 be a (0,1)-implication, and $\rightarrow_1 = \rightarrow_2 \triangleq \rightarrow$, then the universal triple I method (i.e. the triple I method) is P_1 -reversible.*

Proof. Since $\rightarrow_1 = \rightarrow_2$, we have that \rightarrow_1 satisfies (C6). Considering \rightarrow_1 is a (0,1)-implication, we get from Lemma 4.2 that $1 \rightarrow_1 b \geq b$ holds (i.e. \rightarrow_1 satisfies (C5)), thus it follows from Theorem 4.3 that the conclusion is correct. \square

From Corollary 4.7, Propositions 2.15, 2.19 and 2.21, we can easily get Corollaries 4.8, 4.9 and 4.10.

Corollary 4.8. Let $\rightarrow_2 \in \{I_{FD}, I_L, I_{Go}, I_G, I_{RR}\}$, and $\rightarrow_1 = \rightarrow_2 \triangleq \rightarrow$, then the universal triple I method (i.e. the triple I method) is P_1 -reversible.

Corollary 4.9. Let \rightarrow_2 be an R-implication, and $\rightarrow_1 = \rightarrow_2 \triangleq \rightarrow$, then the universal triple I method (i.e. the triple I method) is P_1 -reversible.

Corollary 4.10. Let $([0, 1], T, \rightarrow)$ be a residuated lattice, and $\rightarrow_1 = \rightarrow_2 \triangleq \rightarrow$, then the universal triple I method (i.e. the triple I method) is P_1 -reversible.

Remark 4.11. In [33, 34], Wang discussed the reversibility property of triple I method (for the FMP problem), which was only aiming at the case of I_{FD} . And Theorem 4 in [33] and Theorem 4.4.12 in [34] pointed out the fact that the triple I method via I_{FD} is P_1 -reversible. It is evident that this conclusion is the same as the related one of Corollary 4.8 in this paper.

Remark 4.12. In Theorem 5 of [36] and Theorem 6 of [24], Wang and Pei all gave the same result that the triple I method (for the FMP problem) is P_1 -reversible if the implication takes an R-implication, which coincides with Corollary 4.9 in this paper. Moreover, it follows from Proposition 2.15 in this paper that Theorem 5 of [36] (or Theorem 6 in [24]) is a special case of Corollary 4.7 in this paper.

Remark 4.13. In Theorem 5 of [23], Pei got the result that if $([0, 1], T, \rightarrow)$ is a residuated lattice, then the triple I method (for the FMP problem) employing \rightarrow is P_1 -reversible, which is the same as Corollary 4.10 in this paper. Moreover, it follows from Proposition 2.19 in this paper that Theorem 5 of [23] is a special case of Corollary 4.7 in this paper.

4.2. Reversibility Property for Reduction Type Operator. We shall investigate the case that \rightarrow_2 takes a reduction type operator (in detail, I_M, I_{La}).

Theorem 4.14. If \rightarrow_2 takes I_M , and \rightarrow_1 satisfies (C5) and (C10), then the universal triple I method is P_1 -reversible, especially for the case of $\rightarrow_1 \in \{I_G, I_M, I_{La}\}$.

Proof. It follows from Lemma 3.7 that when $A^* = A$, the Min-solution $B^*(y) = \sup_{x \in X} \{A(x) \wedge (A(x) \rightarrow_1 B(y))\}$.

Since P_1 is satisfied, there exists $a \in X$ such that $A(a) = 1$, then we get $B^*(y) \geq 1 \wedge (1 \rightarrow_1 B(y)) \geq B(y)$ (noting that \rightarrow_1 satisfies (C5)), i.e. $B^*(y) \geq B(y)$. For any $x \in X$, if $A(x) \leq B(y)$, then $A(x) \wedge (A(x) \rightarrow_1 B(y)) \leq A(x) \leq B(y)$; if $A(x) > B(y)$, then $A(x) \wedge (A(x) \rightarrow_1 B(y)) \leq A(x) \rightarrow_1 B(y) \leq B(y)$ (noting that \rightarrow_1 satisfies (C10)), thus $B^*(y) \leq B(y)$.

Therefore, $B^* = B$. □

Remark 4.15. If \rightarrow_2 takes I_M , and \rightarrow_1 satisfies (C5) but does not satisfy (C10), then it is similar to Remark 4.6 that the universal triple I method can not ensure to be P_1 -reversible. Especially, if $\rightarrow_1 \in \{I_{FD}, I_L, I_{Go}, I_{RR}, I_{KD}, I_Z\}$, we have the same conclusion.

If \rightarrow_2 takes I_{La} , then it is similar to the case of I_{KD} that the universal triple I method can not ensure to be P -reversible where $P \in \{P_1, P_2, P_3\}$.

4.3. Reversibility Property for Other Type Operators. Noting that I_Z is an other type operator (and also a typical QL-implication), we shall analyze the case that \rightarrow_2 takes I_Z .

Theorem 4.16. *If \rightarrow_2 takes I_Z , and \rightarrow_1 satisfies (C5), (C7) together with (C12) $a \rightarrow_1 a = 1$ whenever $0 < a \leq 0.5$ ($a \in [0, 1]$), then the universal triple I method is P_2 -reversible.*

Proof. When $A^* = A$, we get from Lemma 3.7 that the Min-solution is $B^*(y) = \sup_{x \in E_y} \{A(x) \wedge (A(x) \rightarrow_1 B(y))\}$ where $E_y = \{x \in X \mid A'(x) \vee 0.5 < A(x) \rightarrow_1 B(y)\}$.

(i) Suppose $B(y) = 0$. We shall observe the structure of E_y . If $A(x) > 0$, considering \rightarrow_1 satisfies (C7), we have $A'(x) < A(x) \rightarrow_1 B(y) \leq A'(x) \vee 0 = A'(x)$, which is a contradiction. If $A(x) = 0$, then $A'(x) \vee 0.5 = 1 < A(x) \rightarrow_1 B(y)$, which is also a contradiction. Thus we get $E_y = \emptyset$, therefore $B^*(y) = 0 = B(y)$.

(ii) Suppose $B(y) > 0$. At first, we shall show $B^*(y) \geq B(y)$. (a) Suppose $B(y) > 0.5$. Since P_2 holds, there exists $a \in X$ such that $A(a) = 1$, thus $A(a) \rightarrow_1 B(y) \geq B(y) > A'(a) \vee 0.5$ (noting that \rightarrow_1 satisfies (C5)). Thus $a \in E_y$, and then $B^*(y) \geq A(a) \wedge (A(a) \rightarrow_1 B(y)) \geq B(y)$, i.e. $B^*(y) \geq B(y)$. (b) Suppose $0 < B(y) \leq 0.5$. Since P_2 is satisfied, there exists $b \in X$ such that $A(b) = B(y)$, so $A(b) \rightarrow_1 B(y) = 1 > A'(b) \vee 0.5$ (noting that \rightarrow_1 satisfies (C12)). Thus $b \in E_y$, and then $B^*(y) \geq A(b) \wedge (A(b) \rightarrow_1 B(y)) = A(b) = B(y)$, i.e. $B^*(y) \geq B(y)$.

Furthermore, we shall prove $B^*(y) \leq B(y)$ when $B(y) > 0$. Inspecting the process above, it is easy to know E_y is not empty. For any $x \in E_y$, $A'(x) \vee 0.5 < A(x) \rightarrow_1 B(y)$ holds. If $A(x) \leq B(y)$, then $A(x) \wedge (A(x) \rightarrow_1 B(y)) \leq A(x) \leq B(y)$. If $A(x) > B(y)$, then $A'(x) \vee 0.5 < A(x) \rightarrow_1 B(y) \leq A'(x) \vee B(y)$ (noting that \rightarrow_1 satisfies (C7)), thus $A'(x) < B(y)$, and $A(x) \wedge (A(x) \rightarrow_1 B(y)) \leq A(x) \wedge [A'(x) \vee B(y)] = A(x) \wedge B(y) \leq B(y)$. To sum up, we have $B^*(y) \leq B(y)$.

Therefore $B^* = B$ whenever $B(y) > 0$. Summarizing above, $B^* = B$ holds. \square

From Theorem 4.16, we can easily get Proposition 4.17.

Proposition 4.17. *Let \rightarrow_2 take I_Z , and $\rightarrow_1 \in \{I_{FD}, I_G\}$, then the universal triple I method is P_2 -reversible.*

Remark 4.18. If \rightarrow_2 takes I_Z , and \rightarrow_1 satisfies (C5), (C12) but does not satisfy (C7), then it is similar to Remark 4.6 that $B^*(y) > B(y)$ may hold when $A(x) > B(y) > 0$, and thus the universal triple I method can not ensure to be P_2 -reversible. Especially, if $I_1 \in \{I_L, I_{Go}, I_{RR}\}$, we have the same result.

Proposition 4.19. *If \rightarrow_2 takes I_Z , $\rightarrow_1 \in \{I_{KD}, I_M, I_{La}, I_Z\}$ and A is a normal fuzzy set. Then, when $A^* = A$, $B^*(y) = 0$ if $B(y) \leq 0.5$, $B^*(y) = B(y)$ if $B(y) > 0.5$.*

Proof. Suppose $A^* = A$. It is obvious that the Min-solution $B^*(y)$ is the same as Theorem 4.16. Note that I_{KD}, I_M, I_{La}, I_Z obviously satisfy (C5) and (C7) (but do not satisfy (C12)), then from the proving process of Theorem 4.16, it is easy to get $B^*(y) = 0$ if $B(y) = 0$, and $B^*(y) = B(y)$ if $B(y) > 0.5$. We shall show $B^*(y) = 0$

when $0 < B(y) \leq 0.5$. We only prove the case of $\rightarrow_1 = I_{KD}$ as an example. Suppose $\rightarrow_1 = I_{KD}$, then

$$E_y = \{x \in X \mid A'(x) \vee 0.5 < A'(x) \vee B(y)\} = \{x \in X \mid A'(x) \vee 0.5 < B(y)\}.$$

When $0 < B(y) \leq 0.5$, we can easily get $E_y = \emptyset$, and then $B^*(y) = 0$. \square

It follows from Proposition 4.19 that we can easily get Theorem 4.20.

Theorem 4.20. *Let \rightarrow_2 take I_Z , and $\rightarrow_1 \in \{I_{KD}, I_M, I_{La}, I_Z\}$, then the universal triple I method is P_3 -reversible.*

Remark 4.21. If \rightarrow_2 takes I_Z , and \rightarrow_1 satisfies (C5), (C7) but does not satisfy (C12), then $B^*(y) \neq B(y)$ may hold when $0 < B(y) \leq 0.5$, and the universal triple I method can not ensure to be P_2 -reversible. Especially, if $\rightarrow_1 \in \{I_{KD}, I_M, I_{La}, I_Z\}$, we have the same result (from Proposition 4.19 and Theorem 4.20); but under more strict condition (i.e. P_3), the universal triple I method possesses the reversibility property. In [27], Song investigated the triple I method based on I_Z and its reversibility property, and got Theorem 4 and Corollary 5 for the reversibility property of triple I method (for the FMP problem), which are respectively the same as the related conclusions of Proposition 4.19 and Theorem 4.20 in this paper.

4.4. Summarizations of Reversibility Property of Universal Triple I Methods. We shall summarize the reversibility property of universal triple I methods. By Remark 3.8, when $\rightarrow_2 = I_M$, the universal triple I method degenerates into the CRI method, then we get the reversibility property of related CRI methods by Theorem 4.14 and Remark 4.15 (see Proposition 4.22). When $\rightarrow_1 = \rightarrow_2$, the universal triple I method degenerates into the triple I method. Inspecting the results mentioned above, we get the reversibility property of triple I methods (see Proposition 4.23).

Proposition 4.22. *Let \rightarrow_2 take I_M , $\rightarrow_1 \in \{I_G, I_M, I_{La}\}$, then the universal triple I method (i.e. the CRI method) is P_1 -reversible.*

Proposition 4.23. *Take $\rightarrow_1 = \rightarrow_2 \triangleq \rightarrow$, then the universal triple I method (that is, the triple I method) is P_1 -reversible if $\rightarrow \in \{I_{FD}, I_L, I_{Go}, I_G, I_{RR}, I_M\}$, and P_3 -reversible if $\rightarrow \in \{I_Z\}$.*

Remark 4.24. In Theorem 3.1 of [5], Hou and Li pointed out the fact that the CRI method where $\rightarrow \in \{I_G, I_M, I_{La}\}$ is P_1 -reversible, and the one where $\rightarrow \in \{I_{FD}, I_L, I_{Go}, I_{RR}, I_{KD}, I_Z\}$ is not, which are the same as the related conclusions of Proposition 4.22 and Remark 4.15 in this paper. In Theorem 3.2 of [5], Hou and Li drew the conclusions that the triple I method (for the FMP problem) where $\rightarrow \in \{I_{FD}, I_L, I_{Go}, I_G, I_{RR}, I_M\}$ is P_1 -reversible, and the one where $\rightarrow \in \{I_{KD}, I_{La}, I_Z\}$ is not, which coincide with the related conclusions of Proposition 4.5, Remark 4.6 and Proposition 4.23 in this paper. From Propositions 4.22 and 4.23, the triple I method is superior to the CRI method from the viewpoint of reversibility property.

Table 1 summarizes the reversibility property of universal triple I method (except the case of $\rightarrow_2 \in \{I_{KD}, I_{La}\}$), where P_i represents that corresponding universal

| $\rightarrow_2 \backslash \rightarrow_1$ | I_{FD} | I_L | I_{Go} | I_G | I_{RR} | I_{KD} | I_M | I_{La} | I_Z |
|--|----------|-------|----------|-------|----------|----------|-------|----------|-------|
| I_{FD} | P_1 | | | P_1 | | P_1 | P_1 | P_1 | P_1 |
| I_L | P_1 | P_1 | P_1 | P_1 | P_1 | P_1 | P_1 | P_1 | P_1 |
| I_{Go} | | | P_1 | P_1 | | | P_1 | P_1 | |
| I_G | | | | P_1 | | | P_1 | P_1 | |
| I_{RR} | P_1 | | | P_1 | P_1 | P_1 | P_1 | P_1 | P_1 |
| I_M | | | | P_1 | | | P_1 | P_1 | |
| I_Z | P_2 | | | P_2 | | P_3 | P_3 | P_3 | P_3 |

TABLE 1. The Reversibility Property of Some Universal Triple I Methods

triple I method (employing first implication \rightarrow_1 and second implication \rightarrow_2) is P_i -reversible. The universal triple I method possesses reversibility property under very strict condition if \rightarrow_2 takes I_Z (which is an other type operator); and it hardly is reversible if $\rightarrow_2 \in \{I_{KD}, I_{La}\}$. However, when \rightarrow_2 takes a (0,1)-implication (which belongs to an expansion type operator), the reversibility property seems excellent, which is embodied as that the universal triple I method is P -reversible for a lot of first implications. By the way, when \rightarrow_1 takes a reduction type operator (e.g., I_M, I_{La}), the reversibility property seems fine.

From another viewpoint, when \rightarrow_2 takes an R-implication (e.g., I_{FD}, I_L, I_{Go}, I_G), the reversibility property of universal triple I method seems fine. When \rightarrow_2 is an S-implication (e.g., I_{FD}, I_L, I_{KD}) or a QL-implication (e.g., I_{FD}, I_L, I_{KD}, I_Z), the reversibility property is uncertain, where the case of $\rightarrow_2 \in \{I_{KD}\}$ is unacceptable, and the case of $\rightarrow_2 \in \{I_Z\}$ is complicated, and finally the case of $\rightarrow_2 \in \{I_{FD}, I_L\}$ is excellent.

5. Response Ability of Fuzzy Controllers Based on Universal Triple I Methods

In this section, we shall investigate the universal triple I methods from the viewpoint of fuzzy controllers.

First of all, we shall review briefly Mamdanian fuzzy control algorithm (which approximates to an interpolation function), where takes the SISO fuzzy controller as an example, thus some necessary concepts and signs are introduced.

Let X and Y be the input and output universe, respectively. Denote $\mathbb{A} = \{A_i\}_{(1 \leq i \leq n)}$, $\mathbb{B} = \{B_i\}_{(1 \leq i \leq n)}$ where $A_i \in F(X)$, $B_i \in F(Y)$. \mathbb{A}, \mathbb{B} are regarded as linguistic variables, thus the fuzzy reasoning rules can be expressed as follows:

$$\text{If } x \text{ is } A_i, \text{ then } y \text{ is } B_i, \quad i = 1, \dots, n, \quad (9)$$

where $x \in X, y \in Y$ are called base variables.

According to the Mamdanian fuzzy control algorithm, the inference relation of the i -th rule can be regarded as a fuzzy relation from X to Y ($i = 1, \dots, n$), denoting by $A_i(x) \rightarrow_1 B_i(y)$ (where \rightarrow_1 is an implication). And such n rules can be connected by 'or' relation (i.e, taking 'max' operator for rules), thus the whole rule is

$$IR(x, y) \triangleq \bigvee_{i=1}^n (A_i(x) \rightarrow_1 B_i(y)).$$

Given $A^* \in F(X)$, the inference conclusion $B^* \in F(Y)$ can be obtained as $B^* \triangleq A^* \circ IR$, by the CRI method, in which

$$B^*(y) = \sup_{x \in X} \{A^*(x) \wedge IR(x, y)\}, \quad y \in Y. \quad (10)$$

For a fuzzy controller, since its input is a crisp quantity, it should be transformed into fuzzy set to utilize (10) by

$$A^*(x) = \begin{cases} 1, & x = x^* \\ 0, & x \neq x^* \end{cases} \triangleq A_{x^*}^*,$$

which is called a singleton fuzzification.

Furthermore, it is known that B^* should be turned into a crisp quantity by using some defuzzification methods, and the commonly used method is the so-called method of centroid:

$$y^* = \frac{\int_Y y B^*(y) dy}{\int_Y B^*(y) dy}. \quad (11)$$

Then, a natural problem arise: What is interpolation representation of such fuzzy controller if we replace the CRI method with the universal triple I method? This is the task of this section.

5.1. SISO Fuzzy Controllers Based on Universal Triple I Methods. Here we employ the universal triple I method instead of the CRI method. Thus, for fuzzy reasoning rules (9), it is easy to get that (4) should be turned into:

$$IR(x, y) \rightarrow_2 (A^*(x) \rightarrow_2 B^*(y)). \quad (12)$$

Therefore, there is an output $y^* = F(x^*)$ for each input x^* . Thus a SISO fuzzy controller based on universal triple I method is obtained.

Definition 5.1. Let Z be any nonempty set and $\mathbb{C} = \{C_i\}_{(1 \leq i \leq n)}$ a family of normal fuzzy sets on Z , where the peak-point of C_i is z_i (i.e. the unique point satisfying $C_i(z_i) = 1$ in Z). \mathbb{C} is called a fuzzy partition of Z if $(\forall z \in Z)(\sum_{i=1}^n C_i(z) = 1)$ holds, and C_i is defined as a base element in \mathbb{C} . Thus \mathbb{C} is also called a group of base elements of Z .

Remark 5.2. Definition 5.1 obviously implies $(\forall i, j)(i \neq j \Rightarrow z_i \neq z_j)$ and that \mathbb{C} has Kronecker property (i.e. $C_i(z_j) = \delta_{ij}$ where $\delta_{ij} = 1$ if $i = j$, $\delta_{ij} = 0$ if $i \neq j$).

To investigate interpolation mechanism of fuzzy controller, suppose that \mathbb{A} and \mathbb{B} are respectively fuzzy partitions of X and Y (in which A_i, B_i are integrable functions). We assume that X and Y are all real number intervals, e.g. $X = [a, b]$ and $Y = [c, d]$ where $a < x_1 < x_2 < \dots < x_n < b$, $c < y_1 < y_2 < \dots < y_n < d$, in which x_i, y_i are respectively peak-points of A_i, B_i .

Let $h_1 = y_1 - c$, $h_i = y_i - y_{i-1}$ ($i = 2, 3, \dots, n$) and $h = \max_{1 \leq i \leq n} \{h_i\}$. Since \mathbb{A} and \mathbb{B} are all fuzzy partitions, they have Kronecker property: $A_i(x_j) = \delta_{ij} = B_i(y_j)$ ($i, j = 1, \dots, n$). By the definition of definite integral, we achieve for the method of centroid:

$$y^* = \frac{\int_Y y B^*(y) dy}{\int_Y B^*(y) dy} \approx \frac{\sum_{i=1}^n y_i B^*(y_i) h_i}{\sum_{i=1}^n B^*(y_i) h_i}. \quad (13)$$

Similar to Theorem 3.5, we can prove Proposition 5.3.

Proposition 5.3. *Suppose that \rightarrow_2 is a $(0,1)$ -implication, and that T the function residual to \rightarrow_2 , then the Min-solution derived from (12) can be expressed as follows:*

$$B^*(y) = \sup_{x \in X} \{T(A^*(x), IR(x, y))\}, \quad y \in Y.$$

Theorem 5.4. *Let \rightarrow_2 be a $(0,1)$ -implication, then the Min-solution $B^*(y) = IR(x^*, y)$ for the SISO fuzzy controller based on universal triple I method.*

Proof. Let \rightarrow_2 be a $(0,1)$ -implication. Then there uniquely exists T which is the residual function w.r.t. \rightarrow_2 , and $T(1, b) = b$, $T(0, b) = 0$ ($b \in [0, 1]$). It follows from Proposition 5.3 that the Min-solution $B^*(y) = \sup_{x \in X} \{T(A^*(x), IR(x, y))\}$, $y \in Y$. As for input x^* , we get $A_{x^*}^* = \begin{cases} 1, & x = x^* \\ 0, & x \neq x^* \end{cases}$. Because $T(1, b) = b$, $T(0, b) = 0$ ($b \in [0, 1]$), we obtain $T(A^*(x), IR(x, y)) = T(1, IR(x^*, y)) = IR(x^*, y)$ for the case of $x = x^*$, and $T(A^*(x), IR(x, y)) = T(0, IR(x, y)) = 0$ for the case of $x \in X - \{x^*\}$. Therefore $B^*(y) = \sup_{x \in X} \{T(A^*(x), IR(x, y))\} = IR(x^*, y)$. \square

Theorem 5.4 gives the equivalent form of Min-solution in SISO fuzzy controller.

Lemma 5.5. *In the SISO fuzzy controller based on universal triple I method,*

(i) *if \rightarrow_1 is an expansion type operator, then $IR(x^*, y_j) > 0$ for any $x^* \in X$ ($j = 1, \dots, n$);*

(ii) *if \rightarrow_1 is a reduction type operator satisfying*

$$(C13) \quad a \rightarrow 1 = a \quad (a \in [0, 1]),$$

then there exists y_j such that $IR(x^, y_j) > 0$ for any $x^* \in X$ ($j \in \{1, \dots, n\}$);*

(iii) *if \rightarrow_1 is an other type operator satisfying*

$$(C14) \quad a \rightarrow 1 > 0 \quad (a \in [0, 1]),$$

then $IR(x^, y_j) > 0$ for any $x^* \in X$ ($j = 1, \dots, n$);*

(iv) *if \rightarrow_1 is an R-implication or S-implication, then $IR(x^*, y_j) > 0$ for any $x^* \in X$ ($j = 1, \dots, n$);*

(v) *if \rightarrow_1 is a QL-implication, then $IR(x^*, y_j) > 0$ for any $x^* \in X$ ($j = 1, \dots, n$).*

Proof. In the SISO fuzzy controller based on universal triple I method, suppose any $x^* \in X$, then $IR(x^*, y) = \bigvee_{i=1}^n (A_i(x^*) \rightarrow_1 B_i(y))$.

(i) Suppose that \rightarrow_1 is an expansion type operator. Then $a \rightarrow_1 1 = 1$, and thus $IR(x^*, y_j) = \bigvee_{i=1}^n (A_i(x^*) \rightarrow_1 B_i(y_j)) = 1 > 0$ holds ($j = 1, \dots, n$) since $B_i(y_j) = \delta_{ij}$ ($i, j = 1, \dots, n$).

(ii) Suppose that \rightarrow_1 is a reduction type operator satisfying (C13). Then we have $a \rightarrow_1 0 = 0$ ($a \in [0, 1]$). Suppose, on the contrary, that there exists $x^* \in X$ such that $IR(x^*, y_j) = 0$ for any y_j ($j = 1, \dots, n$). Since $B_i(y_j) = \delta_{ij}$ ($i, j = 1, \dots, n$) and $a \rightarrow_1 1 = a$, $a \rightarrow_1 0 = 0$, we get $0 = IR(x^*, y_j) = \bigvee_{i=1}^n (A_i(x^*) \rightarrow_1 B_i(y_j)) = A_j(x^*)$ ($j = 1, \dots, n$), and then $\sum_{j=1}^n A_j(x^*) = 0$. But, from the previous assumption, $\mathbb{A} = \{A_i\}_{(1 \leq i \leq n)}$ is a fuzzy partition on X , then it follows from Definition 5.1 that $\sum_{j=1}^n A_j(x^*) = 1$, which is a contradiction. Therefore, there exists y_j such that $IR(x^*, y_j) > 0$ for any $x^* \in X$ ($j \in \{1, \dots, n\}$).

(iii) Suppose that \rightarrow_1 is an other type operator satisfying (C14). Since $B_i(y_j) = \delta_{ij}$ ($i, j = 1, \dots, n$), we get $IR(x^*, y_j) = \bigvee_{i=1}^n (A_i(x^*) \rightarrow_1 B_i(y_j)) \geq A_j(x^*) \rightarrow_1 B_j(y_j) = A_j(x^*) \rightarrow_1 1 > 0$ holds ($j = 1, \dots, n$).

(iv) Since an R-implication or S-implication is also an expansion type operator, then similarly we get the conclusion.

(v) Suppose that \rightarrow_1 is a QL-implication, then it is easy to get $a \rightarrow_1 1 > 0$, thus we also get the conclusion. \square

Theorem 5.6. *Let \rightarrow_2 be a (0,1)-implication. If \rightarrow_1 is a reduction type operator satisfying (C13), then there exists a group of base functions $\mathbb{A}^* = \{A_i^*\}_{(1 \leq i \leq n)}$ such that the SISO fuzzy controller based on universal triple I method is approximately a univariate piecewise interpolation function regarding A_i^* as its base functions, i.e. $F(x) = \sum_{i=1}^n A_i^*(x)y_i$, and \mathbb{A}^* is a fuzzy partition of X . Moreover, if $\{y_i\}_{(1 \leq i \leq n)}$ is an equidistant partition, then \mathbb{A}^* degenerates into \mathbb{A} , i.e. $F(x) = \sum_{i=1}^n A_i(x)y_i$.*

Proof. Since \rightarrow_2 is a (0,1)-implication, it follows from Theorem 5.4 that Min-solution $B^*(y) = IR(x^*, y) = \bigvee_{k=1}^n (A_k(x^*) \rightarrow_1 B_k(y))$. Since \rightarrow_1 is a reduction type operator satisfying (C13), then $a \rightarrow_1 1 = a$, $a \rightarrow_1 0 = 0$. Noting $B_k(y_i) = \delta_{ki}$ ($k, i = 1, \dots, n$), it follows from (13) that

$$\begin{aligned} y^* &\approx \frac{\sum_{i=1}^n y_i B^*(y_i) h_i}{\sum_{i=1}^n B^*(y_i) h_i} = \frac{\sum_{i=1}^n h_i [\bigvee_{k=1}^n (A_k(x^*) \rightarrow_1 B_k(y_i))] y_i}{\sum_{i=1}^n h_i [\bigvee_{k=1}^n (A_k(x^*) \rightarrow_1 B_k(y_i))]} \\ &= \frac{\sum_{i=1}^n h_i A_i(x^*) y_i}{\sum_{i=1}^n h_i A_i(x^*)}, \end{aligned} \quad (14)$$

where there exists y_i such that $B^*(y_i) = IR(x^*, y_i) > 0$ ($i \in \{1, \dots, n\}$) by Lemma 5.5(ii), so $\sum_{i=1}^n B^*(y_i) h_i > 0$ and then (14) makes sense.

Denote $A_i^*(x^*) \triangleq h_i A_i(x^*) / (\sum_{i=1}^n h_i A_i(x^*))$, then $y^* \approx \sum_{i=1}^n A_i^*(x^*) y_i$. Let $\mathbb{A}^* \triangleq \{A_i^*\}_{(1 \leq i \leq n)}$, $F(x) \triangleq \sum_{i=1}^n A_i^*(x) y_i$. We get

$$F(x_i) = \sum_{k=1}^n A_k^*(x_i) y_k = \frac{\sum_{k=1}^n h_k A_k(x_i) y_k}{\sum_{k=1}^n h_k A_k(x_i)} = y_i \quad (i = 1, \dots, n),$$

noting that $A_k(x_i) = \delta_{ki}$ ($i, k = 1, \dots, n$), then $F(x)$ is a univariate piecewise interpolation function which regards A_i^* as its base functions.

Furthermore, $\sum_{i=1}^n A_i^*(x) = \sum_{i=1}^n [h_i A_i(x) / (\sum_{i=1}^n h_i A_i(x))] = 1$ holds for any $x \in X$, thus \mathbb{A}^* is a fuzzy partition of X . At last, if $\{y_i\}_{(1 \leq i \leq n)}$ is an equidistant partition (i.e. $(\forall i)(h_i = h)$), then it is evident that $A_i^* = A_i$, $\mathbb{A}^* = \mathbb{A}$, and hence $F(x) = \sum_{i=1}^n A_i(x) y_i$. \square

Note that I_M, I_{La} are reduction type operators satisfying (C13), which implies Corollary 5.7 (by virtue of Theorem 5.6).

Corollary 5.7. *Let \rightarrow_2 be a (0,1)-implication. If $\rightarrow_1 \in \{I_M, I_{La}\}$, then there exists a group of base functions $\mathbb{A}^* = \{A_i^*\}_{(1 \leq i \leq n)}$ such that the SISO fuzzy controller based on universal triple I method is approximately a univariate piecewise interpolation function regarding A_i^* as its base functions and \mathbb{A}^* is a fuzzy partition of X . Moreover, if $\{y_i\}_{(1 \leq i \leq n)}$ is an equidistant partition, then \mathbb{A}^* degenerates into \mathbb{A} .*

Since I_Z is an other type operator satisfying (C14), and it is also a typical QL-implication, we shall investigate the case that \rightarrow_1 takes I_Z .

Theorem 5.8. *Let \rightarrow_2 be a $(0,1)$ -implication. If \rightarrow_1 is I_Z , then there exists a group of base functions $A^* = \{A_i^*\}_{(1 \leq i \leq n)}$ such that the SISO fuzzy controller based on universal triple I method is approximately a univariate piecewise fitted function regarding A_i^* as its base functions, i.e. $F(x) = \sum_{i=1}^n A_i^*(x)y_i$.*

Proof. It is similar to Theorem 5.6 that $B^*(y) = \bigvee_{k=1}^n (A_k(x^*) \rightarrow_1 B_k(y))$. Since \rightarrow_1 is I_Z and $B_k(y_i) = \delta_{ki}$ ($k, i = 1, \dots, n$), it follows from (13) that

$$\begin{aligned} y^* &\approx \frac{\sum_{i=1}^n y_i B^*(y_i) h_i}{\sum_{i=1}^n B^*(y_i) h_i} = \frac{\sum_{i=1}^n h_i [\bigvee_{k=1}^n ((1 - A_k(x^*)) \vee (A_k(x^*) \wedge B_k(y_i)))] y_i}{\sum_{i=1}^n h_i [\bigvee_{k=1}^n ((1 - A_k(x^*)) \vee (A_k(x^*) \wedge B_k(y_i)))]} \\ &= \frac{\sum_{i=1}^n h_i [A_i(x^*) \vee (\bigvee_{k=1}^n (1 - A_k(x^*)))] y_i}{\sum_{i=1}^n h_i [A_i(x^*) \vee (\bigvee_{k=1}^n (1 - A_k(x^*)))]}, \end{aligned} \quad (15)$$

where $B^*(y_i) = IR(x^*, y_i) > 0$ ($i = 1, \dots, n$) according to Lemma 5.5(iii)(v), so $\sum_{i=1}^n B^*(y_i) h_i > 0$ and then (15) makes sense.

Denote

$$C_i(x^*) \triangleq A_i(x^*) \vee (\bigvee_{k=1}^n (1 - A_k(x^*))), \quad A_i^*(x^*) \triangleq h_i C_i(x^*) / (\sum_{i=1}^n h_i C_i(x^*)),$$

then $y^* \approx \sum_{i=1}^n A_i^*(x^*) y_i$. Let $A^* \triangleq \{A_i^*\}_{(1 \leq i \leq n)}$ and $F(x) \triangleq \sum_{i=1}^n A_i^*(x) y_i$. It is easy to verify that it can't make $F(x_i) = y_i$ hold for any i , thus $F(x)$ is a univariate piecewise fitted function which regards A_i^* as its base functions. \square

Theorem 5.9. *Let \rightarrow_2 be a $(0,1)$ -implication. If \rightarrow_1 is an expansion type operator, then the SISO fuzzy controller based on universal triple I method is approximately a step response function.*

Proof. It is similar to Theorem 5.6 that $B^*(y) = \bigvee_{k=1}^n (A_k(x^*) \rightarrow_1 B_k(y))$. Since \rightarrow_1 is an expansion type operator, then $a \rightarrow_1 1 = 1$. Noting $B_k(y_i) = \delta_{ki}$ ($k, i = 1, \dots, n$), it follows from (13) that

$$y^* \approx \frac{\sum_{i=1}^n h_i [\bigvee_{k=1}^n (A_k(x^*) \rightarrow_1 B_k(y_i))] y_i}{\sum_{i=1}^n h_i [\bigvee_{k=1}^n (A_k(x^*) \rightarrow_1 B_k(y_i))]} = \frac{\sum_{i=1}^n h_i y_i}{\sum_{i=1}^n h_i} \triangleq c_0, \quad (16)$$

where $B^*(y_i) = IR(x^*, y_i) > 0$ ($i = 1, \dots, n$) according to Lemma 5.5(i), thus $\sum_{i=1}^n B^*(y_i) h_i > 0$ and then (16) makes sense. \square

We can prove Theorem 5.10 using Theorem 5.9 and Proposition 2.11.

Theorem 5.10. *Let \rightarrow_2 be a $(0,1)$ -implication. If \rightarrow_1 is an R -implication or S -implication, then the SISO fuzzy controller based on universal triple I method is approximately a step response function.*

Note that $I_{FD}, I_L, I_{Go}, I_G, I_{RR}, I_{KD}$ are expansion type operators, we can get Corollary 5.11.

Corollary 5.11. *Let \rightarrow_2 be a $(0,1)$ -implication. If $\rightarrow_1 \in \{I_{FD}, I_L, I_{Go}, I_G, I_{RR}, I_{KD}\}$, then the SISO fuzzy controller based on universal triple I method is approximately a step response function.*

Remark 5.12. When $\rightarrow_2 \in \{I_{FD}, I_L, I_{Go}, I_G, I_{RR}, I_M\}$, the related results (e.g., Theorems 5.4, 5.6, 5.8, 5.9 and 5.10) are also correct, since $I_{FD}, I_L, I_{Go}, I_G, I_{RR}$ are $(0,1)$ -implications, and the case of $\rightarrow_2 = I_M$ is similar to the one of $\rightarrow_2 = I_G$ (which has the same expression of Min-solution).

5.2. DISO Fuzzy Controllers Based on Universal Triple I Methods. Let X, Y be the universes of input variables and Z the universe of output variable. Denote $\mathbb{A} = \{A_i\}_{(1 \leq i \leq n)}$, $\mathbb{B} = \{B_i\}_{(1 \leq i \leq n)}$ and $\mathbb{C} = \{C_i\}_{(1 \leq i \leq n)}$, where $A_i \in F(X)$, $B_i \in F(Y)$, $C_i \in F(Z)$. $\mathbb{A}, \mathbb{B}, \mathbb{C}$ are regarded as linguistic variables, thus the fuzzy reasoning rules can be expressed as follows:

$$\text{If } x \text{ is } A_i \text{ and } y \text{ is } B_i, \text{ then } z \text{ is } C_i, \quad i = 1, \dots, n, \quad (17)$$

where $x \in X, y \in Y, z \in Z$ are called base variables. Similarly, the inference relation of i -th inference rule can be changed into $(A_i(x) \wedge B_i(y)) \rightarrow_1 C_i(z)$, and we get the whole inference rule

$$IR(x, y, z) \triangleq \bigvee_{i=1}^n ((A_i(x) \wedge B_i(y)) \rightarrow_1 C_i(z)).$$

For a DISO fuzzy controller, the input value is a crisp quantity $(x^*, y^*) \in X \times Y$. We treat (x^*, y^*) by singleton fuzzification, and get $A_{x^*}^* = \begin{cases} 1, & x = x^* \\ 0, & x \neq x^* \end{cases}$, $B_{y^*}^* = \begin{cases} 1, & y = y^* \\ 0, & y \neq y^* \end{cases}$. Then we achieve C^* by universal triple I method of fuzzy reasoning from the input $A_{x^*}^*$ and $B_{y^*}^*$, where (12) should be turned into:

$$IR(x, y, z) \rightarrow_2 ((A^*(x) \wedge B^*(y)) \rightarrow_2 C^*(z)). \quad (18)$$

Lastly, to defuzzify C^* , we adopt the method of centroid, that is,

$$z^* = \int_Z z C^*(z) dz / \int_Z C^*(z) dz.$$

Therefore, there is an output $z^* = G(x^*, y^*)$ for each input (x^*, y^*) . Then a DISO fuzzy controller based on universal triple I method is achieved.

To analyze interpolation mechanism of DISO fuzzy controller, suppose that $\mathbb{A}, \mathbb{B}, \mathbb{C}$ are respectively the fuzzy partitions of X, Y and Z (where A_i, B_i, C_i are integrable functions). We assume that X, Y and Z are all real number intervals, e.g. $X = [a, b]$, $Y = [c, d]$ and $Z = [e, f]$ where $a < x_1 < x_2 < \dots < x_n < b$, $c < y_1 < y_2 < \dots < y_n < d$ and $e < z_1 < z_2 < \dots < z_n < f$, in which x_i, y_i, z_i are respectively peak-points of A_i, B_i, C_i .

Let $h_1 = z_1 - e$, $h_i = z_i - z_{i-1}$ ($i = 2, 3, \dots, n$) and $h = \max_{1 \leq i \leq n} \{h_i\}$. Since \mathbb{A}, \mathbb{B} and \mathbb{C} are all fuzzy partitions, they have Kronecker property: $A_i(x_j) = B_i(y_j) = C_i(z_j) = \delta_{ij}$ ($i, j = 1, \dots, n$). By the definition of definite integral, we obtain for the method of centroid:

$$z^* = \frac{\int_Z z C^*(z) dz}{\int_Z C^*(z) dz} \approx \frac{\sum_{i=1}^n z_i C^*(z_i) h_i}{\sum_{i=1}^n C^*(z_i) h_i}. \quad (19)$$

It is similar to Proposition 5.3 that Proposition 5.13 can be obtained.

Proposition 5.13. *Suppose that \rightarrow_2 is a $(0,1)$ -implication, and that T the function residual to \rightarrow_2 , then the Min-solution derived from (18) can be expressed as follows:*

$$C^*(z) = \sup_{(x,y) \in X \times Y} \{T((A^*(x) \wedge B^*(y)), IR(x, y, z))\}, \quad z \in Z.$$

Theorem 5.14. *Let \rightarrow_2 be a $(0,1)$ -implication, then the Min-solution $C^*(z) = IR(x^*, y^*, z)$ for the DISO fuzzy controller based on universal triple I method.*

Proof. Let \rightarrow_2 be a $(0,1)$ -implication. Then there uniquely exists T which is the residual function w.r.t. \rightarrow_2 , and $T(1, b) = b$, $T(0, b) = 0$ ($b \in [0, 1]$). It follows from Proposition 5.13 that the Min-solution is $C^*(z) = \sup_{(x,y) \in X \times Y} \{T((A^*(x) \wedge B^*(y)), IR(x, y, z))\}$, $z \in Z$. As for input (x^*, y^*) , we get $A_{x^*}^*$ and $B_{y^*}^*$. Because $T(1, b) = b$, $T(0, b) = 0$ hold for any $b \in [0, 1]$, we have: $T((A^*(x) \wedge B^*(y)), IR(x, y, z)) = T(1, IR(x^*, y^*, z)) = IR(x^*, y^*, z)$ for the case of $(x, y) = (x^*, y^*)$, and $T((A^*(x) \wedge B^*(y)), IR(x, y, z)) = T(0, IR(x, y, z)) = 0$ for the case of $(x, y) \in X \times Y - \{(x^*, y^*)\}$. Therefore $C^*(z) = IR(x^*, y^*, z)$. \square

Similar to Lemma 5.5, we can prove Lemma 5.15.

Lemma 5.15. *In the DISO fuzzy controller based on universal triple I method,*

- (i) *if \rightarrow_1 is an expansion type operator, then $IR(x^*, y^*, z_j) > 0$ for any $(x^*, y^*) \in X \times Y$ ($j = 1, \dots, n$);*
- (ii) *if \rightarrow_1 is a reduction type operator satisfying (C13), then there exists z_j such that $IR(x^*, y^*, z_j) > 0$ for any $(x^*, y^*) \in X \times Y$ ($j \in \{1, \dots, n\}$);*
- (iii) *if \rightarrow_1 is an other type operator satisfying (C14), then $IR(x^*, y^*, z_j) > 0$ for any $(x^*, y^*) \in X \times Y$ ($j = 1, \dots, n$);*
- (iv) *if \rightarrow_1 is an R-implication, an S-implication or a QL-implication, then $IR(x^*, y^*, z_j) > 0$ for any $(x^*, y^*) \in X \times Y$ ($j = 1, \dots, n$).*

Theorem 5.16. *Let \rightarrow_2 be a $(0,1)$ -implication. If \rightarrow_1 is a reduction type operator satisfying (C13), then there exists a group of base functions $\Phi = \{\varphi_i\}_{(1 \leq i \leq n)}$ such that the DISO fuzzy controller based on universal triple I method is approximately a binary piecewise interpolation function taking φ_i as its base functions, i.e. $G(x, y) = \sum_{i=1}^n \varphi_i(x, y) z_i$.*

Proof. Note that \rightarrow_2 is a $(0,1)$ -implication, it follows from Theorem 5.14 that Min-solution $C^*(z) = IR(x^*, y^*, z) = \bigvee_{k=1}^n ((A_k(x^*) \wedge B_k(y^*)) \rightarrow_1 C_k(z))$.

Since \rightarrow_1 is a reduction type operator satisfying (C13), then $a \rightarrow_1 1 = a$, $a \rightarrow_1 0 = 0$. Noting $C_k(z_i) = \delta_{ki}$ ($i, k = 1, \dots, n$), it follows from (19) that

$$\begin{aligned} z^* &\approx \frac{\sum_{i=1}^n z_i C^*(z_i) h_i}{\sum_{i=1}^n C^*(z_i) h_i} = \frac{\sum_{i=1}^n z_i [\bigvee_{k=1}^n ((A_k(x^*) \wedge B_k(y^*)) \rightarrow_1 C_k(z_i))] h_i}{\sum_{i=1}^n [\bigvee_{k=1}^n ((A_k(x^*) \wedge B_k(y^*)) \rightarrow_1 C_k(z_i))] h_i} \\ &= \frac{\sum_{i=1}^n z_i (A_i(x^*) \wedge B_i(y^*)) h_i}{\sum_{i=1}^n (A_i(x^*) \wedge B_i(y^*)) h_i}, \end{aligned} \quad (20)$$

where there exists z_i such that $C^*(z_i) = IR(x^*, y^*, z_i) > 0$ ($i \in \{1, \dots, n\}$) by Lemma 5.15(ii), so $\sum_{i=1}^n C^*(z_i) h_i > 0$ and then (20) makes sense.

Denote

$$C_i(x^*, y^*) \triangleq A_i(x^*) \wedge B_i(y^*), \quad \varphi_i(x^*, y^*) \triangleq h_i C_i(x^*, y^*) / (\sum_{i=1}^n h_i C_i(x^*, y^*)),$$

then we get $z^* \approx \sum_{i=1}^n \varphi_i(x^*, y^*) z_i$. Let $\Phi \triangleq \{\varphi_i\}_{(1 \leq i \leq n)}$, $G(x, y) \triangleq \sum_{i=1}^n \varphi_i(x, y) z_i$. Considering $A_k(x_i) = B_k(y_i) = \delta_{ki}$ ($i, k = 1, \dots, n$), we get

$$G(x_i, y_i) = \frac{\sum_{k=1}^n z_k (A_k(x_i) \wedge B_k(y_i)) h_k}{\sum_{k=1}^n (A_k(x_i) \wedge B_k(y_i)) h_k} = \frac{z_i h_i}{h_i} = z_i, \quad i = 1, \dots, n,$$

then $G(x, y)$ is a binary piecewise interpolation function. \square

Corollary 5.17. *Let \rightarrow_2 be a (0,1)-implication. If $\rightarrow_1 \in \{I_M, I_{La}\}$, then there exists a group of base functions $\Phi = \{\varphi_i\}_{(1 \leq i \leq n)}$ such that the DISO fuzzy controller based on universal triple I method is approximately a binary piecewise interpolation function taking φ_i as its base functions.*

Similarly Theorems 5.18, 5.19, 5.20 and Corollary 5.21 can be obtained.

Theorem 5.18. *Let \rightarrow_2 be a (0,1)-implication. If $\rightarrow_1 = I_Z$, then there exists a group of base functions $\Phi = \{\varphi_i\}_{(1 \leq i \leq n)}$ such that the DISO fuzzy controller based on universal triple I method is approximately a binary piecewise fitted function regarding φ_i as its base functions, i.e. $G(x, y) = \sum_{i=1}^n \varphi_i(x, y) z_i$.*

Theorem 5.19. *Let \rightarrow_2 be a (0,1)-implication. If \rightarrow_1 is an expansion type operator, then the DISO fuzzy controller based on universal triple I method is approximately a step response function.*

Theorem 5.20. *Let \rightarrow_2 be a (0,1)-implication. If \rightarrow_1 is an R-implication or S-implication, then the DISO fuzzy controller based on universal triple I method is approximately a step response function.*

Corollary 5.21. *Let \rightarrow_2 be a (0,1)-implication. If $\rightarrow_1 \in \{I_{FD}, I_L, I_{Go}, I_G, I_{RR}, I_{KD}\}$, then the DISO fuzzy controller based on universal triple I method is approximately a step response function.*

Remark 5.22. Similar to Remark 5.12, when $\rightarrow_2 \in \{I_{FD}, I_L, I_{Go}, I_G, I_{RR}, I_M\}$, the related results (e.g., Theorems 5.14, 5.16, 5.18, 5.19 and 5.20) are also right.

Remark 5.23. Since an R-implication is a (0,1)-implication, the related results in this section is also applicable to the case that \rightarrow_2 takes an R-implication.

Remark 5.24. When \rightarrow_2 takes I_M , the universal triple I method degenerates into the CRI method, thus we can easily get the response ability of SISO and DISO fuzzy controllers based on the CRI method (e.g., from Remarks 5.12, 5.22, Theorems 5.6, 5.8, 5.9, 5.16, 5.18 and 5.19). Obviously, there are more usable fuzzy controllers based on the universal triple I method than the ones based on the CRI method.

Remark 5.25. The response ability of fuzzy controllers based on universal triple I methods can be divided into 3 kinds as follows:

(i) If \rightarrow_2 is a (0,1)-implication and \rightarrow_1 is a reduction type operator satisfying (C13), then the fuzzy controller based on universal triple I method is approximately an interpolation function, thus it can be universal approximator and then usable in practice.

(ii) If \rightarrow_2 is a (0,1)-implication, and \rightarrow_1 takes I_Z (noting that I_Z is an other type operator, and also a typical QL-implication), then the fuzzy controller based on universal triple I method is approximately a fitted function, hence it may be usable.

(iii) If \rightarrow_2 is a (0,1)-implication, and \rightarrow_1 is an expansion type operator (including R-implication or S-implication), then the fuzzy controller based on universal triple I method is approximately a step response function, thus it only has step response ability, therefore it can hardly be used in practice.

6. Several Discussions of the Universal Triple I Method

To begin with, we shall discuss the important value of generalization from the triple I method to the universal triple I method. The reversibility property of universal triple I method is totally determined by the second and third implications in (3) (corresponding to \rightarrow_2 in (4)) if $\rightarrow_2 \in \{I_{KD}, I_{La}\}$; and is unitedly determined by \rightarrow_1 and \rightarrow_2 , if \rightarrow_2 is a (0,1)-implication or $\rightarrow_2 \in \{I_M, I_Z\}$ (see e.g., Theorems 4.3, 4.4, 4.14 and 4.16). Therefore, it is reasonable to let the first implication be \rightarrow_1 , and the second and third implications be \rightarrow_2 (i.e., generalize (3) to (4)).

Furthermore, when \rightarrow_1 and \rightarrow_2 are allowed to take different implications, more usable fuzzy controllers are achieved. For example, it follows from Remark 5.25 that we obtain 18 usable fuzzy controllers based on universal triple I method (in which $(\rightarrow_1, \rightarrow_2) \in \{I_M, I_{La}, I_Z\} \times \{I_{FD}, I_L, I_{Go}, I_G, I_{RR}, I_M\}$). In fact, if we get more (0,1)-implications (or reduction type operators satisfying (C13)), we can achieve more usable fuzzy controllers. However, we get from [6, 15] that there are only 2 usable fuzzy controllers based on the triple I method. Therefore, the practicability of the universal triple I method is superior to the triple I method.

To sum up, it is of significance to generalize the triple I method to the universal triple I method.

At last, we shall analyze the duty of first implication \rightarrow_1 and second implication \rightarrow_2 , together with how to choose \rightarrow_1 and \rightarrow_2 . It is easy to know that the form of universal triple I solution is basically determined only if \rightarrow_2 is chosen (i.e. \rightarrow_2 takes an implication), and hence \rightarrow_2 determines the reasoning mechanism to a large extent (see e.g. Theorem 3.5, Lemma 3.7 and Proposition 5.3). Meanwhile, \rightarrow_1 often exists as the form of $(A(x) \rightarrow_1 B(y))$ (or $IR(x, y)$ and so on) embodying the function of rule base (see e.g. Theorem 3.5, Propositions 5.3 and 5.13). Summarizing above, the second implication and first implication respectively embody the reasoning mechanism and function of rule base. What is more, the second implication has leading status in virtue of its effect on direction of inference.

Note that a universal triple I method is decided if \rightarrow_1 and \rightarrow_2 are chosen. Naturally, how to reasonably choose \rightarrow_1 and \rightarrow_2 becomes a key research topic in the universal triple I method. We will not discuss it here in detail, but give some instructional principles. From the results mentioned above, when \rightarrow_2 takes the

(0,1)-implication (or more strict R-implication), the universal triple I method has good property from the viewpoints of reversibility property and response ability, thus \rightarrow_2 can prefer to be the (0,1)-implication (or R-implication). Besides, it is natural to let the universal triple I method possess both the reversibility property and good response ability to the greatest extent for choosing \rightarrow_1 and \rightarrow_2 .

7. Conclusions

The universal triple I method is investigated from the viewpoints of both fuzzy reasoning and fuzzy controllers. The main contributions are as follows.

(i) The universal triple I principle is brought forward, which improves the previous triple I principle. Then, unified form of universal triple I method is established (to allow different implications to be employed in the same manner), where \rightarrow_2 takes a (0,1)-implication or an R-implication. The CRI method and the triple I method can be regarded as special cases of the universal triple I method.

(ii) The reversibility property of universal triple I method is analyzed from expansion, reduction and other type operators. It is found that the universal triple I method can be reversible so long as we appropriately choose \rightarrow_1 and \rightarrow_2 , and the reversibility property seems excellent if \rightarrow_2 takes a (0,1)-implication.

(iii) We investigate the response ability of fuzzy controllers based on universal triple I methods. When \rightarrow_2 takes a (0,1)-implication, and \rightarrow_1 takes I_Z or a reduction type operator satisfying (C13), the corresponding fuzzy controller can be practicable. There are more usable fuzzy controllers based on the universal triple I method than the ones based on the CRI method or the triple I method.

(iv) It is pointed out that, in universal triple I method, \rightarrow_1 and \rightarrow_2 respectively embody the function of rule base and the reasoning mechanism. Moreover, it is suggested that \rightarrow_2 should prefer to take the (0,1)-implication (or R-implication).

In the universal triple I method, how do we reasonably choose \rightarrow_1 and \rightarrow_2 ? There is only preparatory research involving it in this paper. Moreover, for the fuzzy controllers based on universal triple I methods, we analyze the case that the combination operator of inference rules takes 'max' operator, then how about the 'min' operator (where the whole inference rule is $\bigwedge_{i=1}^n (A_i(x) \rightarrow_1 B_i(y))$)? These problems will be investigated in the further research.

Furthermore, the properties related to universal triple I method (and corresponding fuzzy controllers), such as continuity, robustness as well as stability (see [9, 13, 16]), are also vital topics, which will be our research emphases in the future.

Acknowledgements. Authors would like to express their sincere thanks to the valuable comments and suggestions from editors and reviewers.

REFERENCES

- [1] S. M. Chen, Y. K. Ko, Y. C. Chang and J. S. Pan, *Weighted fuzzy interpolative reasoning based on weighted increment transformation and weighted ratio transformation techniques*, IEEE Transactions on Fuzzy Systems, **17(6)** (2009), 1412–1427.
- [2] J. C. Fodor, *On contrapositive symmetry of implications in fuzzy logic*, First European Congress on Fuzzy and Intelligent Technologies, Aachen, (1993), 1342–1348.

- [3] S. Gottwald, *A treatise on many-valued logics*, Research Studies Press, Baldock, 2001.
- [4] P. Hájek, *Metamathematics of fuzzy logic*, Kluwer Academic Publishers, Dordrecht, 1998.
- [5] J. Hou and H. X. Li, *Reductivity of some fuzzy inference methods*, Fuzzy Systems and Mathematics, **19(4)** (2005), 90–95.
- [6] J. Hou, F. You and H. X. Li, *Fuzzy systems constructed by triple I algorithm and their response ability*, Progress in Natural Science, **15(1)** (2005), 29–37.
- [7] S. Kirindis and V. Chatzis, *A robust fuzzy local information C-means clustering algorithm*, IEEE Transactions on Image Processing, **19(5)**(2010), 1328–1337.
- [8] E. P. Klement, R. Mesiar and E. Pap, *Triangular Norms*, Kluwer Academic Publishers, Dordrecht, 2000.
- [9] H. K. Lam and M. Narimani, *Quadratic-stability analysis of fuzzy-model-based control systems using staircase membership functions*, IEEE Transactions on Fuzzy Systems, **18(1)** (2010), 125–137.
- [10] H. X. Li, *Interpolation mechanism of fuzzy control*, Science in China (Series E), **41(3)** (1998), 312–320.
- [11] H. X. Li, *Probability representations of fuzzy systems*, Science in China (Series F), **49(3)** (2006), 339–363.
- [12] H. X. Li and E. S. Lee, *Interpolation representations of fuzzy logic systems*, Computers and Mathematics with Applications, **45(10)** (2003), 1683–1693.
- [13] D. C. Li, Y. M. Li and Y. J. Xie, *Robustness of interval-valued fuzzy inference*, Information Sciences, **181(20)** (2011), 4754–4764.
- [14] H. X. Li, F. You and J. Y. Peng, *Fuzzy controllers based on some fuzzy implication operators and their response functions*, Progress in Natural Science, **14(1)** (2004), 15–20.
- [15] H. X. Li, J. Y. Peng, J. Y. Wang, J. Hou and Y. Z. Zhang, *Fuzzy systems based on triple I algorithm and their response ability*, Journal of Systems Science and Complexity, **26(5)** (2006), 578–590.
- [16] H. W. Liu and G. J. Wang, *Continuity of triple I methods based on several implications*, Computers and Mathematics with Applications, **56(8)** (2008), 2079–2087.
- [17] X. P. Liu, Y. M. Tang, G. T. Shen and X. Chen, *A formal model of collaborative discussion for problem-solving*, Chinese Journal of Electronics, **21(3)** (2012), 453–459.
- [18] M. Mas, M. Monserrat, J. Torrens and E. Trillas, *A survey on fuzzy implication functions*, IEEE Transactions on Fuzzy Systems, **15(1)** (2007), 1107–1121.
- [19] J. M. Mendel, *Fuzzy logic systems for engineering: a tutorial*, Proceedings of the IEEE, **83(3)** (1995), 345–377.
- [20] V. Novák, I. Perfilieva and J. Močkoř, *Mathematical principles of fuzzy logic*, Kluwer Academic Publishers, Boston, 1999.
- [21] D. W. Pei, *Ideal implications in fuzzy logic and fuzzy control*, Journal of Xi'an Petroleum Institute, **15(6)** (2000), 44–47.
- [22] D. W. Pei, *R_0 implication: characteristics and applications*, Fuzzy Sets and Systems, **131(3)** (2002), 297–302.
- [23] D. W. Pei, *Full implication triple I algorithms and their consistency in fuzzy reasoning*, Journal of Mathematical Research and Exposition, **24(2)** (2004), 359–368.
- [24] D. W. Pei, *Unified full implication algorithms of fuzzy reasoning*, Information Sciences, **178(2)** (2008), 520–530.
- [25] J. Y. Peng, H. X. Li, J. Hou, F. You and J. Y. Wang, *Fuzzy controllers based on pointwise optimization fuzzy inference and its interpolation mechanism*, Journal of Systems Science and Complexity, **25(3)** (2005), 311–322.
- [26] F. J. Ren, *Automatic abstracting important sentences*, International Journal of Information Technology and Decision Making, **4(1)** (2005), 141–152.
- [27] S. J. Song, C. B. Feng and E. S. Lee, *Triple I method of fuzzy reasoning*, Computers and Mathematics with Applications, **44(12)** (2002), 1567–1579.
- [28] Y. M. Tang and X. P. Liu, *Task partition for function tree according to innovative functional reasoning*, Twelfth International Conference on CSCWD, Xian, China, (2008), 189–195.

- [29] Y. M. Tang and X. P. Liu, *Fuzzy systems constructed by triple I method or CRI method and their response functions*, Journal of Hefei University of Technology, **33(2)** (2010), 182–187.
- [30] Y. M. Tang and X. P. Liu, *Differently implicational universal triple I method of (1, 2, 2) type*, Computers and Mathematics with Applications, **59(6)** (2010), 1965–1984.
- [31] Y. M. Tang, F. J. Ren, X. Sun and Y. X. Chen, *Reverse universal triple I method of (1,1,2) type for the Lukasiewicz implication*, Seventh Conference on NLPKE, Tokushima, Japan, (2011), 23–30.
- [32] I. B. Turksen and Y. Tian, *Combination of rules or their consequences in fuzzy expert systems*, Fuzzy Sets and Systems, **58(1)** (1993), 3–40.
- [33] G. J. Wang, *Fully implicational triple I method for fuzzy reasoning*, Science in China (Series E), **29(1)** (1999), 43–53.
- [34] G. J. Wang, *Non-classical mathematical logic and approximate reasoning*, Science in China Press, Beijing, 2000.
- [35] G. J. Wang, *Introduction to mathematical logic and resolution principle*, Science in China Press, Beijing, 2003.
- [36] G. J. Wang and L. Fu, *Unified forms of triple I method*, Computers and Mathematics with Applications, **49(5)** (2005), 923–932.
- [37] L. A. Zadeh, *Outline of a new approach to the analysis of complex systems and decision processes*, IEEE Transactions on Systems Man and Cybernetics, **3(1)** (1973), 28–44.
- [38] J. C. Zhang and X. Y. Yang, *Some properties of fuzzy reasoning in propositional fuzzy logic systems*, Information Sciences, **180(23)** (2010), 4661–4671.

YIMING TANG*, INFORMATION AND COMMUNICATION ENGINEERING POSTDOCTORAL RESEARCH STATION, SCHOOL OF COMPUTER AND INFORMATION, HEFEI UNIVERSITY OF TECHNOLOGY, HEFEI 230009, CHINA

E-mail address: tym608@163.com

FUJI REN, INSTITUTE OF TECHNOLOGY AND SCIENCE, THE UNIVERSITY OF TOKUSHIMA, MINAMI JOSANJIMA, TOKUSHIMA, 770-8506, JAPAN

E-mail address: ren2fuji@gmail.com

*CORRESPONDING AUTHOR