

THE CATEGORY OF \top -CONVERGENCE SPACES AND ITS CARTESIAN-CLOSEDNESS

Q. YU AND J. FANG

ABSTRACT. In this paper, we define a kind of lattice-valued convergence spaces based on the notion of \top -filters, namely \top -convergence spaces, and show the category of \top -convergence spaces is Cartesian-closed. Further, in the lattice valued context of a complete MV -algebra, a close relation between the category of \top -convergence spaces and that of strong L -topological spaces is established. In details, we show that the category of strong L -topological spaces is concretely isomorphic to that of strong L -topological \top -convergence spaces categorically and bireflectively embedded in that of \top -convergence spaces.

1. Introduction

As pointed out by E. Lowen and R. Lowen in [14], the category of stratified $[0, 1]$ -topological spaces (or fuzzy topological spaces in the original terminology of [13]) is not completely satisfactory for certain application in Algebra topology or Functional analysis, here $[0, 1]$ is the unital interval. The main reason is the fact that it is not Cartesian-closed and hence there is no natural function space for the sets of morphisms. In order to overcome this deficiency, by starting from convergence theory in stratified $[0, 1]$ -topological spaces developed by R. Lowen in [13], E. Lowen et al. [14, 15] considered fuzzy convergence spaces as a generalization of Choquet's convergence spaces [1] and obtained the resulting Cartesian-closed category containing the category of stratified $[0, 1]$ -topological spaces as a fully embedded subcategory.

For more general lattice L instead of the unital interval $[0, 1]$, stemming from stratified L -topological spaces, Jäger [10] developed a theory of convergence based on the notion of stratified L -filters, where L is a complete Heyting Algebra. The resulting category, namely the category of stratified L -generalized convergence spaces, has the desired structural property of Cartesian-closedness and contains the category of stratified L -topological spaces as an embedded reflective subcategory. The convergence theory was developed to a significant extent in recent years [3, 18, 19, 11, 12, 16, 20, 21]. On this basis, in the same lattice valued context, Fang [2] defined a subcategory of the category of stratified L -generalized convergence spaces, namely the category of stratified L -ordered convergence spaces, which also is Cartesian-closed and contains the category of stratified L -topological spaces as an embedded reflective subcategory.

Received: March 2016; Revised: September 2016; Accepted: November 2016

Key words and phrases: \top -filter, \top -convergence, Cartesian-closedness, Topological category, Reflection, Strong L -topology.

Thus in case of the underlying lattice L being a complete Heyting Algebra, we see that Jäger and Fang all got nice conclusions. Regretfully, the idempotency about the meet operation \wedge of L is indispensable for Cartesian-closedness of the resulting categories in [2, 10]. The requirement of idempotency is not very convenient because the semigroup operation in most of underlying lattices (e.g., complete MV -algebras) is not idempotent, but commutative.

Apart from the Cartesian-closedness depending on the idempotency of the meet operation \wedge , all convergence spaces mentioned above, such as fuzzy convergence space, stratified L -generalized convergence spaces and stratified L -ordered convergence spaces, start from a kind of stratified L -topological spaces. In fact, besides stratified L -topological spaces, there exists another kind of lattice-valued topological spaces, namely strong L -topological spaces introduced by Zhang [22], which actually are probabilistic topological spaces [8, 9] in a complete MV -algebra.

Hence when the underlying lattice is possessed of a semigroup operation with non-idempotency, it is necessary to find a kind of lattice-valued convergence spaces starting from strong L -topological spaces such that the resulting category is Cartesian-closed and contains the category of strong L -topological spaces as an embedded reflective subcategory.

By this paper, we try to propose a kind of lattice-valued convergence spaces based on the notion of \top -filters which was introduced by Höhle [9], namely \top -convergence spaces, and show that when the lattice theoretical setting is a complete MV -algebra, the category of \top -convergence spaces is Cartesian-closed and the idempotency of the semigroup operation is not required here. Further, we also want to establish a close relation between the category of \top -convergence spaces and that of strong L -topological spaces in case the lattice L is a complete MV -algebra. In fact, we will show that the category of strong L -topological \top -convergence spaces is concretely isomorphic to that of strong L -topological spaces categorically, and the category of \top -convergence spaces contains that of strong L -topological spaces as an embedded bireflective subcategory.

The paper is organized as follows: In Section 2, we provide the lattice theoretical context and recall some notions used in this paper. In Section 3, a concept of \top -convergence spaces is proposed and the category of \top -convergence spaces is introduced. Then after showing the category of \top -convergence spaces is topological, the Cartesian-closedness of the category of \top -convergence spaces is obtained. In Section 4, it is presented the relation between the category of \top -convergence spaces and that of strong L -topological spaces.

2. Preliminaries

A triple $(L, \leq, *)$ is called a *complete residuated lattice*, if (L, \leq) is a complete lattice with \top and \perp respectively being the top and the bottom element of L , and $*$: $L \times L \rightarrow L$, called a tensor on L , is a commutative, associative binary operation such that

- (1) $*$ is monotone on each variable,

- (2) For each $\alpha \in L$, the monotone mapping $\alpha * (-) : L \rightarrow L$ has a right adjoint $\alpha \rightarrow (-) : L \rightarrow L$ in the sense that $\alpha * \beta \leq \gamma \iff \beta \leq \alpha \rightarrow \gamma$ for all $\beta, \gamma \in L$,
- (3) The top element \top is a unit element for $*$, i.e. $\top * \alpha = \alpha$ for all $\alpha \in L$.

For a given complete residuated lattice, the binary operation \rightarrow on L can be computed by $\alpha \rightarrow \beta = \bigvee \{ \gamma \in L \mid \alpha * \gamma \leq \beta \}$ for all $\alpha, \beta \in L$. The binary operation \rightarrow is called the *implication* operation with respect to $*$. Some basic properties of the tensor $*$ and the implication operation \rightarrow are collected in the following lemma; they can be found in many works, for instance [4, 19, 22].

Lemma 2.1. *Let $(L, \leq, *)$ be a complete residuated lattice. Then for all $\alpha, \beta, \gamma, \delta \in L$, $\{\beta_i\}_{i \in I} \subseteq L$, the following conditions hold:*

- (a) $\top \rightarrow \alpha = \alpha$,
- (b) $\alpha * (\alpha \rightarrow \beta) \leq \beta$,
- (c) $\alpha \leq \beta$ if and only if $\alpha \rightarrow \beta = \top$,
- (d) $(\alpha \rightarrow \beta) * (\beta \rightarrow \gamma) \leq \alpha \rightarrow \gamma$,
- (e) $(\alpha \rightarrow \beta) * (\gamma \rightarrow \delta) \leq (\alpha * \gamma) \rightarrow (\beta * \delta)$ and $(\alpha \rightarrow \beta) \wedge (\gamma \rightarrow \delta) \leq (\alpha \wedge \gamma) \rightarrow (\beta \wedge \delta)$,
- (f) $\alpha * \bigvee_{i \in I} \beta_i = \bigvee_{i \in I} (\alpha * \beta_i)$,
- (g) $\alpha \rightarrow \bigwedge_{i \in I} \beta_i = \bigwedge_{i \in I} (\alpha \rightarrow \beta_i)$, hence $(\alpha \rightarrow \beta) \leq (\alpha \rightarrow \gamma)$ whenever $\beta \leq \gamma$,
- (h) $(\bigvee_{i \in I} \beta_i) \rightarrow \beta = \bigwedge_{i \in I} (\beta_i \rightarrow \beta)$, hence $(\alpha \rightarrow \beta) \geq (\gamma \rightarrow \beta)$ whenever $\alpha \leq \gamma$.

A complete residuated lattice $(L, \leq, *)$, denoted by L simply, is called a *complete MV-algebra*, if L satisfies the condition:

$$(MV) \quad \alpha \vee \beta = (\alpha \rightarrow \beta) \rightarrow \beta, \quad \forall \alpha, \beta \in L.$$

A canonical example is the unital interval $[0, 1]$ with the tensor $\alpha * \beta = \max\{\alpha + \beta - 1, 0\}$, which means that $([0, 1], \leq, *)$ is a complete MV-algebra such that 1 is the unital element and the implication \rightarrow with respect to $*$ is given by $\alpha \rightarrow \beta = \min\{1 - \alpha + \beta, 1\}$ for all $\alpha, \beta \in [0, 1]$.

Throughout this paper, we will assume L to be a complete MV-algebra although most of results are valid for more general lattice-valued cases.

In this paper, we will often use, without explicitly mentioning, the following properties of a complete MV-algebra.

Lemma 2.2. [7] *For all $\alpha \in L$ and $\{\beta_j\}_{j \in J} \subseteq L$, then the following properties are valid:*

- (M1) $\alpha \wedge (\bigvee_{j \in J} \beta_j) = \bigvee_{j \in J} (\alpha \wedge \beta_j)$,
- (M2) $\alpha \vee (\bigwedge_{j \in J} \beta_j) = \bigwedge_{j \in J} (\alpha \vee \beta_j)$,
- (M3) $\alpha * (\bigwedge_{j \in J} \beta_j) = \bigwedge_{j \in J} (\alpha * \beta_j)$,
- (M4) $\alpha \rightarrow (\bigvee_{j \in J} \beta_j) = \bigvee_{j \in J} (\alpha \rightarrow \beta_j)$.

An L -subset on a set X is a map from X to L , and the family of all L -subsets on X will be denoted by L^X , called the L -power set of X . For any $x \in X$, $A(x)$ is interpreted as the degree to which x is in A . By 1_X and 0_X , we denote the constant L -subsets on X taking the value \top and \perp , respectively. We don't distinguish an

element $\alpha \in L$ and the constant function $\alpha : X \rightarrow L$ such that $\alpha(x) = \alpha$ for all $x \in X$. As usual, for a universal set X , the set of all subsets of X is denoted by $\mathcal{P}(X)$, called the *power set* of X .

All algebraic operations on L can be extended to the L -power set L^X pointwisely. That is, for all $A, B \in L^X$ and $x \in X$,

- (1) $(A \wedge B)(x) = A(x) \wedge B(x)$,
- (2) $(A \vee B)(x) = A(x) \vee B(x)$,
- (3) $(A * B)(x) = A(x) * B(x)$,
- (4) $(A \rightarrow B)(x) = A(x) \rightarrow B(x)$.

Let $\varphi : X \rightarrow Y$ be a map. Define $\varphi^\rightarrow : L^X \rightarrow L^Y$ and $\varphi^\leftarrow : L^Y \rightarrow L^X$ respectively by $\varphi^\rightarrow(A)(y) = \bigvee_{\varphi(x)=y} A(x)$ for all $A \in L^X$ and $y \in Y$, $\varphi^\leftarrow(B) = B \circ \varphi$

for all $B \in L^Y$.

For a set X , there exists a binary map $\mathcal{S}_X(-, -) : L^X \times L^X \rightarrow L$ defined by $\mathcal{S}_X(A, B) = \bigwedge_{x \in X} (A(x) \rightarrow B(x))$ for each pair $(A, B) \in L^X \times L^X$, where \rightarrow is the

implication operation corresponding to $*$. For all $A, B \in L^X$, $\mathcal{S}_X(A, B)$ can be interpreted as the degree to which A is a subset of B . It was called *fuzzy inclusion order* [22] or *subsethood degree* [4] of L -subsets.

Lemma 2.3. [2] *Let X and Y be nonempty sets. For any $A, B, C \in L^X$ and $E, F \in L^Y$, then the following statements hold:*

- (1) $A \leq B$ if and only if $\top = \mathcal{S}_X(A, B)$.
- (2) $\mathcal{S}_X(A, B) \leq \mathcal{S}_X(B, C) \rightarrow \mathcal{S}_X(A, C)$.
- (3) $\mathcal{S}_X(A, B \wedge C) = \mathcal{S}_X(A, B) \wedge \mathcal{S}_X(A, C)$ and $\mathcal{S}_X(B \vee C, A) = \mathcal{S}_X(B, A) \wedge \mathcal{S}_X(C, A)$, hence $\mathcal{S}_X(C, A) \leq \mathcal{S}_X(B, A)$ when $B \leq C$.
- (4) If $\varphi : X \rightarrow Y$ is a map, then

$$\mathcal{S}_X(A, B) \leq \mathcal{S}_Y(\varphi^\rightarrow(A), \varphi^\rightarrow(B)) \text{ and } \mathcal{S}_Y(E, F) \leq \mathcal{S}_X(\varphi^\leftarrow(E), \varphi^\leftarrow(F)).$$

J. Gutiérrez García and M.A. De Prada Vicente [5, 6] introduced the notion of characteristic value of a family of L -subsets extending that of characteristic value of a prefilter in [13] and provided the equivalent form of κ -condition [9]. Thereby they obtained the equivalent definitions of \top -filter and \top -filter base as follows.

Definition 2.4. [5, 6] Let X be a nonempty set. A \top -filter is a nonempty subset \mathbb{F} of L^X with the following properties:

- (F1) If $A \in L^X$ with $\bigvee_{C \in \mathbb{F}} \mathcal{S}_X(C, A) = \top$, then $A \in \mathbb{F}$,
- (F2) $A_1 \wedge A_2 \in \mathbb{F}$ for all $A_1, A_2 \in \mathbb{F}$,
- (F3) $\bigvee_{x \in X} A(x) = \top$ for all $A \in \mathbb{F}$.

The set of all \top -filters on X is denoted by $\mathcal{F}_L^\top(X)$.

Example 2.5. Let $[x]_\top = \{A \in L^X \mid A(x) = \top\}$ for given a point $x \in X$. Then $[x]_\top$ is a \top -filter, and called the *point \top -filter* of x . In case $X = \{x\}$, a single point set, $[x]_\top$ is the unique \top -filter on the X .

Definition 2.6. [6] A nonempty subset \mathbb{B} of L^X is a \top -filter base, if it satisfies the following conditions:

- ($\mathbb{B}1$) If $B_1, B_2 \in \mathbb{B}$, then $\bigvee_{B \in \mathbb{B}} \mathcal{S}_X(B, B_1 \wedge B_2) = \top$,
- ($\mathbb{B}2$) $\bigvee_{x \in X} B(x) = \top$ for all $B \in \mathbb{B}$.

Remark 2.7. [6] Every \top -filter base \mathbb{B} generates a \top -filter $\mathbb{F}_{\mathbb{B}}$ defined by

$$\mathbb{F}_{\mathbb{B}} = \{A \in L^X \mid \bigvee_{B \in \mathbb{B}} \mathcal{S}_X(B, A) = \top\}.$$

In this case, $\mathbb{B} \subseteq \mathbb{F}_{\mathbb{B}}$ holds and \mathbb{B} is called a base of $\mathbb{F}_{\mathbb{B}}$. We could know every \top -filter \mathbb{F} is a base of itself. And let \mathbb{B} be any base of a \top -filter \mathbb{F} , then $\mathbb{F} = \{A \in L^X \mid \bigvee_{B \in \mathbb{B}} \mathcal{S}_X(B, A) = \top\}$ is true.

Definition 2.8. [6] Let $\varphi : X \rightarrow Y$ be a map, $\mathbb{F} \in \mathcal{F}_L^{\top}(X)$, $\mathbb{G} \in \mathcal{F}_L^{\top}(Y)$.

- (1) A \top -filter $\varphi^{\rightarrow}(\mathbb{F})$ on Y generated by the \top -filter base

$$\{\varphi^{\rightarrow}(B) \in L^Y \mid B \in \mathbb{F}\}$$

is called the *image* of \mathbb{F} under φ .

- (2) If the class $\{\varphi^{\leftarrow}(C) \in L^X \mid C \in \mathbb{G}\}$ satisfies $\bigvee_{y \in \varphi(X)} C(y) = \top$ for all $C \in \mathbb{G}$,

then $\{\varphi^{\leftarrow}(C) \in L^X \mid C \in \mathbb{G}\}$ is a \top -filter base on X and a \top -filter on X generated by it is called the *inverse image* of \mathbb{G} under φ , denoted by $\varphi^{\leftarrow}(\mathbb{G})$. In this case, we say the inverse image $\varphi^{\leftarrow}(\mathbb{G})$ exists sometimes.

Remark 2.9. [6] Let $\varphi : X \rightarrow Y$ be a map, $\mathbb{F} \in \mathcal{F}_L^{\top}(X)$ and $\mathbb{G} \in \mathcal{F}_L^{\top}(Y)$.

- (1) By Remark 2.7, we have

$$\varphi^{\rightarrow}(\mathbb{F}) = \{C \in L^Y \mid \bigvee_{B \in \mathbb{F}} \mathcal{S}_Y(\varphi^{\rightarrow}(B), C) = \top\}.$$

In fact, $\varphi^{\rightarrow}(\mathbb{F})$ has another expression $\{C \in L^Y \mid \varphi^{\leftarrow}(C) \in \mathbb{F}\}$.

- (2) From Definition 2.8(2), we know that the inverse image $\varphi^{\leftarrow}(\mathbb{G})$ doesn't always exist, but if $\varphi^{\leftarrow}(\mathbb{G})$ exists, we have the following expression

$$\varphi^{\leftarrow}(\mathbb{G}) = \{A \in L^X \mid \bigvee_{B \in \mathbb{G}} \mathcal{S}_X(\varphi^{\leftarrow}(B), A) = \top\}.$$

- (3) The following are satisfied:

- (i) if $\mathbb{F} \subseteq \mathbb{H}$ for $\mathbb{H} \in \mathcal{F}_L^{\top}(X)$, then $\varphi^{\rightarrow}(\mathbb{F}) \subseteq \varphi^{\rightarrow}(\mathbb{H})$,
- (ii) $\varphi^{\leftarrow} \circ \varphi^{\rightarrow}(\mathbb{F}) \subseteq \mathbb{F}$.

For more information on the categorical terminology we refer the reader to [17]. By a category we mean a construct \mathcal{C} whose objects are structured sets, i.e. pairs (X, ξ) where X is a set and ξ a \mathcal{C} -structure on X , whose morphisms $\varphi : (X, \xi) \rightarrow (Y, \eta)$ are suitable maps from X to Y and whose composition is the usual composition of maps. The forgetful functors will not be mentioned explicitly. We simply write \mathbf{X} for a categorical object (X, ξ) sometimes.

Definition 2.10. [17] A category \mathcal{C} is said to be *topological* if the following conditions are satisfied:

- (1) *Existence of initial structures:* For any set X , any family $((X_i, \xi_i))_{i \in I}$ of \mathcal{C} -objects indexed by a class I and any family $(f_i : X \rightarrow X_i)_{i \in I}$ of maps indexed by I there exists a unique \mathcal{C} -structure ξ on X which is *initial* with respect to $(f_i : X \rightarrow X_i)_{i \in I}$ in the sense that for any \mathcal{C} -object (Y, η) , a map $g : (Y, \eta) \rightarrow (X, \xi)$ is a \mathcal{C} -morphism iff for every $i \in I$ the composite map $f_i \circ g : (Y, \eta) \rightarrow (X_i, \xi_i)$ is a \mathcal{C} -morphism.
- (2) *Fibre-smallness:* For any set X , the class $\{\xi \mid (X, \xi) \text{ is a } \mathcal{C}\text{-object}\}$ of all \mathcal{C} -structures with the underlying set X , called *\mathcal{C} -fibre* of X , is a set.
- (3) *Terminal separator property:* For any set X with cardinality at most one, there exists exactly one \mathcal{C} -structure on X .

Definition 2.11. [17] A category \mathcal{C} is said to be *Cartesian-closed* provided that the following conditions are satisfied:

- (1) For each pair (\mathbf{X}, \mathbf{Y}) of \mathcal{C} -objects there exists a product $\mathbf{X} \times \mathbf{Y}$ in \mathcal{C} .
- (2) For any \mathcal{C} -objects \mathbf{X} and \mathbf{Y} , there exists some \mathcal{C} -object $\mathbf{Y}^{\mathbf{X}}$ (called *power object*) and some \mathcal{C} -morphism $ev_{\mathbf{X}, \mathbf{Y}} : \mathbf{Y}^{\mathbf{X}} \times \mathbf{X} \rightarrow \mathbf{Y}$ (called *evaluation morphism*) such that for each \mathcal{C} -object \mathbf{Z} and each \mathcal{C} -morphism $\varphi : \mathbf{Z} \times \mathbf{X} \rightarrow \mathbf{Y}$, there exists a unique \mathcal{C} -morphism $\varphi^* : \mathbf{Z} \rightarrow \mathbf{Y}^{\mathbf{X}}$ such that $ev_{\mathbf{X}, \mathbf{Y}} \circ (\varphi^* \times id_{\mathbf{X}}) = \varphi$.

Definition 2.12. [17] Let \mathcal{A} be a subcategory of a category \mathcal{C} . \mathcal{A} is said to be *reflective* in \mathcal{C} provided that for each $\mathbf{X} \in |\mathcal{C}|$ there exists an \mathcal{A} -object $\mathbf{X}_{\mathcal{A}}$ and a \mathcal{C} -morphism $\gamma_{\mathbf{X}} : \mathbf{X} \rightarrow \mathbf{X}_{\mathcal{A}}$ such that for each \mathcal{A} -object \mathbf{Y} and each \mathcal{C} -morphism $\varphi : \mathbf{X} \rightarrow \mathbf{Y}$ there is a unique \mathcal{A} -morphism $\bar{\varphi} : \mathbf{X}_{\mathcal{A}} \rightarrow \mathbf{Y}$ such that $\varphi = \bar{\varphi} \circ \gamma_{\mathbf{X}}$. If the \mathcal{C} -morphism $\gamma_{\mathbf{X}} : \mathbf{X} \rightarrow \mathbf{X}_{\mathcal{A}}$ is bimorphism, then \mathcal{A} is said to be *bireflective* in \mathcal{C} , and $\gamma_{\mathbf{X}}$ is called *bireflection*.

3. The Cartesian-closedness of \top -Conv

In this section, we define a kind of lattice-valued convergence spaces based on the notion of \top -filter, namely \top -convergence spaces. The class of all \top -convergence spaces and continuous maps forms a category. We prove the category is topological and Cartesian-closed which are very nice structural properties.

Definition 3.1. Let X be a nonempty set. A map $\lim : \mathcal{F}_L^{\top}(X) \rightarrow \mathcal{P}(X)$ satisfying the following conditions:

- (TC1) $\forall x \in X, x \in \lim[x]_{\top}$,
- (TC2) $\forall \mathbb{F}, \mathbb{G} \in \mathcal{F}_L^{\top}(X), \mathbb{F} \subseteq \mathbb{G} \Rightarrow \lim \mathbb{F} \subseteq \lim \mathbb{G}$,

is called a \top -convergence on X , and the pair (X, \lim) is called a \top -convergence space. The set of all \top -convergences on X is denoted by $C^{\top}(X)$. We say \mathbb{F} converges to x instead of $x \in \lim \mathbb{F}$.

A map $\varphi : (X, \lim^X) \rightarrow (Y, \lim^Y)$ between \top -convergence spaces is said to be *continuous* provided that $x \in \lim^X \mathbb{F}$ means $\varphi(x) \in \lim^Y \varphi^{\Rightarrow}(\mathbb{F})$ for all $x \in X$ and $\mathbb{F} \in \mathcal{F}_L^{\top}(X)$. The class of all \top -convergence spaces and continuous maps forms a category, which is denoted \top -Conv.

Theorem 3.2. *The category $\top\text{-Conv}$ is topological.*

Proof. Firstly, the existence of initial structures can be proved as follows. Let X be a nonempty set, $\{(X_j, \lim_j)\}_{j \in J}$ a family of \top -convergence spaces and $\{\varphi_j : X \rightarrow (X_j, \lim_j)\}_{j \in J}$ a family of maps. A structure map $\lim^X : \mathcal{F}_L^\top(X) \rightarrow \mathcal{P}(X)$ on X is defined by $\lim^X \mathbb{F} = \{x \in X \mid \varphi_j(x) \in \lim_j \varphi_j^\Rightarrow(\mathbb{F}), \forall j \in J\}$ for all $\mathbb{F} \in \mathcal{F}_L^\top(X)$. Then it is easy to verify that \lim^X is a \top -convergence on X , and we only check its property of being initial below.

Let (Y, \lim^Y) be a \top -convergence space and $\psi : Y \rightarrow X$ be a map. For any $y \in \lim^Y \mathbb{G}$, here $y \in Y$ and $\mathbb{G} \in \mathcal{F}_L^\top(Y)$, we can get

$$\begin{aligned} & \varphi_j \circ \psi : (Y, \lim^Y) \rightarrow (X_j, \lim_j) \text{ is continuous for every } j \in J \\ \iff & \varphi_j(\psi(y)) = \varphi_j \circ \psi(y) \in \lim_j (\varphi_j \circ \psi)^\Rightarrow(\mathbb{G}) = \lim_j \varphi_j^\Rightarrow(\psi^\Rightarrow(\mathbb{G})), \forall j \in J \\ \iff & \psi(y) \in \lim^X \psi^\Rightarrow(\mathbb{G}) \text{ (by the definition of } \lim^X) \\ \iff & \psi : (Y, \lim^Y) \rightarrow (X, \lim^X) \text{ is continuous.} \end{aligned}$$

Next, since the class of all \top -convergences on X belongs to the set $\mathbf{2}^{\mathcal{P}(X)^{\mathcal{F}_L^\top(X)}}$, here $\mathbf{2} = \{0, 1\}$, the $\top\text{-Conv}$ -fibre of X is a set.

Finally, let $X = \{x\}$ be a singleton. $\mathcal{F}_L^\top(X) = \{[x]_\top\}$ holds by Example 2.5. Then the structure map $\lim : \mathcal{F}_L^\top(X) \rightarrow \mathcal{P}(X)$ is only determined by $\lim[x]_\top = \{x\}$. Thus \lim is the unique \top -convergence on X .

From all above, we get that the category $\top\text{-Conv}$ is topological. \square

Every topological category has products [17], so the condition (1) in Definition 2.11 is automatically fulfilled by the category of \top -convergence spaces.

Let (X, \lim^X) and (Y, \lim^Y) be \top -convergence spaces and $p_X : X \times Y \rightarrow X$ and $p_Y : X \times Y \rightarrow Y$ be the projection maps. The product of (X, \lim^X) and (Y, \lim^Y) is denoted by $(X \times Y, \lim^X \times \lim^Y)$ explicitly. Of course, for any $\mathbb{F} \in \mathcal{F}_L^\top(X \times Y)$, $(x, y) \in (\lim^X \times \lim^Y) \mathbb{F}$ if and only if $x \in \lim^X p_X^\Rightarrow(\mathbb{F})$ and $y \in \lim^Y p_Y^\Rightarrow(\mathbb{F})$.

The set of all continuous mappings from (X, \lim^X) to (Y, \lim^Y) is denoted by $C_\top(X, Y)$. In the category **Set** of sets and maps, there exists the evaluation map $ev_{X,Y} : C_\top(X, Y) \times X \rightarrow Y$ defined by $ev_{X,Y}(\varphi, x) = \varphi(x)$ for all $(\varphi, x) \in C_\top(X, Y) \times X$. In order to explore the Cartesian-closedness of $\top\text{-Conv}$, we need some lemmas and propositions in preparation for it.

Lemma 3.3. *Let $\mathbb{F}_i \in \mathcal{F}_L^\top(X_i)$ and \mathbb{B}_i be a base of \mathbb{F}_i , here $i = 1, 2$. Then*

$$\mathbb{B} = \{B_1 \times B_2 \mid B_1 \in \mathbb{B}_1 \text{ and } B_2 \in \mathbb{B}_2\}$$

is a \top -filter base, where for any $B_i \in \mathbb{B}_i$ ($i=1,2$),

$$B_1 \times B_2((x_1, x_2)) = B_1(x_1) \wedge B_2(x_2), \forall (x_1, x_2) \in X_1 \times X_2.$$

Proof. For any $A, C \in \mathbb{B}$, there exist $A_1, C_1 \in \mathbb{B}_1$ and $A_2, C_2 \in \mathbb{B}_2$ such that $A = A_1 \times A_2$ and $C = C_1 \times C_2$. $\bigvee_{B_i \in \mathbb{B}_i} \mathcal{S}_{X_i}(B_i, A_i) = \top$ ($i = 1, 2$) follows immediately from

Remark 2.7. From this, we observe that $\bigvee_{\substack{B_i \in \mathbb{B}_i \\ i=1,2}} \mathcal{S}_{X_1}(B_1, A_1) \wedge \mathcal{S}_{X_2}(B_2, A_2) = \top$. Since

for each $B_i \in \mathbb{B}_i$ ($i = 1, 2$), $\mathcal{S}_{X_1 \times X_2}(B_1 \times B_2, A_1 \times A_2) \geq \mathcal{S}_{X_1}(B_1, A_1) \wedge \mathcal{S}_{X_2}(B_2, A_2)$ holds owing to Lemma 2.1 (e), we conclude that

$$\bigvee_{\substack{B_i \in \mathbb{B}_i \\ i=1,2}} \mathcal{S}_{X_1 \times X_2}(B_1 \times B_2, A_1 \times A_2) = \top.$$

Certainly, $\bigvee_{\substack{B_i \in \mathbb{B}_i \\ i=1,2}} \mathcal{S}_{X_1 \times X_2}(B_1 \times B_2, C_1 \times C_2) = \top$. From all above, we obtain that

$$\begin{aligned} & \bigvee_{\substack{B_i \in \mathbb{B}_i \\ i=1,2}} \mathcal{S}_{X_1 \times X_2}(B_1 \times B_2, A \wedge C) \\ &= \bigvee_{\substack{B_i \in \mathbb{B}_i \\ i=1,2}} \mathcal{S}_{X_1 \times X_2}(B_1 \times B_2, (A_1 \times A_2) \wedge (C_1 \times C_2)) \\ &= \bigvee_{\substack{B_i \in \mathbb{B}_i \\ i=1,2}} \mathcal{S}_{X_1 \times X_2}(B_1 \times B_2, A_1 \times A_2) \wedge \mathcal{S}_{X_1 \times X_2}(B_1 \times B_2, C_1 \times C_2) \\ & \quad \text{(by using Lemma 2.3 (3))} \\ &= \top, \end{aligned}$$

i.e. \mathbb{B} satisfies (B1).

For any $B_1 \in \mathbb{B}_1$ and $B_2 \in \mathbb{B}_2$, we have

$$\bigvee_{\substack{x_i \in X_i \\ i=1,2}} B_1 \times B_2((x_1, x_2)) = \bigvee_{\substack{x_i \in X_i \\ i=1,2}} B_1(x_1) \wedge B_2(x_2) = \top$$

since $\bigvee_{x_i \in X_i} B_i(x_i) = \top$ for $i = 1, 2$ holds by \mathbb{B}_i satisfying (B2). Then the condition (B2) is satisfied by \mathbb{B} . \square

Let $\mathbb{F}_i \in \mathcal{F}_L^\top(X_i)$ for $i = 1, 2$. Because every \top -filter is a base of itself, from Lemma 3.3 above, $\{B_1 \times B_2 \mid B_i \in \mathbb{F}_i, i = 1, 2\}$ is a \top -filter base, which generates a \top -filter, denoted by $\mathbb{F}_1 \times \mathbb{F}_2$. And from Remark 2.7, $\mathbb{F}_1 \times \mathbb{F}_2$ can be determined by

$$\mathbb{F}_1 \times \mathbb{F}_2 = \{A \in L^{X_1 \times X_2} \mid \bigvee_{\substack{B_i \in \mathbb{F}_i \\ i=1,2}} \mathcal{S}_{X_1 \times X_2}(B_1 \times B_2, A) = \top\}.$$

Let $\varphi_i : X_i \rightarrow Y_i$ ($i=1,2$) be a map. By Remark 2.7 and Definition 2.8, we know $\{\varphi_i^\rightarrow(B_i) \in L^{Y_i} \mid B_i \in \mathbb{F}_i\}$ is a base of $\varphi_i^\rightarrow(\mathbb{F}_i)$. So

$$\{\varphi_1^\rightarrow(B_1) \times \varphi_2^\rightarrow(B_2) \mid B_i \in \mathbb{F}_i, i = 1, 2\}$$

is a \top -filter base from Lemma 3.3 and generates a \top -filter $\varphi_1^\rightarrow(\mathbb{F}_1) \times \varphi_2^\rightarrow(\mathbb{F}_2)$.

In addition, we could check that $(\varphi_1 \times \varphi_2)^\rightarrow(B_1 \times B_2) = \varphi_1^\rightarrow(B_1) \times \varphi_2^\rightarrow(B_2)$ is true. Then $\{(\varphi_1 \times \varphi_2)^\rightarrow(B_1 \times B_2) \mid B_i \in \mathbb{F}_i, i = 1, 2\}$ equals

$$\{\varphi_1^\rightarrow(B_1) \times \varphi_2^\rightarrow(B_2) \mid B_i \in \mathbb{F}_i, i = 1, 2\}$$

and is also a \top -filter base. And it generates a \top -filter $(\varphi_1 \times \varphi_2)^\rightarrow(\mathbb{F}_1 \times \mathbb{F}_2)$ by Definition 2.8. So we can get the following lemma.

Lemma 3.4. Let $\varphi : X_1 \rightarrow Y_1$ and $\psi : X_2 \rightarrow Y_2$ be two maps, $\mathbb{F}_1 \in \mathcal{F}_L^\top(X_1)$, $\mathbb{F}_2 \in \mathcal{F}_L^\top(X_2)$. Then $(\varphi \times \psi)^\rightarrow(\mathbb{F}_1 \times \mathbb{F}_2) = \varphi^\rightarrow(\mathbb{F}_1) \times \psi^\rightarrow(\mathbb{F}_2)$.

Lemma 3.5. Let $(X \times Y, \lim^X \times \lim^Y)$ be the product of two \top -Conv-objects (X, \lim^X) and (Y, \lim^Y) . If $p_X : X \times Y \rightarrow X$ and $p_Y : X \times Y \rightarrow Y$ be the projection maps, then for any $\mathbb{F} \in \mathcal{F}_L^\top(X)$ and $\mathbb{G} \in \mathcal{F}_L^\top(Y)$ the following are valid:

$$(1) p_X^\rightarrow(\mathbb{F} \times \mathbb{G}) = \mathbb{F}, \quad (2) p_Y^\rightarrow(\mathbb{F} \times \mathbb{G}) = \mathbb{G}.$$

Proof. In the proof, we only show the conclusion (1) for example. In general, $p_X^\rightarrow(A \times B) = A$ for each $A \in \mathbb{F}$, $B \in \mathbb{G}$ since for all $x \in X$,

$$p_X^\rightarrow(A \times B)(x) = \bigvee_{p_X(x,y)=x} A \times B((x,y)) = A(x) \wedge \bigvee_{y \in Y} B(y) = A(x) \wedge \top = A(x),$$

where $\bigvee_{y \in Y} B(y) = \top$ owing to $B \in \mathbb{G}$. Thus for each $C \in L^X$, $C \in p_X^\rightarrow(\mathbb{F} \times \mathbb{G})$ if and only if $\bigvee_{A \in \mathbb{F}, B \in \mathbb{G}} \mathcal{S}_X(p_X^\rightarrow(A \times B), C) = \top$, i.e. $\bigvee_{A \in \mathbb{F}} \mathcal{S}_X(A, C) = \top$, which is equivalent to $C \in \mathbb{F}$. Consequently, $p_X^\rightarrow(\mathbb{F} \times \mathbb{G}) = \mathbb{F}$. \square

Lemma 3.6 (Jäger [10]). If $\varphi \in C_\top(X, Y)$, then $\varphi^\rightarrow(A) = ev_{X,Y}^\rightarrow(1_\varphi \times A)$ holds for any $A \in L^X$, here $1_\varphi \in L^{C_\top(X,Y)}$ such that $1_\varphi(\varphi) = \top$ and $1_\varphi(\psi) = \perp$ for $\psi \neq \varphi$.

Now, we begin to confirm the existence of power object and the continuity of the evaluation map by the following propositions.

Proposition 3.7. Let (X, \lim^X) and (Y, \lim^Y) be two \top -Conv-objects. A map

$$\lim_C : \mathcal{F}_L^\top(C_\top(X, Y)) \rightarrow \mathcal{P}(C_\top(X, Y))$$

is defined for all $\mathbb{H} \in \mathcal{F}_L^\top(C_\top(X, Y))$, by

$$\lim_C \mathbb{H} = \{\varphi \in C_\top(X, Y) \mid \forall x \in X, \forall \mathbb{F} \in \mathcal{F}_L^\top(X), \\ x \in \lim^X \mathbb{F} \Rightarrow \varphi(x) \in \lim^Y ev_{X,Y}^\rightarrow(\mathbb{H} \times \mathbb{F})\}.$$

Then \lim_C is a \top -convergence on $C_\top(X, Y)$.

Proof. We have to check the map \lim_C satisfies the axioms (TC1) and (TC2). The axiom (TC2) follows immediately from the definition of \lim_C .

To check the map \lim_C satisfies the axioms (TC1), we have to show $\varphi \in \lim_C[\varphi]_\top$ holds for each $\varphi \in C_\top(X, Y)$, where $[\varphi]_\top = \{D \in L^{C_\top(X,Y)} \mid D(\varphi) = \top\}$ is the point \top -filter of φ on $C_\top(X, Y)$. And it suffices to check $x \in \lim^X \mathbb{F}$ implies $\varphi(x) \in \lim^Y ev_{X,Y}^\rightarrow([\varphi]_\top \times \mathbb{F})$ for all $x \in X$ and $\mathbb{F} \in \mathcal{F}_L^\top(X)$ by the definition of \lim_C . In fact, we firstly observe that for each $C \in \varphi^\rightarrow(\mathbb{F})$,

$$\begin{aligned} \top &= \bigvee_{A \in \mathbb{F}} \mathcal{S}_Y(\varphi^\rightarrow(A), C) \\ &= \bigvee_{A \in \mathbb{F}} \mathcal{S}_Y(ev_{X,Y}^\rightarrow(1_\varphi \times A), C) \quad (\text{by Lemma 3.6}) \\ &\leq \bigvee_{E \in [\varphi]_\top \times \mathbb{F}} \mathcal{S}_Y(ev_{X,Y}^\rightarrow(E), C) \quad (\text{here, } 1_\varphi \times A \in ([\varphi]_\top \times \mathbb{F}) \text{ for } A \in \mathbb{F}), \end{aligned}$$

which deduces $C \in ev_{\vec{X},Y}([\varphi]_{\top} \times \mathbb{F})$ by Remark 2.9 (1). Thus from the arbitrariness of C , $\varphi^{\rightarrow}(\mathbb{F}) \subseteq ev_{\vec{X},Y}([\varphi]_{\top} \times \mathbb{F})$ is obtained.

Finally, since $x \in \lim^X \mathbb{F}$ means $\varphi(x) \in \lim^Y \varphi^{\rightarrow}(\mathbb{F})$ by the continuity of φ for each $x \in X$, we have $\varphi(x) \in \lim^Y \varphi^{\rightarrow}(\mathbb{F}) \subseteq \lim^Y ev_{\vec{X},Y}([\varphi]_{\top} \times \mathbb{F})$ by using the property of \lim^Y satisfying the axiom (TC2). \square

Proposition 3.8. *Let (X, \lim^X) and (Y, \lim^Y) be two \top -Conv-objects. Then the evaluation map $ev_{X,Y} : (C_{\top}(X, Y) \times X, \lim_C \times \lim^X) \rightarrow (Y, \lim^Y)$ is continuous.*

Proof. Firstly, we point out the fact that $ev_{\vec{X},Y}(p_{C_{\top}(X,Y)}^{\rightarrow}(\mathbb{K}) \times p_X^{\rightarrow}(\mathbb{K})) \subseteq ev_{\vec{X},Y}(\mathbb{K})$ holds for each $\mathbb{K} \in \mathcal{F}_L^{\top}(C_{\top}(X, Y) \times X)$. To confirm the fact, it suffices to show $p_{C_{\top}(X,Y)}^{\rightarrow}(\mathbb{K}) \times p_X^{\rightarrow}(\mathbb{K}) \subseteq \mathbb{K}$. Since $p_{C_{\top}(X,Y)}^{\rightarrow}(A) \times p_X^{\rightarrow}(B) \geq A \wedge B$ holds for all $A, B \in \mathbb{K}$, we have

$$\begin{aligned} \bigvee_{C \in \mathbb{K}} \mathcal{S}_{C_{\top}(X,Y) \times X}(C, D) &\geq \bigvee_{A, B \in \mathbb{K}} \mathcal{S}_{C_{\top}(X,Y) \times X}(A \wedge B, D) \\ &\geq \bigvee_{A, B \in \mathbb{K}} \mathcal{S}_{C_{\top}(X,Y) \times X}(p_{C_{\top}(X,Y)}^{\rightarrow}(A) \times p_X^{\rightarrow}(B), D) \\ &= \top \end{aligned}$$

for every $D \in p_{C_{\top}(X,Y)}^{\rightarrow}(\mathbb{K}) \times p_X^{\rightarrow}(\mathbb{K})$. Thus whenever $D \in p_{C_{\top}(X,Y)}^{\rightarrow}(\mathbb{K}) \times p_X^{\rightarrow}(\mathbb{K})$, $D \in \mathbb{K}$ follows from (F1), which is to say $p_{C_{\top}(X,Y)}^{\rightarrow}(\mathbb{K}) \times p_X^{\rightarrow}(\mathbb{K}) \subseteq \mathbb{K}$ holds.

Now, we show the continuity of $ev_{X,Y}$ as follows:

Take any $(\varphi, x) \in C_{\top}(X, Y) \times X$ and any $\mathbb{K} \in \mathcal{F}_L^{\top}(C_{\top}(X, Y) \times X)$ such that $(\varphi, x) \in (\lim_C \times \lim^X)\mathbb{K}$. Then $\varphi \in \lim_C p_{C_{\top}(X,Y)}^{\rightarrow}(\mathbb{K})$ and $x \in \lim^X p_X^{\rightarrow}(\mathbb{K})$ hold. We have $\varphi(x) \in \lim^Y ev_{\vec{X},Y}(p_{C_{\top}(X,Y)}^{\rightarrow}(\mathbb{K}) \times p_X^{\rightarrow}(\mathbb{K}))$ from this and the definition of \lim_C . Finally, by using the fact above and the axiom (TC2), we conclude

$$ev_{X,Y}(\varphi, x) = \varphi(x) \in \lim^Y ev_{\vec{X},Y}(\mathbb{K}).$$

Consequently, $ev_{X,Y}$ is continuous. \square

Lemma 3.9. *Let (X, \lim^X) , (Y, \lim^Y) and (Z, \lim^Z) be \top -convergence spaces. If the map $\psi : (Z \times X, \lim^Z \times \lim^X) \rightarrow (Y, \lim^Y)$ is continuous, then for each $z \in Z$, the map $\psi(z, -) : (X, \lim^X) \rightarrow (Y, \lim^Y)$ is also continuous.*

Proof. To show the continuity of $\psi(z, -)$ for each $z \in Z$, take any $x \in X$ and $\mathbb{F} \in \mathcal{F}_L^{\top}(X)$ such that $x \in \lim^X \mathbb{F}$. $(z, x) \in (\lim^Z \times \lim^X)([z]_{\top} \times \mathbb{F})$ follows from $z \in \lim^Z [z]_{\top}$ and Lemma 3.5. Further, by the continuity of ψ , we observe that

$$\psi(z, -)(x) = \psi(z, x) \in \lim^Y \psi^{\rightarrow}([z]_{\top} \times \mathbb{F}).$$

Notice that for $A \in [z]_{\top}$ and $B \in \mathbb{F}$, $\psi^{\rightarrow}(A \times B) \geq \psi(z, -)^{\rightarrow}(B)$ holds because

$$\psi^{\rightarrow}(A \times B)(y) = \bigvee_{\psi(u,v)=y} A(u) \wedge B(v) \geq \bigvee_{\psi(z,v)=y} B(v) = \psi(z, -)^{\rightarrow}(B)(y)$$

for all $y \in Y$. From this, we observe that for $C \in \psi^{\Rightarrow}([z]_{\top} \times \mathbb{F})$,

$$\bigvee_{B \in \mathbb{F}} \mathcal{S}_Y(\psi(z, -)^{\rightarrow}(B), C) \geq \bigvee_{A \in [z]_{\top}, B \in \mathbb{F}} \mathcal{S}_Y(\psi^{\rightarrow}(A \times B), C) = \top,$$

which deduces $C \in \psi(z, -)^{\Rightarrow}(\mathbb{F})$. Thus, $\psi^{\Rightarrow}([z]_{\top} \times \mathbb{F}) \subseteq \psi(z, -)^{\Rightarrow}(\mathbb{F})$ follows from the arbitrariness of C . Finally, by the axiom (TC2) we obtain that for each $z \in Z$,

$$\psi(z, -)(x) \in \lim^Y \psi^{\Rightarrow}([z]_{\top} \times \mathbb{F}) \subseteq \lim^Y \psi(z, -)^{\Rightarrow}(\mathbb{F}).$$

In sum, the continuity of $\psi(z, -)$ is proved as desired. \square

Let (X, \lim^X) , (Y, \lim^Y) and (Z, \lim^Z) be \top -convergence spaces and

$$\psi : (Z \times X, \lim^Z \times \lim^X) \rightarrow (Y, \lim^Y)$$

be a continuous map. We define a map $\psi^* : Z \rightarrow C_{\top}(X, Y)$ by $\psi^*(z) = \psi(z, -)$ for all $z \in Z$. Then by Lemma 3.9, ψ^* is *well-defined* and the following lemma confirm that $\psi^* : (Z, \lim^Z) \rightarrow (C_{\top}(X, Y), \lim_C)$ is continuous.

Lemma 3.10. *Let (X, \lim^X) , (Y, \lim^Y) and (Z, \lim^Z) be \top -convergence spaces. If $\psi : (Z \times X, \lim^Z \times \lim^X) \rightarrow (Y, \lim^Y)$ is a continuous mapping, then the mapping $\psi^* : (Z, \lim^Z) \rightarrow (C_{\top}(X, Y), \lim_C)$ is continuous.*

Proof. Take any $z \in Z$ and $\mathbb{G} \in \mathcal{F}_L^{\top}(Z)$ such that $z \in \lim^Z \mathbb{G}$. We have to show

$$\psi^*(z) = \psi(z, -) \in \lim_C \psi^{*\Rightarrow}(\mathbb{G}).$$

For this, assume that $x \in \lim^X \mathbb{F}$, here $x \in X$ and $\mathbb{F} \in \mathcal{F}_L^{\top}(X)$. Then we have

$$(z, x) \in (\lim^Z \times \lim^X)(\mathbb{G} \times \mathbb{F}).$$

Immediately, $\psi(z, -)(x) = \psi(z, x) \in \lim^Y \psi^{\Rightarrow}(\mathbb{G} \times \mathbb{F})$ follows from the continuity of ψ . Further by means of $ev_{X,Y} \circ (\psi^* \times id_X) = \psi$ and Lemma 3.4, we observe

$$\lim^Y \psi^{\Rightarrow}(\mathbb{G} \times \mathbb{F}) = \lim^Y (ev_{X,Y} \circ (\psi^* \times id_X))^{\Rightarrow}(\mathbb{G} \times \mathbb{F}) = \lim^Y ev_{X,Y}^{\Rightarrow}(\psi^{*\Rightarrow}(\mathbb{G}) \times \mathbb{F}).$$

Hence $\psi^*(z) = \psi(z, -) \in \lim_C \psi^{*\Rightarrow}(\mathbb{G})$ is obtained from the definition of \lim_C . \square

Let (X, \lim^X) and (Y, \lim^Y) be two \top -**Conv**-objects. From Propositions 3.7, 3.8 and Lemma 3.10, there are the \top -**Conv**-object $(C_{\top}(X, Y), \lim_C)$ and the continuous map $ev_{X,Y} : (C_{\top}(X, Y) \times X, \lim_C \times \lim^X) \rightarrow (Y, \lim^Y)$ such that for each \top -**Conv**-object (Z, \lim^Z) and each continuous map

$$\psi : (Z \times X, \lim^Z \times \lim^X) \rightarrow (Y, \lim^Y),$$

there exists a unique continuous map $\psi^* : (Z, \lim^Z) \rightarrow (C_{\top}(X, Y), \lim_C)$ satisfying the equality $ev_{X,Y} \circ (\psi^* \times id_X) = \psi$. Thus by the definition of Cartesian-closedness, we obtain the following theorem.

Theorem 3.11. *The category \top -**Conv** of \top -convergence spaces is Cartesian-closed.*

4. Relation Between \top -convergences and Strong L -topologies

In this section, we show that there is a close relation between \top -convergences and strong L -topologies on a universal set X . In details, we will demonstrate that the category of strong L -topological spaces is concretely isomorphic to that of strong L -topological \top -convergence spaces categorically and embedded in the category $\top\text{-Conv}$ of \top -convergence spaces as a bireflective subcategory.

Firstly, we recall the definition of strong L -topological spaces [22] as follows.

Definition 4.1. Let X be a nonempty set. A *strong L -topological space* is a pair (X, τ) , where τ a subset of L^X such that the following conditions are satisfied:

- (ST1) $0_X, 1_X \in \tau$,
- (ST2) $U_1 \wedge U_2 \in \tau$ for all $U_1, U_2 \in \tau$,
- (ST3) $\bigvee_{j \in J} U_j \in \tau$ for every family $\{U_j \mid j \in J\} \subseteq \tau$,
- (ST4) $\alpha * U \in \tau$ for all $\alpha \in L$ and $U \in \tau$,
- (ST5) $\alpha \rightarrow U \in \tau$ for all $\alpha \in L$ and $U \in \tau$.

If (X, τ) is a strong L -topological space, then τ is called a *strong L -topology* on the set X . The set of all strong L -topologies on X is denoted by $ST_L(X)$. A map $\varphi : (X, \tau) \rightarrow (Y, \delta)$ between strong L -topological spaces is called *continuous map* provided that $\varphi^{\leftarrow}(V) \in \tau$ for each $V \in \delta$. We denote the category of strong L -topological spaces and continuous maps by $\mathbf{STOP}(L)$.

Definition 4.2. Let X be a nonempty set. If $\mathcal{U} = \{\mathbb{U}^x\}_{x \in X}$ is a family of \top -filters satisfying the axiom

$$(N) \quad \forall x \in X, \forall B \in \mathbb{U}^x, B(x) = \top,$$

we call \mathcal{U} a *system of \top -neighborhoods* on X . And if a system of \top -neighborhoods \mathcal{U} still satisfies the axiom

$$(TT) \quad \text{For any } x \in X \text{ and each } B \in \mathbb{U}^x, \text{ there exists } B^* \in \mathbb{U}^x \text{ with } B^* \leq B \text{ such that for every } y \in X, \text{ there exists } B_y \in \mathbb{U}^y \text{ satisfying } B^*(y) \leq \mathcal{S}_X(B_y, B),$$

\mathcal{U} is called a *strong L -topological system of \top -neighborhoods* on X .

Example 4.3. (1) Let (X, \lim) be a \top -convergence space. We denote $\mathcal{U}_{\lim} = \{\mathbb{U}_{\lim}^x\}_{x \in X}$, where $\mathbb{U}_{\lim}^x = \bigcap \{\mathbb{F} \in \mathcal{F}_L^\top(X) \mid x \in \lim \mathbb{F}\}$ for each $x \in X$. Then \mathcal{U}_{\lim} is a system of \top -neighborhoods.

(2) Let (X, τ) be a strong L -topological space. For each $x \in X$, \mathbb{U}_τ^x is defined by $\mathbb{U}_\tau^x = \{B \in L^X \mid \bigvee_{U \in \tau} U(x) * \mathcal{S}_X(U, B) = \top\}$. Then $\{\mathbb{U}_\tau^x\}_{x \in X}$ is a strong L -topological system of \top -neighborhoods on X . The detailed contents of the proof see [9].

Definition 4.4. A \top -convergence space (X, \lim) is said to be *strong L -topological* provided that $\{\mathbb{U}_{\lim}^x\}_{x \in X}$ satisfies the axioms (TT) and

$$(TP) \quad x \in \lim \mathbb{U}_{\lim}^x \text{ for all } x \in X.$$

The set of all strong L -topological \top -convergences on X is denoted by $STC^\top(X)$. The category of strong L -topological \top -convergence spaces and continuous maps is denoted by $\top\text{-STConv}$.

Example 4.5. Let (X, τ) be a strong L -topological space. The structure map $\lim_\tau : \mathcal{F}_L^\top(X) \rightarrow \mathcal{P}(X)$ induced by τ is defined by

$$\lim_\tau \mathbb{F} = \{x \in X \mid \mathbb{U}_\tau^x \subseteq \mathbb{F}\}, \text{ for all } \mathbb{F} \in \mathcal{F}_L^\top(X), \quad (1)$$

Then (X, \lim_τ) is a strong L -topological \top -convergence space.

Höhle [9] demonstrated that a strong L -topological space (X, τ) could induce a strong L -topological system of \top -neighborhoods $\{\mathbb{U}_\tau^x\}_{x \in X}$ (see Example 4.3 (2)). Besides, he also showed a strong L -topological system of \top -neighborhoods \mathcal{U} can induce a strong L -topology given by

$$\tau_{\mathcal{U}} = \{U \in L^X \mid U(x) \leq \bigvee_{B \in \mathcal{U}^x} \mathcal{S}_X(B, U), \forall x \in X\}.$$

Significantly, for the proofs of the above contents, it is dispensable that the condition Lemma 2.2 (M4) (i.e. \rightarrow preserving arbitrary union) which holds in the context of a complete MV -algebra.

Notice that if (X, \lim) is a \top -convergence space, $\{\mathbb{U}_{\lim}^x\}_{x \in X}$ is a system of \top -neighborhoods on X . Naturally, we can get a strong L -topology

$$\tau_{\lim} = \{U \in L^X \mid U(x) \leq \bigvee_{B \in \mathbb{U}_{\lim}^x} \mathcal{S}_X(B, U), \forall x \in X\}. \quad (2)$$

In fact, there exists a bijection between the set $STC^\top(X)$ of all strong L -topological \top -convergence structures on a set X and the set $ST_L(X)$ of all strong L -topologies on the X . We need a lemma in preparation for it.

Lemma 4.6. *Let (X, \lim) be a \top -convergence space. Then*

- (1) *for every $x \in X$, $\mathbb{U}_{\tau_{\lim}}^x \subseteq \mathbb{U}_{\lim}^x$ is valid.*
- (2) *if (X, \lim) is a strong L -topological, then for any $x \in X$, $\mathbb{U}_{\lim}^x = \mathbb{U}_{\tau_{\lim}}^x$ holds.*

Proof. (1) Take any $x \in X$ and let $B \in \mathbb{U}_{\tau_{\lim}}^x$, i.e. $\bigvee_{U \in \tau_{\lim}^x} U(x) * \mathcal{S}_X(U, B) = \top$ by Example 4.3 (2). From the formula (2) and Lemma 2.3 (2), we observe that $U \in \tau_{\lim}$ means

$$\begin{aligned} U(x) &\leq \bigvee_{C \in \mathbb{U}_{\lim}^x} \mathcal{S}_X(C, U) \\ &\leq \bigvee_{C \in \mathbb{U}_{\lim}^x} (\mathcal{S}_X(U, B) \rightarrow \mathcal{S}_X(C, B)) \\ &= \mathcal{S}_X(U, B) \rightarrow \bigvee_{C \in \mathbb{U}_{\lim}^x} \mathcal{S}_X(C, B), \end{aligned}$$

which is to say $U(x) * \mathcal{S}_X(U, B) \leq \bigvee_{C \in \mathbb{U}_{\lim}^x} \mathcal{S}_X(C, B)$. By the arbitrariness of U , we get $\top = \bigvee_{U \in \tau_{\lim}^x} U(x) * \mathcal{S}_X(U, B) \leq \bigvee_{C \in \mathbb{U}_{\lim}^x} \mathcal{S}_X(C, B)$. From this and (F1), $B \in \mathbb{U}_{\lim}^x$ holds, i.e. $\mathbb{U}_{\tau_{\lim}}^x \subseteq \mathbb{U}_{\lim}^x$ is obtained and the proof of the conclusion is completed.

(2) Let (X, \lim) be a strong L -topological space. For any $x \in X$, we only need to show $\mathbb{U}_{\lim}^x \subseteq \mathbb{U}_{\tau_{\lim}}^x$ by using (1). Now take any $B \in \mathbb{U}_{\lim}^x$ and define $\bar{B} \in L^X$

by $\overline{B}(y) = \bigvee_{A \in \mathbb{U}_{\text{lim}}^y} \mathcal{S}_X(A, B)$ for each $y \in X$. Then $\overline{B}(x) = \top$ and $\overline{B} \leq B$ follows respectively from $\overline{B}(x) = \bigvee_{A \in \mathbb{U}_{\text{lim}}^x} \mathcal{S}_X(A, B) \geq \mathcal{S}_X(B, B) = \top$ and for each $y \in X$

$$\overline{B}(y) = \bigvee_{A \in \mathbb{U}_{\text{lim}}^y} \mathcal{S}_X(A, B) \leq \bigvee_{A \in \mathbb{U}_{\text{lim}}^y} (A(y) \rightarrow B(y)) = \bigvee_{A \in \mathbb{U}_{\text{lim}}^y} (\top \rightarrow B(y)) = B(y).$$

In the following, we additionally confirm $\overline{B} \in \tau_{\text{lim}}$ is also true by means of the axiom (TT). In fact, for each $y \in X$, the axiom (TT) tell us $A \in \mathbb{U}_{\text{lim}}^y$ implies there is $A^* \in \mathbb{U}_{\text{lim}}^y$ with $A^* \leq A$ such that for all $z \in X$, there exists $A_z \in \mathbb{U}_{\text{lim}}^z$ to satisfy $A^*(z) \leq \mathcal{S}_X(A_z, A)$. Then we get $\mathcal{S}_X(A, B) \leq \mathcal{S}_X(A^*, \overline{B})$ because for all $z \in X$

$$\begin{aligned} \mathcal{S}_X(A, B) &\leq \mathcal{S}_X(A_z, A) \rightarrow \mathcal{S}_X(A_z, B) \\ &\leq A^*(z) \rightarrow \mathcal{S}_X(A_z, B) \\ &\leq A^*(z) \rightarrow \left(\bigvee_{C \in \mathbb{U}_{\text{lim}}^z} \mathcal{S}_X(C, B) \right) \\ &= A^*(z) \rightarrow \overline{B}(z). \end{aligned}$$

From this, we conclude that for all $y \in X$,

$$\overline{B}(y) = \bigvee_{A \in \mathbb{U}_{\text{lim}}^y} \mathcal{S}_X(A, B) \leq \bigvee_{A^* \in \mathbb{U}_{\text{lim}}^y} \mathcal{S}_X(A^*, \overline{B}) \leq \bigvee_{C \in \mathbb{U}_{\text{lim}}^y} \mathcal{S}_X(C, \overline{B}).$$

Namely, $\overline{B} \in \tau_{\text{lim}}$ by the definition of τ_{lim} . From all above, we have

$$\bigvee_{U \in \tau_{\text{lim}}} U(x) * \mathcal{S}_X(U, B) \geq \overline{B}(x) * \mathcal{S}_X(\overline{B}, B) = \top,$$

i.e. $B \in \mathbb{U}_{\tau_{\text{lim}}}^x$ holds. Finally, $\mathbb{U}_{\text{lim}}^x \subseteq \mathbb{U}_{\tau_{\text{lim}}}^x$ follows from the arbitrariness of B . \square

Theorem 4.7. *Let X be a nonempty set.*

- (1) *Given a strong L -topology τ on X , then $\tau_{\text{lim}, \tau} = \tau$.*
- (2) *Given a strong L -topological \top -convergence lim on X , then $\text{lim}_{\tau_{\text{lim}}} = \text{lim}$.*

Proof. (1) By the formula (2), we know

$$\tau_{\text{lim}, \tau} = \{U \in L^X \mid U(x) \leq \bigvee_{A \in \mathbb{U}_{\text{lim}, \tau}^x} \mathcal{S}_X(A, U), \forall x \in X\}.$$

From Example 4.3 and the formula (1), we observe that for all $x \in X$,

$$\mathbb{U}_{\text{lim}, \tau}^x = \bigcap \{\mathbb{F} \in \mathcal{F}_L^\top(X) \mid x \in \text{lim}_\tau \mathbb{F}\} = \bigcap \{\mathbb{F} \in \mathcal{F}_L^\top(X) \mid \mathbb{U}_\tau^x \subseteq \mathbb{F}\} = \mathbb{U}_\tau^x.$$

Above all, $\tau_{\text{lim}, \tau} = \{U \in L^X \mid U(x) \leq \bigvee_{A \in \mathbb{U}_\tau^x} \mathcal{S}_X(A, U), \forall x \in X\}$. In fact, $U \in \tau$ if and only if $U(x) \leq \bigvee_{B \in \mathbb{U}_\tau^x} \mathcal{S}_X(B, U)$ holds for every $x \in X$ (see [9]). Then $\tau_{\text{lim}, \tau} = \tau$ is true.

(2) In order to show $\text{lim}_{\tau_{\text{lim}}} = \text{lim}$, take any $\mathbb{F} \in \mathcal{F}_L^\top(X)$.

$$\text{lim}_{\tau_{\text{lim}}} \mathbb{F} = \{x \in X \mid \mathbb{U}_{\tau_{\text{lim}}}^x \subseteq \mathbb{F}\} = \{x \in X \mid \mathbb{U}_{\text{lim}}^x \subseteq \mathbb{F}\}$$

follows from Lemma 4.6 (2). Further, for each $x \in X$ and $\mathbb{F} \in \mathcal{F}_L^\top(X)$, we have

$$x \in \lim_{\tau_{\text{lim}}} \mathbb{F} \iff \mathbb{U}_{\text{lim}}^x \subseteq \mathbb{F} \iff x \in \lim \mathbb{F}.$$

Consequently, $\lim_{\tau_{\text{lim}}} = \lim$ is true. \square

Corollary 4.8. *Let X be a nonempty set. Then there is a bijection between the set of all strong L -topological \top -convergences and the set of all strong L -topologies on X .*

In order to explore the relation between the category of strong L -topological spaces and that of \top -convergence spaces, the following theorem is indispensable.

Theorem 4.9. *Let (X, τ) and (Y, δ) be strong L -topological spaces, (X, \lim^X) and (Y, \lim^Y) be \top -convergence spaces. Then the following are valid:*

(1) *If $\varphi : (X, \tau) \rightarrow (Y, \delta)$ is a continuous map, then $\varphi : (X, \lim_\tau) \rightarrow (Y, \lim_\delta)$ is a continuous map.*

(2) *If $\varphi : (X, \lim^X) \rightarrow (Y, \lim^Y)$ is a continuous map, then $\varphi : (X, \tau_{\text{lim}^X}) \rightarrow (Y, \delta_{\text{lim}^Y})$ is a continuous map.*

Proof. (1) First, we point out the fact given in [9] that a map $\varphi : (X, \tau) \rightarrow (Y, \delta)$ is continuous if and only if φ is continuous at each $x \in X$ in the sense that $\mathbb{U}_\delta^{\varphi(x)} \subseteq \varphi^\Rightarrow(\mathbb{U}_\tau^x)$ holds. Let $x \in X$ and $\mathbb{F} \in \mathcal{F}_L^\top(X)$ such that $x \in \lim_\tau \mathbb{F}$, equivalently $\mathbb{U}_\tau^x \subseteq \mathbb{F}$. Then $\varphi^\Rightarrow(\mathbb{U}_\tau^x) \subseteq \varphi^\Rightarrow(\mathbb{F})$ follows from (i) in Remark 2.9 (3). Because the mapping $\varphi : (X, \tau) \rightarrow (Y, \delta)$ is continuous, which is to say $\mathbb{U}_\delta^{\varphi(x)} \subseteq \varphi^\Rightarrow(\mathbb{U}_\tau^x)$ holds from the fact above. Hence $\mathbb{U}_\delta^{\varphi(x)} \subseteq \varphi^\Rightarrow(\mathbb{F})$ is true, which already means that $\varphi(x) \in \lim_\delta \varphi^\Rightarrow(\mathbb{F})$. Consequently, the continuity of φ from (X, \lim_τ) to (Y, \lim_δ) is proved as desired.

(2) Firstly, we are going to show that $\varphi^\Leftarrow(\mathbb{U}_{\text{lim}^Y}^{\varphi(x)}) \subseteq \mathbb{U}_{\text{lim}^X}^x$ holds, here

$$\mathbb{U}_{\text{lim}^X}^x = \bigcap \{ \mathbb{F} \in \mathcal{F}_L^\top(X) \mid x \in \lim^X \mathbb{F} \}$$

defined in Example 4.3 (1). For all $C \in \mathbb{U}_{\text{lim}^Y}^{\varphi(x)}$, $C(\varphi(x)) = \top$ holds by Example 4.3 (1). So $\varphi^\Leftarrow(\mathbb{U}_{\text{lim}^Y}^{\varphi(x)})$ exists from Definition 2.8 (2). For any $\mathbb{F} \in \mathcal{F}_L^\top(X)$ with $x \in \lim^X \mathbb{F}$, by the continuity of φ from (X, \lim^X) to (Y, \lim^Y) , we have $\varphi(x) \in \lim^Y \varphi^\Rightarrow(\mathbb{F})$. Then $\mathbb{U}_{\text{lim}^Y}^{\varphi(x)} \subseteq \varphi^\Rightarrow(\mathbb{F})$ holds. From this and (ii) in Remark 2.9 (3),

$$\varphi^\Leftarrow(\mathbb{U}_{\text{lim}^Y}^{\varphi(x)}) \subseteq \varphi^\Leftarrow \circ \varphi^\Rightarrow(\mathbb{F}) \subseteq \mathbb{F}$$

is valid. Hence we get

$$\varphi^\Leftarrow(\mathbb{U}_{\text{lim}^Y}^{\varphi(x)}) \subseteq \bigcap \{ \mathbb{F} \in \mathcal{F}_L^\top(X) \mid x \in \lim^X \mathbb{F} \} = \mathbb{U}_{\text{lim}^X}^x.$$

Now, we prove $\varphi : (X, \tau_{\text{lim}^X}) \rightarrow (Y, \delta_{\text{lim}^Y})$ is continuous. Take any $V \in \delta_{\text{lim}^Y}$, and hence $V(\varphi(x)) \leq \bigvee_{B \in \mathbb{U}_{\text{lim}^Y}^{\varphi(x)}} \mathcal{S}_Y(B, V)$ for $x \in X$. In this case, we observe that

$$\begin{aligned}
\bigvee_{A \in \mathbb{U}_{\lim^x}^x} \mathcal{S}_X(A, \varphi^\leftarrow(V)) &\geq \bigvee_{A \in \varphi^\leftarrow(\mathbb{U}_{\lim^Y}^{\varphi(x)})} \mathcal{S}_X(A, \varphi^\leftarrow(V)) \\
&\geq \bigvee_{B \in \mathbb{U}_{\lim^Y}^{\varphi(x)}} \mathcal{S}_X(\varphi^\leftarrow(B), \varphi^\leftarrow(V)) \\
&\geq \bigvee_{B \in \mathbb{U}_{\lim^Y}^{\varphi(x)}} \mathcal{S}_Y(B, V) \quad (\text{by Lemma 2.3(4)}) \\
&\geq V(\varphi(x)) = \varphi^\leftarrow(V)(x),
\end{aligned}$$

which is to say $\varphi^\leftarrow(V)(x) \leq \bigvee_{A \in \mathbb{U}_{\lim^x}^x} \mathcal{S}_X(A, \varphi^\leftarrow(V))$ holds for all $x \in X$. By definition of τ_{\lim^x} , $\varphi^\leftarrow(V) \in \tau_{\lim^x}$ holds. Consequently, $\varphi : (X, \tau_{\lim^x}) \rightarrow (Y, \delta_{\lim^Y})$ is continuous. \square

From Corollary 4.8 and Theorem 4.9, the following theorem is obtained.

Theorem 4.10. *The category $\mathbf{STOP}(L)$ of strong L -topological spaces is concretely isomorphic to the category $\mathbb{T}\text{-}\mathbf{STConv}$ of strong L -topological \mathbb{T} -convergence spaces, categorically.*

The following theorem shows the category $\mathbb{T}\text{-}\mathbf{STConv}$ is bireflective in the category $\mathbb{T}\text{-}\mathbf{Conv}$ in some detail.

Theorem 4.11. *The category $\mathbb{T}\text{-}\mathbf{STConv}$ of strong L -topological \mathbb{T} -convergence spaces is bireflective in the category $\mathbb{T}\text{-}\mathbf{Conv}$ of \mathbb{T} -convergence spaces.*

Proof. Let (X, \lim) be a $\mathbb{T}\text{-}\mathbf{Conv}$ -object. We claim that $\text{id}_X : (X, \lim) \rightarrow (X, \widetilde{\lim})$ is the bireflection of (X, \lim) w.r.t. $\mathbb{T}\text{-}\mathbf{STConv}$, where $\widetilde{\lim} = \lim_{\tau_{\lim}}$, i.e. for each $\mathbb{F} \in \mathcal{F}_L^\mathbb{T}(X)$, $\widetilde{\lim}\mathbb{F} = \{x \in X \mid \mathbb{U}_{\tau_{\lim}}^x \subseteq \mathbb{F}\}$. By Example 4.5, we know \lim is a strong L -topological \mathbb{T} -convergence on X . By Definition 2.12, we need to confirm

- (1) $\text{id}_X : (X, \lim) \rightarrow (X, \widetilde{\lim})$ is continuous.
- (2) For any continuous map $\varphi : (X, \lim) \rightarrow (Y, \lim^Y)$, where (Y, \lim^Y) is a $\mathbb{T}\text{-}\mathbf{STConv}$ -object, $\varphi : (X, \widetilde{\lim}) \rightarrow (Y, \lim^Y)$ is continuous.

In fact, for the conclusion (1), take any $\mathbb{F} \in \mathcal{F}_L^\mathbb{T}(X)$ and $x \in X$ such that $x \in \lim\mathbb{F}$. Then $\mathbb{U}_{\lim}^x \subseteq \mathbb{F}$ holds. And by Lemma 4.6 (1), $\mathbb{U}_{\tau_{\lim}}^x \subseteq \mathbb{U}_{\lim}^x$. Hence we further obtain $\mathbb{U}_{\tau_{\lim}}^x \subseteq \mathbb{F}$, i.e. $\text{id}_X(x) = x \in \widetilde{\lim}\mathbb{F} = \lim(\text{id}_X)^\Rightarrow(\mathbb{F})$, which can deduces that id_X is continuous.

For the conclusion (2), assume a map $\varphi : (X, \lim) \rightarrow (Y, \lim^Y)$ is continuous, where (Y, \lim^Y) is a $\mathbb{T}\text{-}\mathbf{STConv}$ -object and hence satisfies $\lim^Y = \lim_{\delta_{\lim^Y}}$ by Theorem 4.7. By Theorem 4.9 (1) and (2), $\varphi : (X, \tau_{\lim}) \rightarrow (Y, \delta_{\lim^Y})$ is continuous first, and then the map $\varphi : (X, \lim_{\tau_{\lim}}) \rightarrow (Y, \lim_{\delta_{\lim^Y}})$ is continuous also, which is to say that $\varphi : (X, \widetilde{\lim}) \rightarrow (Y, \lim^Y)$ is continuous. \square

By Theorems 4.10 and 4.11, we can get the following corollary.

Corollary 4.12. *The category $\mathbf{STOP}(L)$ of strong L -topological spaces is embedded in the category $\mathbb{T}\text{-}\mathbf{Conv}$ of \mathbb{T} -convergence spaces as a bireflective subcategory.*

5. Conclusions

In this paper, we proposed the concept of \top -convergence spaces based on \top -filters. Afterwards, in the lattice valued context of a complete MV -algebra without the idempotency of the semigroup operation, we proved the category of \top -convergence spaces is topological and further is Cartesian-closed. Additionally, the category of strong L -topological spaces can be bireflectively embedded in the category of \top -convergence spaces.

Interestingly, we could study the corresponding subcategories of \top -**Conv** by generalizing the well-known categories of Kent convergence spaces, of limit spaces, of pseudo-topological spaces to the many-valued setting as well as their categorical relations. From the paper [3], we have known the category **STOP**(L) is a reflective subcategory of the category L -**OCS** of L -ordered convergence spaces. Then, is there any link between the category \top -**Conv** and the category L -**OCS**? We will study these problems in the future.

Acknowledgements. Authors would like to express their sincere thanks to the anonymous reviewers for their valuable comments and suggestions. This work was supported by NSFC (11201437,11471297,11401547).

REFERENCES

- [1] G. Choquet, *Convergences*, Ann. Univ. Grenoble, **23** (1948), 57–112.
- [2] J. M. Fang, *Stratified L -ordered convergence structures*, Fuzzy Sets and Systems, **161** (2010), 2130–2149.
- [3] J. M. Fang, *Relationships between L -ordered convergence structures and strong L -topologies*, Fuzzy Sets and Systems, **161** (2010), 2923–2944.
- [4] J. A. Goguen, *L -fuzzy sets*, J. Math. Anal. Appl, **18** (1967), 145–174.
- [5] J. Gutiérrez García and M. A. De Prada Vicente, *Characteristic values of \top -filter*, Fuzzy Sets and Systems, **156** (2005), 55–67.
- [6] J. Gutiérrez García and M. A. De Prada Vicente, *A unified approach to the concept of fuzzy L -uniform space*, In: Topological and Algebraic Structures in Fuzzy Sets—A Handbook of Recent Developments in the Mathematics of Fuzzy Sets, (S.E. Rodabaugh, E.P. Klement, ed.), Kluwer Academic Publishers, Boston, Dordrecht, London, (2003), 79–114.
- [7] U. Höhle, *Commutative residuated ℓ -monoids*, In: Non-classical Logics and Their Applications to Fuzzy Subsets: A Handbook of the Mathematical Foundations of Fuzzy Set Theory, (U. Höhle, E.P. Klement, Eds.), Klumer Academic Publishers, Dordrecht, Boston, London (1995), 53–106.
- [8] U. Höhle, *Probabilistische topologien*, Manuscripta Math., **26** (1978), 223–245.
- [9] U. Höhle, *Probabilistic topologies induced by L -fuzzy uniformities*, Manuscripta Math., **38** (1982), 289–323.
- [10] G. Jäger, *A category of L -fuzzy convergence spaces*, Quaestiones Mathematicae, **24** (2001), 501–517.
- [11] G. Jäger, *Subcategories of lattice-valued convergence spaces*, Fuzzy Sets and Systems, **156** (2005), 1–24.
- [12] G. Jäger, *Lattice-valued convergence spaces and regularity*, Fuzzy Sets and Systems, **159** (2008), 2488–2502.
- [13] R. Lowen, *Convergence in fuzzy topological spaces*, Gen. Top. Appl., **10** (1979), 147–160.
- [14] E. Lowen and R. Lowen, *A topological universe extension of FTS* , In: Applications of category theory to fuzzy sets, (S.E. Rodabaugh, E.P. Klement, U. Höhle, Eds.), Kluwer, Dordrecht (1992), 153–176.

- [15] E. Lowen, R. Lowen and P. Wuyts, *The categorical topology approach to fuzzy topology and fuzzy convergence*, Fuzzy Sets and Systems, **40** (1991), 347–343.
- [16] L. Li, Q. Jin and K. Hu, *On stratified L -convergence spaces: Fischer's diagonal axiom*, Fuzzy Sets and Systems, **267** (2015), 31–40.
- [17] G. Preuss, *Foundations of topology*, Kluwer Academic Publishers, Dordrecht, Boston, London (2002), 30–92.
- [18] B. Pang and J. M. Fang, *L -fuzzy Q -convergence structures*, Fuzzy Sets and System, **182** (2011), 53–65.
- [19] B. Pang, *On (L, M) -fuzzy convergence spaces*, Fuzzy Sets and Systems, **238** (2014), 46–70.
- [20] W. Yao, *On many-valued stratified L -fuzzy convergence spaces*, Fuzzy Sets and Systems, **159** (2008), 2503–2519.
- [21] W. Yao, *Moore-Smith convergence in (L, M) -fuzzy topology*, Fuzzy Sets and Systems, **190** (2012), 47–62.
- [22] D. Zhang, *An enriched category approach to many valued topology*, Fuzzy Sets and Systems, **158** (2007), 349–366.

QIAN YU, DEPARTMENT OF MATHEMATICS, OCEAN UNIVERSITY OF CHINA, 238 SONGLING ROAD, 266100, QINGDAO, P.R. CHINA
E-mail address: 459778696@qq.com

JINMING FANG*, DEPARTMENT OF MATHEMATICS, OCEAN UNIVERSITY OF CHINA, 238 SONGLING ROAD, 266100, QINGDAO, P.R. CHINA
E-mail address: jinming-fang@163.com

*CORRESPONDING AUTHOR

Archive of SID

THE CATEGORY OF T-CONVERGENCE SPACES AND ITS CARTESIAN-CLOSEDNESS

Q. YU AND J. FANG

رسته فضاهای T – همگرا و بسته بودن دکارتی آنها

چکیده. در این مقاله، نوعی از فضاهای همگرای شبکه – مقدار بر اساس مفهوم T-فیلترها، یعنی فضاهای T-همگرا را تعریف می کنیم و نشان می دهیم که رسته فضاهای T-همگرا بسته دکارتی است. علاوه، در زمینه شبکه مقداری یک MV – جبر کامل، رابطه ی نزدیکی بین رسته فضاهای T-همگرا و رسته فضاهای L – توپولوژیکی قوی برقرار شده است. به تفصیل، نشان می دهیم که رسته فضاهای L – توپولوژیکی قوی با رسته فضاهای T-همگرای L – توپولوژیکی قوی بطور محسوسی یکریخت رسته ای است و بطور دو جانبه در فضاهای T-همگرا نشانده می شود.

Archive of SID