

Solving LR fuzzy linear matrix equation[†]X. Guo¹ and D. Shang²¹ College of Mathematics and Statistics, Northwest Normal University, Lanzhou 730070, China² Public Health College, Gansu University of Traditional Chinese Medicine, Lanzhou 730000, China

guoxb@nwnu.edu.cn, sdqglz@qq.com

Abstract

In this paper, the fuzzy matrix equation $A\tilde{X}B = \tilde{C}$ in which A, B are $n \times n$ crisp matrices respectively and \tilde{C} is an $n \times n$ arbitrary LR fuzzy numbers matrix, is investigated. A new numerical procedure for calculating the fuzzy solution is designed and a sufficient condition for the existence of strong fuzzy solution is derived. Some examples are given to illustrate the proposed method.

Keywords: Fuzzy numbers, matrix analysis, fuzzy matrix equation, fuzzy approximate solutions.

1 Introduction

Imprecision is often involved in many real-world engineering systems. Fuzzy systems have been an essential role in the fuzzy modeling, which can formulate uncertainty in actual environment. In many matrix systems, some of the system parameters are vague or imprecise, and fuzzy number is a better tool than crisp one for modeling the problem, and hence solving a fuzzy matrix system is becoming more important. The concept of fuzzy numbers and their arithmetic operations were first introduced and investigated by Zadeh [40], Dubois et al. [19] and Nahmias [31]. A different approach to fuzzy numbers and the structure of fuzzy number spaces was given by Puri and Ralescu [37], Goetschell et al. [23] and Wu Congxin et al. [38, 39].

Since Friedman et al. [21] proposed a general model for solving an $n \times n$ fuzzy linear systems $A\tilde{x} = \tilde{b}$ by an embedding approach in 1998, lots of research works have been done about some more complicated fuzzy linear systems such as dual fuzzy linear systems (DFLS), general fuzzy linear systems (GFLS), fully fuzzy linear systems (FFLS) and general dual fuzzy linear systems (GDFLS) see [1–6, 11, 16–17, 21, 30, 41]. And some new theory and method for fuzzy linear systems still appeared recently [7, 12, 20, 22, 32–34].

However, for a fuzzy linear matrix equation which always has a wide use in control theory and control engineering, few work has been done in the past decades. In 2009, Allahviranloo et al. [9] discussed the fuzzy linear matrix equations (FLME) with the form of $A\tilde{X}B = \tilde{C}$ by embedding approach. In 2011, Gong and Guo [24] investigated a class of inconsistent fuzzy matrix equations $A\tilde{X} = \tilde{B}$ and obtained its fuzzy least squares solutions by the same way. In 2017, Guo et al. [27] presented a computing method for fuzzy matrix equations $\tilde{X}A = \tilde{B}$. Recently, Amirfakhrian et al. [10] proposed a new method for solving fuzzy matrix equations $A\tilde{X}B = \tilde{C}$ based on triangular fuzzy numbers.

In traditional fuzzy linear systems, the uncertain elements were usually denoted by the parametric form of fuzzy numbers. Thus the extended linear equations always contains parameter $r, 0 \leq r \leq 1$, which makes their computation inconvenient in some sense. On the other hand, the weak fuzzy solution of fuzzy linear systems does not exist sometimes [8]. To make the multiplication of fuzzy numbers easy and handle the full fuzzy linear systems, Dubois and Prade [19] introduced the LR fuzzy number in 1978. We know that triangular fuzzy numbers is just specious cases of LR fuzzy numbers. In 2012, Otadi and Mosleh [35] set up a model for solving fully fuzzy matrix equations $\tilde{A}\tilde{X} = \tilde{B}$. In 2013, Guo et al. [26] proposed a computing method for solving fuzzy Sylvester matrix equation $A\tilde{X} + \tilde{X}B = \tilde{C}$ with LR fuzzy numbers. For fully fuzzy Sylvester matrix equation $\tilde{A}\tilde{X} + \tilde{X}\tilde{B} = \tilde{C}$, Daud et al. [15] and Dookhitram et al. [18]

Corresponding Author: X. Guo

Received: December 2017; Revised: April 2018; Accepted: April 2019.

carried on a thorough investigations in recent years. In 2014, Gong et al. [25] studied the general dual fuzzy matrix systems $A\tilde{X} + \tilde{B} = C\tilde{X} + \tilde{D}$ according to arithmetic operations of LR fuzzy numbers. In 2017, Kaur and Kumar [29] made a note on Gong et al.'s paper. Recently, Guo et al.[28] made a further investigation for dual fuzzy matrix systems $A\tilde{X} + \tilde{B} = C\tilde{X} + \tilde{D}$.

In this paper, we propose a general model for solving the fuzzy matrix equation $A\tilde{X}B = \tilde{C}$ in which A, B are $n \times n$ crisp matrices and \tilde{C} is an $n \times n$ arbitrary LR fuzzy numbers matrix respectively. We extend the fuzzy matrix equation into a crisp system of linear matrix equations. We obtain the fuzzy minimal solution of the original fuzzy matrix equation from solving the crisp function matrix system. Moreover, we discussed the existence condition of the strong fuzzy minimal solution. Finally, we given some examples to illustrate the proposed method. The structure of this paper is organized as follows:

In Section 2, we recall the fuzzy number and present the concept of the LR fuzzy linear matrix equation. In Section 3, the model to the fuzzy matrix equation is proposed in detail and the minimal fuzzy solution of the fuzzy linear matrix equation is obtained. Some examples are given to illustrate our method in Section 4 and the conclusion is drawn in Section 5.

2 Preliminaries

2.1 The LR fuzzy number

There are several definitions for the concept of LR fuzzy numbers (see [19, 31, 40]).

Definition 2.1. A fuzzy number is a fuzzy set $u : R \rightarrow I = [0, 1]$ that satisfies:

- (1) u is upper semicontinuous,
- (2) u is fuzzy convex, i.e., $u(\lambda x + (1 - \lambda)y) \geq \min\{u(x), u(y)\}$ for all $x, y \in R, \lambda \in [0, 1]$,
- (3) u is normal, i.e., there exists $x_0 \in R$ such that $u(x_0) = 1$,
- (4) $suppu = \{x \in R \mid u(x) > 0\}$ is the support of the u , and its closure $cl(suppu)$ is compact.

Let E^1 be the set of all fuzzy numbers on R .

Definition 2.2. A fuzzy number \tilde{M} is said to be a LR fuzzy number if

$$\mu_{\tilde{M}}(x) = \begin{cases} L(\frac{m-x}{\alpha}), & x \leq m, \quad \alpha > 0, \\ R(\frac{x-m}{\beta}), & x \geq m, \quad \beta > 0, \end{cases}$$

where m, α and β are called the mean value, left and right spreads of \tilde{M} , respectively. The function $L(\cdot)$, which is called left shape function satisfies:

- (1) $L(x) = L(-x)$,
- (2) $L(0) = 1$ and $L(1) = 0$,
- (3) $L(x)$ is non increasing on $[0, \infty)$.

The definition of a right shape function $R(\cdot)$ is similar to that of $L(\cdot)$.

Clearly, two LR fuzzy numbers $\tilde{M} = (m, \alpha, \beta)_{LR}$ and $\tilde{N} = (n, \gamma, \delta)_{LR}$ are said to be equal, if and only if $m = n, \alpha = \gamma$ and $\beta = \delta$. Also, $\tilde{M} = (m, \alpha, \beta)_{LR}$ is positive(negative) if and only if $m - \alpha > 0(m + \beta < 0)$.

Definition 2.3. For arbitrary LR fuzzy numbers $\tilde{M} = (m, \alpha, \beta)_{LR}$ and $\tilde{N} = (n, \gamma, \delta)_{LR}$, we have

- (1) Addition

$$\tilde{M} + \tilde{N} = (m, \alpha, \beta)_{LR} + (n, \gamma, \delta)_{LR} = (m + n, \alpha + \gamma, \beta + \delta)_{LR}. \tag{1}$$

- (2) Subtraction

$$\tilde{M} - \tilde{N} = (m, \alpha, \beta)_{LR} - (n, \gamma, \delta)_{LR} = (m - n, \alpha - \delta, \beta - \gamma)_{LR}. \tag{2}$$

- (3) Scalar multiplication

$$\lambda\tilde{M} = \lambda(m, \alpha, \beta)_{LR} \cong \begin{cases} (\lambda m, \lambda\alpha, \lambda\beta), & \lambda \geq 0, \\ (\lambda m, -\lambda\beta, -\lambda\alpha)_{RL}, & \lambda < 0. \end{cases} \tag{3}$$

Proof. Let $\tilde{C} = (C, C^l, C^r) = (c_{ij}, c_{ij}^l, c_{ij}^r)_{n \times n}$, $\tilde{X} = (X, X^l, X^r) = (x_{ij}, x_{ij}^l, x_{ij}^r)_{n \times n}$. We also let $A = A^+ + A^-$ where the elements a_{ij}^+ of matrix A^+ and a_{ij}^- of matrix A^- are determined by this way: if $a_{ij} \geq 0, a_{ij}^+ = a_{ij}$ else $a_{ij}^+ = 0, 1 \leq i, j \leq n$; if $a_{ij} < 0, a_{ij}^- = a_{ij}$ else $a_{ij}^- = 0, 1 \leq i, j \leq n$. For fuzzy matrix equation $A\tilde{X}B = \tilde{C}$, we can express it as

$$(A^+ + A^-)(X, X^l, X^r)B = (C, C^l, C^r). \tag{11}$$

Since

$$k\tilde{x}_{ij} = \begin{cases} (kx_{ij}, kx_{ij}^l, kx_{ij}^r), & k \geq 0, \\ (kx_{ij}, -kx_{ij}^r, -kx_{ij}^l), & k < 0, \end{cases}$$

we have

$$A\tilde{X} = \begin{cases} (AX, AX^l, AX^r), & A \geq 0, \\ (AX, -AX^r, -AX^l), & A < 0. \end{cases}$$

So the equation (11) can be rewritten as

$$A^+(X, X^l, X^r)B + A^-(X, X^l, X^r)B = (A^+X, A^+X^l, A^+X^r)B + (A^-X, -A^-X^r, -A^-X^l)B.$$

Since

$$k\tilde{x}_{ij} = \begin{cases} (kx_{ij}, kx_{ij}^l, kx_{ij}^r), & k \geq 0, \\ (kx_{ij}, -kx_{ij}^r, -kx_{ij}^l), & k < 0, \end{cases}$$

we have

$$\tilde{X}B = \begin{cases} (XB, X^lB, X^rB), & B \geq 0, \\ (XB, -X^rB, -X^lB), & B < 0. \end{cases}$$

Supposing $B = B^+ + B^-$ in which the elements b_{ij}^+ of matrix B^+ and b_{ij}^- of matrix B^- are determined by this way: if $b_{ij} \geq 0, b_{ij}^+ = b_{ij}$ else $b_{ij}^+ = 0, 1 \leq i, j \leq n$; if $b_{ij} < 0, b_{ij}^- = b_{ij}$ else $b_{ij}^- = 0, 1 \leq i, j \leq n$, we have

$$\begin{aligned} & (A^+XB^+, A^+X^lB^+, A^+X^rB^+) + (A^-XB^+, -A^-X^rB^+, -A^-X^lB^+) \\ & + (A^+XB^-, -A^+X^rB^-, -A^+X^lB^-) + (A^-XB^-, A^-X^lB^-, A^-X^rB^-) = \\ & (A^+XB^+ + A^-XB^+ + A^+XB^- + A^-XB^-, A^+X^lB^+ - A^-X^rB^+ - A^+X^rB^- \\ & + A^-X^lB^-, A^+X^rB^+ - A^-X^lB^+ - A^+X^lB^- + A^-X^rB^-) = (C, C^l, C^r). \end{aligned}$$

Thus we get

$$\begin{cases} A^+XB^+ + A^-XB^+ + A^+XB^- + A^-XB^- = C, \\ A^+X^lB^+ - A^-X^rB^+ - A^+X^rB^- + A^-X^lB^- = C^l, \\ A^+X^rB^+ - A^-X^lB^+ - A^+X^lB^- + A^-X^rB^- = C^r. \end{cases} \tag{12}$$

Denoting (12) in matrix form, we have

$$\begin{cases} (A^+ + A^-)X(B^+ + B^-) = C, \\ (A^+X^l, A^+X^r) \begin{pmatrix} B^+ & -B^- \\ -B^- & B^+ \end{pmatrix} + (-A^-X^r, -A^-X^l) \begin{pmatrix} B^+ & -B^- \\ -B^- & B^+ \end{pmatrix} = (C^l, C^r), \end{cases}$$

or

$$\begin{cases} (A^+ + A^-)X(B^+ + B^-) = C, \\ A^+(X^l, X^r) \begin{pmatrix} B^+ & -B^- \\ -B^- & B^+ \end{pmatrix} + A^-(X^l, X^r) \begin{pmatrix} B^+ & -B^- \\ -B^- & B^+ \end{pmatrix} = (C^l, C^r), \end{cases}$$

i.e.,

$$\begin{cases} AXB = C, \\ A \begin{pmatrix} X^l, X^r \end{pmatrix} \begin{pmatrix} B^+ & -B^- \\ -B^- & B^+ \end{pmatrix} = (C^l, C^r). \end{cases}$$

□

In a similar way, we can obtain another model for solving the equation(5).

Theorem 3.4. The fuzzy linear system $A\tilde{X}B = \tilde{C}$ can be extended into the following system of linear matrix equations

$$\left\{ \begin{array}{l} AXB = C, \\ \left(\begin{array}{cc} A^+ & -A^- \\ -A^- & A^+ \end{array} \right) \left(\begin{array}{c} X^l \\ X^r \end{array} \right) B = \left(\begin{array}{c} C^l \\ C^r \end{array} \right), \end{array} \right. \quad (13)$$

where

$$\tilde{X} = (X, X^l, X^r), \tilde{C} = (C, C^l, C^r).$$

And the elements a_{ij}^+ of matrix A^+ and a_{ij}^- of matrix A^- are determined by the following way: if $a_{ij} \geq 0, a_{ij}^+ = a_{ij}$ else $a_{ij}^+ = 0, 1 \leq i, j \leq n$; if $a_{ij} < 0, a_{ij}^- = a_{ij}$ else $a_{ij}^- = 0, 1 \leq i, j \leq n$.

Proof. Let $\tilde{C} = (C, C^l, C^r) = (c_{ij}, c_{ij}^l, c_{ij}^r)_{n \times n}$, $\tilde{X} = (X, X^l, X^r) = (x_{ij}, x_{ij}^l, x_{ij}^r)_{n \times n}$. We also let $B = B^+ + B^-$ where the elements b_{ij}^+ of matrix B^+ and b_{ij}^- of matrix B^- are determined by the way: if $b_{ij} \geq 0, b_{ij}^+ = b_{ij}$ else $b_{ij}^+ = 0, 1 \leq i, j \leq n$; if $b_{ij} < 0, b_{ij}^- = b_{ij}$ else $b_{ij}^- = 0, 1 \leq i, j \leq n$.

For fuzzy matrix equation $A\tilde{X}B = \tilde{C}$, we can express it as

$$A(X, X^l, X^r)(B^+ + B^-) = (C, C^l, C^r). \quad (14)$$

Since

$$\tilde{x}_{ij}k = \begin{cases} (kx_{ij}, kx_{ij}^l, kx_{ij}^r), & k \geq 0, \\ (kx_{ij}, -kx_{ij}^r, -kx_{ij}^l), & k < 0 \end{cases},$$

we have

$$\tilde{X}B = \begin{cases} (XB, X^lB, X^rB), & B \geq 0, \\ (XB, -X^rB, -X^lB), & B < 0 \end{cases}$$

So the equation(14) can be rewritten as

$$A(X, X^l, X^r)B^+ + A(X, X^l, X^r)B^- = A(XB^+, X^lB^+, X^rB^+) + A(XB^-, -X^rB^-, -X^lB^-).$$

Since

$$k\tilde{x}_{ij} = \begin{cases} (kx_{ij}, kx_{ij}^l, kx_{ij}^r), & k \geq 0, \\ (kx_{ij}, -kx_{ij}^r, -kx_{ij}^l), & k < 0, \end{cases}$$

we have

$$A\tilde{X} = \begin{cases} (AX, AX^l, AX^r), & A \geq 0, \\ (AX, -AX^r, -AX^l), & A < 0. \end{cases}$$

Suppose $A = A^+ + A^-$, in which the elements a_{ij}^+ of matrix A^+ and a_{ij}^- of matrix A^- are determined by the way: if $a_{ij} \geq 0, a_{ij}^+ = a_{ij}$ else $a_{ij}^+ = 0, 1 \leq i, j \leq n$; if $a_{ij} < 0, a_{ij}^- = a_{ij}$ else $a_{ij}^- = 0, 1 \leq i, j \leq n$, we have

$$\begin{aligned} & (A^+XB^+, A^+X^lB^+, A^+X^rB^+) + (A^+XB^-, -A^+X^rB^-, -A^+X^lB^-) \\ & + (A^-XB^+, -A^-X^rB^+, -A^-X^lB^+) + (A^-XB^-, A^-X^lB^-, A^-X^rB^-) \\ & = (A^+XB^+ + A^-XB^+ + A^+XB^- + A^-XB^-, A^+X^lB^+ - A^-X^rB^+ - A^+X^rB^- \\ & + A^-X^lB^-, A^+X^rB^+ - A^-X^lB^+ - A^+X^lB^- + A^-X^rB^-) = (C, C^l, C^r). \end{aligned}$$

Thus we get

$$\left\{ \begin{array}{l} A^+XB^+ + A^-XB^+ + A^+XB^- + A^-XB^- = C, \\ A^+X^lB^+ - A^-X^rB^+ - A^+X^rB^- + A^-X^lB^- = C^l, \\ A^+X^rB^+ - A^-X^lB^+ - A^+X^lB^- + A^-X^rB^- = C^r, \end{array} \right. \quad (15)$$

which is the same as the equation(12).

Denoting (15) in matrix form, we have

$$\left\{ \begin{array}{l} (A^+ + A^-)X(B^+ + B^-) = C, \\ \left(\begin{array}{cc} A^+ & -A^- \\ -A^- & A^+ \end{array} \right) \left(\begin{array}{c} X^lB^+ \\ X^rB^+ \end{array} \right) + \left(\begin{array}{cc} A^+ & -A^- \\ -A^- & A^+ \end{array} \right) \left(\begin{array}{c} -X^rB^- \\ -X^lB^- \end{array} \right) = \left(\begin{array}{c} C^l \\ C^r \end{array} \right), \end{array} \right.$$

or

$$\left\{ \begin{array}{l} (A^+ + A^-)X(B^+ + B^-) = C, \\ \left(\begin{array}{cc} A^+ & -A^- \\ -A^- & A^+ \end{array} \right) \left(\begin{array}{c} X^l \\ X^r \end{array} \right) B^+ + \left(\begin{array}{cc} A^+ & -A^- \\ -A^- & A^+ \end{array} \right) \left(\begin{array}{c} X^l \\ X^r \end{array} \right) B^- = \left(\begin{array}{c} C^l \\ C^r \end{array} \right), \end{array} \right.$$

i.e.,

$$\left\{ \begin{array}{l} AXB = C, \\ \left(\begin{array}{cc} A^+ & -A^- \\ -A^- & A^+ \end{array} \right) \left(\begin{array}{c} X^l \\ X^r \end{array} \right) B = \left(\begin{array}{c} C^l \\ C^r \end{array} \right). \end{array} \right. .$$

□

By the matrix theory[13], we can prove the following result.

Theorem 3.5. *Let S belong to $R^{m \times n}$, T belong to $R^{p \times q}$ and C belong to $R^{m \times q}$. Then the minimal solution X^* of the matrix equation $SXT = C$ is expressed by*

$$X^* = S^\dagger CT^\dagger.$$

Proof. Let $Y = XT$, the matrix equation $SXT = C$ is expressed by $SY = C$. We can verify that the minimal solution Y^* of the matrix equation $SY = C$ is expressed by $Y^* = S^\dagger C$.

In fact, we write the matrix equation $SX = C$ in block form of matrix as follows:

$$SY_j = C_j, j = 1, 2, \dots, q$$

where Y_j, C_j are the j th of matrices Y, C respectively.

By the matrix theory, we know that the matrix equation $SY = C$ is consistent if and only if each linear equation $SY_j = C_j, j = 1, 2, \dots, q$ is consistent, and the matrix equation $SY = C$ is inconsistent if and only if at least one of linear equations $SY_j = C_j, j = 1, 2, \dots, q$ is inconsistent.

When $SY = C$ is consistent, the result that $Y^* = S^\dagger C$ is its minimal solution is straightforward.

When $SY = C$ is inconsistent, the expression

$$\|C - SY^*\|_F^2 = \min \|C - SY\|_F^2$$

holds is equivalent to the following conditions:

$$\|C - SY_j^*\|_F^2 = \min \|C_j - SY_j\|_F^2, j = 1, \dots, q$$

where $Y^* = [X_1^*, \dots, Y_q^*]$ is the least squares solution of the matrix equation $SY = C$ and Y_j^* is the least squares solution of linear equation $SY_j = C_j, j = 1, 2, \dots, q$ and

$$\|C - SY\|_F^2 = \sum_{j=1}^q \|C_j - SY_j\|_F^2 = \sum_{j=1}^q \sum_{i=1}^m \|c_{ij} - \sum_{k=1}^p s_{ik}y_{kj}\|_F^2.$$

From above analysis, we know that Y_j^* is the minimal solution of inconsistent linear equation $SY_j = C_j, j = 1, 2, \dots, q$ is equivalent to that $Y^* = [Y_1^*, \dots, Y_q^*]$ is the minimal solution of the inconsistent matrix equation $SY = C$. So we know that the minimal solution of the matrix equation $SY = C$ is expressed uniformly by $Y^* = S^\dagger C$.

On the other hand, we regard the matrix equation $XT = Y$ with $T^\top X^\top = Y^\top$ and denote it as $\overline{TX} = \overline{Y}$.

By the above analysis, we obtain that the minimal solution of the matrix equation $\overline{TX} = \overline{Y}$ is expressed uniformly by $\overline{X} = \overline{T}^\dagger \overline{Y}$, i.e., $X^* = YT^\dagger$. It means $X^* = YT^\dagger$ is the minimal solution of the matrix equation $XT = Y$.

Thus we prove that the minimal solution X^* of the matrix equation $SXT = C$ is expressed by $X^* = S^\dagger CT^\dagger$.

In order to solve the fuzzy matrix equation (5), we need to consider the systems of linear equations (10) or (13). It seems that we have obtained the minimal solutions of the linear system (10) and (13) as

$$\left\{ \begin{array}{l} X = A^\dagger CB^\dagger, \\ \left(\begin{array}{c} X^l \\ X^r \end{array} \right) = A^\dagger \left(\begin{array}{cc} C^l & C^r \end{array} \right) \left(\begin{array}{cc} B^+ & -B^- \\ -B^- & B^+ \end{array} \right)^\dagger \end{array} \right. \quad (16)$$

and

$$\left\{ \begin{array}{l} X = A^\dagger CB^\dagger, \\ \left(\begin{array}{c} X^l \\ X^r \end{array} \right) = \left(\begin{array}{cc} A^+ & -A^- \\ -A^- & A^+ \end{array} \right)^\dagger \left(\begin{array}{c} C^l \\ C^r \end{array} \right) B^\dagger, \end{array} \right. \quad (17)$$

where $(.)^\dagger$ is the Moore-Penrose generalized inverse of matrix $(.)$. □

However, the solution matrix may still not be an appropriate LR fuzzy numbers matrix except for $X^l \geq O, X^r \geq O$. Now we give the definition of LR fuzzy solution to the equation(5) as follows:

Definition 3.6. Let $\tilde{X} = (X, X^l, X^r)$. If (X, X^l, X^r) is the minimal solution of equation(10) or (13), such that $X^l \geq O, X^r \geq O$, we call $\tilde{X} = (X, X^l, X^r)$ is a strong LR fuzzy minimal solution of fuzzy matrix equation(5). Otherwise, $\tilde{X} = (X, X^l, X^r)$ is said a weak LR fuzzy minimal solution of fuzzy matrix equation(5) given by

$$\tilde{X} = \tilde{x}_{ij}$$

where

$$\tilde{x}_{ij} = \begin{cases} (x_{ij}, x_{ij}^l, x_{ij}^r), & x_{ij}^l > 0, \quad x_{ij}^r > 0, \\ (x_{ij}, 0, \max\{-x_{ij}^l, x_{ij}^r\}), & x_{ij}^l < 0, \quad x_{ij}^r > 0, \\ (x_{ij}, \max\{x_{ij}^l, -x_{ij}^r\}, 0), & x_{ij}^l > 0, \quad x_{ij}^r < 0, \\ (x_{ij}, -x_{ij}^l, -x_{ij}^r), & x_{ij}^l < 0, \quad x_{ij}^r < 0. \end{cases} \quad i, j = 1, \dots, n. \quad (18)$$

To illustrate the expression (16) or (17) to be a LR fuzzy solution matrix, we now discuss the generalized inverse of non negative matrix

$$S = \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix}$$

in a special structure.

Lemma 3.7. [13] Let

$$S = \begin{pmatrix} A^+ & -A^- \\ -A^- & A^+ \end{pmatrix}.$$

Then the matrix

$$S^\dagger = \begin{pmatrix} (A^+ - A^-)^\dagger + (A^+ + A^-)^\dagger & (A^+ - A^-)^\dagger - (A^+ + A^-)^\dagger \\ (A^+ - A^-)^\dagger - (A^+ + A^-)^\dagger & (A^+ - A^-)^\dagger + (A^+ + A^-)^\dagger \end{pmatrix} \quad (19)$$

is the Moore-Penrose inverse of the matrix S , where $(A^+ + A^-)^\dagger, (A^+ - A^-)^\dagger$ are the Moore-Penrose inverse of matrices $A^+ + A^-$ and $A^+ - A^-$, respectively.

The key point to make the solution matrix a strong LR fuzzy solution is that $\tilde{X} = (X, X^l, X^r)$ is LR fuzzy matrix, i.e., each element is a LR fuzzy number. By the following analysis, a sufficient condition is that $S^\dagger \geq O, B^\dagger \geq O$ or $A^\dagger \geq O, T^\dagger \geq O$.

Theorem 3.8. If

$$A^\dagger \geq O, \\ (B^+ - B^-)^\dagger + (B^+ + B^-)^\dagger \geq O, (B^+ - B^-)^\dagger - (B^+ + B^-)^\dagger \geq O,$$

the fuzzy matrix equation (2.5) has a strong LR fuzzy minimal solution as follows:

$$\tilde{X} = (X, X^l, X^r)$$

where

$$\begin{cases} X = A^\dagger C B^\dagger, \\ X^l = A^\dagger C^l E + A^\dagger C^r F, \\ X^r = A^\dagger C^l F + A^\dagger C^r E, \\ E = \frac{1}{2}((B^+ - B^-)^\dagger + (B^+ + B^-)^\dagger), \\ F = \frac{1}{2}((B^+ - B^-)^\dagger - (B^+ + B^-)^\dagger). \end{cases} \quad (20)$$

Proof. Since C^l and C^r are the left and right spreads fuzzy matrix \tilde{C} , $C^l \geq O$ and $C^r \geq O$. It means (C^l, C^r) is a non negative matrix.

Let

$$S^\dagger = \begin{pmatrix} E & F \\ F & E \end{pmatrix} = \frac{1}{2} \begin{pmatrix} (B^+ - B^-)^\dagger + (B^+ + B^-)^\dagger & (B^+ - B^-)^\dagger - (B^+ + B^-)^\dagger \\ (B^+ - B^-)^\dagger - (B^+ + B^-)^\dagger & (B^+ - B^-)^\dagger + (B^+ + B^-)^\dagger \end{pmatrix}.$$

We know the condition that $S^\dagger \geq 0$ is equivalent to $E \geq 0$ and $F \geq 0$.

Now that $E \geq O$ and $F \geq O$, the product of three non negative matrices

$$(X^l, X^r) = A^\dagger (C^l, C^r) \begin{pmatrix} B^+ & -B^- \\ -B^- & B^+ \end{pmatrix}^\dagger = A^\dagger (C^l, C^r) \begin{pmatrix} E & F \\ F & E \end{pmatrix} = (A^\dagger C^l E + A^\dagger C^r F, A^\dagger C^l F + A^\dagger C^r E) \geq O$$

is non negative in nature. It means that $X^l \geq O$ and $X^r \geq O$. □

For the model equation(13), we have the following result by the same analysis.

Theorem 3.9. *If*

$$B^\dagger \geq O, \\ (A^+ - A^-)^\dagger + (A^+ + A^-)^\dagger \geq O, (A^+ - A^-)^\dagger - (A^+ + A^-)^\dagger \geq O,$$

the fuzzy matrix equation (2.5) has a strong LR fuzzy minimal solution as follows:

$$\tilde{X} = (X, X^l, X^r)$$

where

$$\begin{cases} X = A^\dagger C B^\dagger, \\ X^l = E C^l B^\dagger + F C^r B^\dagger, \\ X^r = F C^l B^\dagger + E C^r B^\dagger, \\ E = \frac{1}{2}((A^+ - A^-)^\dagger + (A^+ + A^-)^\dagger), \\ F = \frac{1}{2}((A^+ - A^-)^\dagger - (A^+ + A^-)^\dagger). \end{cases} \quad (21)$$

Proof. The proof is straight forward. □

The following Theorems give some results for such S^{-1} and S^\dagger to be nonnegative. As usual, $(\cdot)^\top$ denotes the transpose of a matrix (\cdot) .

Theorem 3.10. [36] *The inverse S^{-1} of a nonnegative matrix S is nonnegative if and only if S is a generalized permutation matrix.*

Theorem 3.11. [14] *Let S be an $2n \times 2n$ nonnegative matrix with rank r . Then the following assertions are equivalent:*

- (a). $S^\dagger \geq 0$.
- (b). *There exists a permutation matrix P , such that PS has the form*

$$PS = \begin{pmatrix} T_1 \\ T_2 \\ \vdots \\ T_r \\ O \end{pmatrix},$$

where each T_i has rank 1 and the rows of T_i are orthogonal to the rows of T_j , whenever $i \neq j$, the zero matrix may be absent.

- (c). $S^\dagger = \begin{pmatrix} GC^\top & GD^\top \\ GD^\top & GC^\top \end{pmatrix}$ for some positive diagonal matrix G . In this case,

$$(C + D)^\dagger = G(C + D)^\top, (C - D)^\dagger = G(C - D)^\top.$$

4 Numerical examples

Example 4.1. *Consider the following fuzzy linear matrix system*

$$\begin{pmatrix} 1 & -1 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} \tilde{x}_{11} & \tilde{x}_{12} \\ \tilde{x}_{21} & \tilde{x}_{22} \end{pmatrix} \begin{pmatrix} -3 & 4 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} (3, 2, 1)_{LR} & (4, 1, 1)_{LR} \\ (5, 2, 2)_{LR} & (3, 1, 2)_{LR} \end{pmatrix}.$$

By the Theorem 3.1., the original fuzzy matrix equation is extended into the following a system of linear matrix equations(10)

$$\begin{cases} AXB = C, \\ A \begin{pmatrix} X^l, X^r \end{pmatrix} \begin{pmatrix} B^+ & -B^- \\ -B^- & B^+ \end{pmatrix} = \begin{pmatrix} C^l, C^r \end{pmatrix}, \end{cases}$$

where

$$\tilde{X} = (X, X^l, X^r), \tilde{C} = (C, C^l, C^r)$$

and

$$B^+ = \begin{pmatrix} 0 & 4 \\ 1 & 0 \end{pmatrix}, B^- = \begin{pmatrix} -3 & 0 \\ 0 & 0 \end{pmatrix}, C = \begin{pmatrix} 3 & 4 \\ 5 & 3 \end{pmatrix}, C^l = \begin{pmatrix} 2 & 1 \\ 2 & 1 \end{pmatrix}, C^r = \begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix}.$$

From (16), the solution of the computing model is

$$X = A^\dagger C B^\dagger = \begin{pmatrix} 1 & -1 \\ 0 & 2 \end{pmatrix}^\dagger \begin{pmatrix} 3 & 4 \\ 5 & 3 \end{pmatrix} \begin{pmatrix} -3 & 4 \\ 1 & 0 \end{pmatrix}^\dagger = \begin{pmatrix} 1.3750 & 9.6250 \\ 0.3750 & 3.6250 \end{pmatrix}$$

and

$$\begin{aligned} (X^l, X^r) &= A^\dagger (C^l, C^r) \begin{pmatrix} B^+ & -B^- \\ -B^- & B^+ \end{pmatrix}^\dagger = \begin{pmatrix} 1 & -1 \\ 0 & 2 \end{pmatrix}^\dagger \begin{pmatrix} 2 & 1 & 1 & 1 \\ 2 & 1 & 2 & 2 \end{pmatrix} \begin{pmatrix} 0 & 4 & 3 & 0 \\ 1 & 0 & 0 & 0 \\ 3 & 0 & 0 & 4 \\ 0 & 0 & 1 & 0 \end{pmatrix}^\dagger \\ &= \begin{pmatrix} 0.3750 & 1.5000 & 0.5000 & 0.8750 \\ 0.1250 & 0.2500 & 0.2500 & 0.6250 \end{pmatrix}. \end{aligned}$$

Since X^l, X^r are nonnegative matrices and $X - X^l > O$, the solution we obtained is an appropriate LR fuzzy matrix

$$\tilde{X} = (X, X^l, X^r) = \left(\begin{pmatrix} 1.3750 & 9.6250 \\ 0.3750 & 3.6250 \end{pmatrix}, \begin{pmatrix} 0.3750 & 1.5000 \\ 0.1250 & 0.2500 \end{pmatrix}, \begin{pmatrix} 0.5000 & 0.8750 \\ 0.2500 & 0.6250 \end{pmatrix} \right)$$

i.e.,

$$\tilde{X} = \begin{pmatrix} \tilde{x}_{11} & \tilde{x}_{12} \\ \tilde{x}_{21} & \tilde{x}_{22} \end{pmatrix} = \begin{pmatrix} (1.3750, 0.3750, 0.5000)_{LR} & (9.6250, 1.5000, 0.8750)_{LR} \\ (0.3750, 0.1250, 0.2500)_{LR} & (3.6250, 0.2500, 0.6250)_{LR} \end{pmatrix},$$

which admits a nonnegative strong LR fuzzy solution of the original fuzzy matrix system.

Example 4.2. Consider another fuzzy linear matrix system

$$\begin{pmatrix} 2 & 1 \\ 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \tilde{x}_{11} & \tilde{x}_{12} \\ \tilde{x}_{21} & \tilde{x}_{22} \end{pmatrix} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} = \begin{pmatrix} (2, 2, 1)_{LR} & (3, 2, 1)_{LR} \\ (3, 1, 1)_{LR} & (2, 1, 2)_{LR} \\ (1, 1, 1)_{LR} & (3, 2, 1)_{LR} \end{pmatrix}.$$

Suppose

$$\tilde{X} = (X, X^l, X^r),$$

$$A = A^+ + A^- = \begin{pmatrix} 2 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & -1 \\ 0 & 0 \end{pmatrix},$$

$$B = B^+ + B^- = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$$

and

$$\tilde{C} = (C, C^l, C^r) = \left(\begin{pmatrix} 2 & 3 \\ 3 & 2 \\ 1 & 3 \end{pmatrix}, \begin{pmatrix} 2 & 2 \\ 1 & 1 \\ 1 & 2 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 1 \end{pmatrix} \right).$$

By the Theorem 3.1., the original fuzzy matrix equation is extended into the following a system of linear matrix equations(10)

$$\begin{cases} AXB = C, \\ A (X^l, X^r) \begin{pmatrix} B^+ & -B^- \\ -B^- & B^+ \end{pmatrix} = (C^l, C^r). \end{cases}$$

From (16), the solution of the computing model is

$$X = A^\dagger C B^\dagger = \begin{pmatrix} 2 & 1 \\ 1 & -1 \\ 0 & 1 \end{pmatrix}^\dagger \begin{pmatrix} 2 & 3 \\ 3 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}^\dagger = \begin{pmatrix} 0.0179 & -0.0179 \\ -0.3393 & 0.3393 \end{pmatrix}$$

and

$$(X^l, X^r) = A^\dagger (C^l, C^r) \begin{pmatrix} B^+ & -B^- \\ -B^- & B^+ \end{pmatrix}^\dagger = \begin{pmatrix} 2 & 1 \\ 1 & -1 \\ 0 & 1 \end{pmatrix}^\dagger \begin{pmatrix} 2 & 2 & 1 & 1 \\ 1 & 1 & 1 & 2 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}^\dagger$$

$$= \begin{pmatrix} 0.4464 & 0.3571 & 0.3571 & 0.4464 \\ 0.0179 & 0.2143 & 0.2143 & 0.0179 \end{pmatrix}.$$

Since X^l, X^r are nonnegative matrices, the solution we obtained is an appropriate LR fuzzy matrix

$$\tilde{X} = (X, X^l, X^r) = \left(\begin{pmatrix} 0.0179 & -0.0179 \\ -0.3393 & 0.3393 \end{pmatrix}, \begin{pmatrix} 0.4464 & 0.3571 \\ 0.0179 & 0.2143 \end{pmatrix}, \begin{pmatrix} 0.3571 & 0.4464 \\ 0.2143 & 0.0179 \end{pmatrix} \right),$$

i.e.,

$$\tilde{X} = \begin{pmatrix} \tilde{x}_{11} & \tilde{x}_{12} \\ \tilde{x}_{21} & \tilde{x}_{22} \end{pmatrix} = \begin{pmatrix} (0.0179, 0.4464, 0.3571)_{LR} & (-0.0179, 0.3571, 0.4464)_{LR} \\ (-0.3393, 0.0179, 0.2143)_{LR} & (0.3393, 0.4464, 0.0179)_{LR} \end{pmatrix},$$

which admits a strong LR fuzzy solution of the original fuzzy linear matrix equation.

5 Conclusions

In this work we presented a model for solving fuzzy linear matrix equations $A\tilde{X}B = \tilde{C}$ where A and B are crisp $m \times m$ and $n \times n$ matrices respectively and \tilde{C} is an $m \times n$ arbitrary LR fuzzy numbers matrix. The model was made of two crisp systems of linear equations which determined the mean value and the left and right spreads of the solution. The LR fuzzy minimal solution of the fuzzy linear matrix equation was derived from solving the crisp systems of linear equations. In addition, the existence condition of strong LR fuzzy solution was studied. Numerical examples showed that our method is feasible to solve this type of fuzzy matrix equations.

Acknowledgement

The work is supported by the Natural Scientific Funds of PR China (no; 61967014; 1861059; 11761062) and the Scientific Research Project of Gansu Province Colleges and Universities (no.2019A-004; 2017A058).

References

- [1] S. Abbasbandy, A. Jafarian, R. Fzzati, *Conjugate gradient method for fuzzy symmetric positive definite system of linear equations*, Applied Mathematics and Computation, **171** (2005), 1184–1191.
- [2] S. Abbasbandy, R. Ezzati, A. Jafarian, *LU decomposition method for solving fuzzy system of linear equations*, Applied Mathematics and Computation, **172** (2006), 633–643.
- [3] S. Abbasbandy, M. Otadi, M. Mosleh, *Minimal solution of general dual fuzzy linear systems*, Chaos, Solitons and Fractals, **29** (2008), 638–652.
- [4] T. Allahviranloo, *Numerical methods for fuzzy system of linear equations*, Applied Mathematics and Computation, **153** (2004), 493–502.
- [5] T. Allahviranloo, *Successive over relaxation iterative method for fuzzy system of linear equations*, Applied Mathematics and Computation, **162** (2005), 189–196.
- [6] T. Allahviranloo, *The adomian decomposition method for fuzzy system of linear equations*, Applied Mathematics and Computation, **163** (2005), 553–563.
- [7] T. Allahviranloo, F. H. Lotfi, M. K. Kiasari, M. Khezerloo, *On the fuzzy solution of LR fuzzy linear systems*, Applied Mathematical Modelling, **37** (2013), 1170–1176.
- [8] T. Allahviranloo, M. Ghanbari, R. Nuraei, *A note on fuzzy linear systems*, Fuzzy Sets and Systems, **177**(1) (2011), 87–92.
- [9] T. Allahviranloo, N. Mikaeilvand, M. Barkhordary, *Fuzzy linear matrix equations*, Fuzzy Optimization and Decision Making, **8** (2009), 165–177.
- [10] M. Amirfakhrian, M. Fallah, R. Rodríguez-López, *A method for solving fuzzy matrix equations*, Soft Computing, **22** (2018), 2095–2103.

- [11] B. Asady, S. Abbasbandy, M. Alavi, *Fuzzy general linear systems*, Applied Mathematics and Computation, **169** (2005), 34–40.
- [12] N. Babbar, A. Kumar, Abhinav Bansal, *Solving fully fuzzy linear system with arbitrary triangular fuzzy numbers*, Soft Computing, **17** (2013) 691-702.
- [13] A. Ben-Israel, T. N. E. Greville, *Generalized inverses: Theory and applications*, second edition, Springer-Verlag, New York, 2003.
- [14] A. Berman, R. J. Plemmons, *Nonnegative matrices in the mathematical sciences*, Academic press, New York, 1979.
- [15] W. S. W. Daud, N. Ahmad, G. Malkawi, *A preliminary study on the solution of fully fuzzy Sylvester matrix equation*, In AIP Conference Proceedings, **1775** (2016), 30–36.
- [16] M. Dehghan, B. Hashemi, M. Ghatee, *Solution of the full fuzzy linear systems using iterative techniques*, Chaos, Solitons and Fractals, **34** (2007) 316–336.
- [17] M. Dehghan, B. Hashemi, *Iterative solution of fuzzy linear systems*, Applied Mathematics and Computation, **175** (2006), 645–674.
- [18] K. Dookhitram, R. Lollchund, R. K. Tripathi, M. Bhuruth, *Fully fuzzy Sylvester matrix equation*, Journal of Intelligent and Fuzzy Systems, **28** (2015), 2199–2211.
- [19] D. Dubois, H. Prade, *Operations on fuzzy numbers*, Journal of Systems Science, **9** (1978), 613–626.
- [20] S. A. Edalatpanah, *Modified iterative methods for solving fully fuzzy linear systems*, In I, Management Association (Ed.) Fuzzy Systems: Concepts, Methodologies, Tools, and Applications, (2017), 55–73.
- [21] M. Friedman, M. Ma, A. Kandel, *Fuzzy linear systems*, Fuzzy Sets and Systems, **96** (1998), 201–209.
- [22] R. Ghanbari, *Solutions of fuzzy LR algebraic linear systems using linear programs*, Applied Mathematical Modelling, **39** (2015), 5164-5173.
- [23] R. Goetschel, W. Voxman, *Elementary calculus*, Fuzzy Sets and Systems, **18** (1986), 31–43.
- [24] Z. T. Gong, X. B. Guo, *Inconsistent fuzzy matrix equations and its fuzzy least squares solutions*, Applied Mathematical Modelling, **35** (2011), 1456–1469.
- [25] Z. T. Gong, X. B. Guo, K. Liu, *Approximate solution of dual fuzzy matrix equations*, Information Sciences, **266** (2014), 112-133.
- [26] X. B. Guo, D. Q. Shang, *Approximate solutions of LR fuzzy Sylvester matrix equations*, Journal of Applied Mathematics, **2013** (2013), 1–10.
- [27] X. B. Guo, K. Zhang, *Solving fuzzy matrix equation of the form $\tilde{X}A = \tilde{B}$* , Journal of Intelligent and Fuzzy Systems, **32** (2017), 2771–2778.
- [28] X. B. Guo, Y. L. Han, *Further investigation to dual fuzzy matrix equation*, Journal of Intelligent and Fuzzy Systems, **33** (2017), 2617–2629.
- [29] J. J. Kaur, A. Kumar, *A note on approximate solution of dual fuzzy matrix equations*, Information Sciences, **418** (2017), 184–185.
- [30] M. Ma, M. Friedman, A. Kandel, *Duality in fuzzy linear systems*, Fuzzy Sets and Systems, **109** (2000), 55–58.
- [31] S. Nahmias, *Fuzzy variables*, Fuzzy Sets and Systems, **1**(2) (1978) 97–111.
- [32] H. S. Najafi, S. A. Edalatpanah, *An improved model for iterative algorithms in fuzzy linear systems*, Computational Mathematics and Modeling, **24** (2013), 443–451.
- [33] H. S. Najafi, S. A. Edalatpanah, *H-Matrices in fuzzy linear systems*, International Journal of Computational Mathematics, **6** (2013), 52–56.

- [34] R. Nuraei, T. Allahviranloo, M. Ghanbari, *Finding an inner estimation of the solution set of a fuzzy linear system*, Applied Mathematical Modelling, **37** (2013), 5148–5161.
- [35] M. Otadi, M. Mosleh, *Solving fully fuzzy matrix equations*, Applied Mathematical Modelling, **36** (2012), 6114–6121.
- [36] R. J. Plemmons, *Regular nonnegative matrices*, Proceedings of the American Mathematical Society, **39** (1973), 26–32.
- [37] M. L. Puri, D. A. Ralescu, *Differentials for fuzzy functions*, Journal of Mathematics Analysis and Application, **91** (1983), 552–558.
- [38] C. X. Wu, M. Ma, *Embedding problem of fuzzy number space: Part I*, Fuzzy Sets and Systems, **44** (1991), 33–38.
- [39] C. X. Wu, M. Ma, *Embedding problem of fuzzy number space: Part III*, Fuzzy Sets and Systems, **46** (1992), 281–286.
- [40] L. A. Zadeh, *Fuzzy sets*, Information and Control, **8** (1965), 338–353.
- [41] B. Zheng, K. Wang, *General fuzzy linear systems*, Applied Mathematics and Computation, **181** (2006), 1276–1286.

Solving LR fuzzy linear matrix equation^{*}

X. Guo and D. Shang

حل معادله ماتریسی خطی فازی LR

چکیده. در این مقاله، معادله ماتریسی فازی $A \tilde{X} B = \tilde{D}$ که در آن A ، B به ترتیب ماتریس‌های قطعی $n \times n$ و \tilde{C} یک ماتریس اعداد فازی LR دلخواه است، مورد بررسی قرار گرفته است. یک روش عددی برای محاسبه جواب فازی طراحی شده است و یک شرط کافی برای وجود جواب فازی قوی بدست آمده است. مثال‌هایی جهت توضیح روش پیشنهادی ارائه گردیده‌اند.