

C^∞ L -fuzzy manifolds with L -gradation of openness and C^∞ LG -fuzzy mappings of them

M. Mostafavi¹

¹Department of Mathematics, University of Qom, Qom, Iran

mmostafavi14279@gmail.com

Abstract

In this paper, we generalize all of the fuzzy structures which we have discussed in [14] to L -fuzzy set theory, where $L = \langle L, \leq, \wedge, \vee, ' \rangle$ denotes a complete distributive lattice with at least two elements. We define the concept of an LG -fuzzy topological space (X, \mathfrak{T}) which X is itself an L -fuzzy subset of a crisp set M and \mathfrak{T} is an L -gradation of openness of L -fuzzy subsets of M which are less than or equal to X . Then we define C^∞ L -fuzzy manifolds with L -gradation of openness and C^∞ LG -fuzzy mappings of them such as LG -fuzzy immersions and LG -fuzzy imbeddings. We fuzzify the concept of the product manifolds with L -gradation of openness and define LG -fuzzy quotient manifolds when we have an equivalence relation on M and investigate the conditions of the existence of the quotient manifolds. We also introduce LG -fuzzy immersed, imbedded and regular submanifolds.

Keywords: C^∞ LG -fuzzy n -manifolds, C^∞ LG -fuzzy mappings, LG -fuzzy quotient manifolds, LG -fuzzy immersion, regular LG -fuzzy submanifolds.

1 Introduction

The concept of fuzzy sets was introduced by Zadeh [21]. Then Chang [1] confined his attention to the more basic concepts of general topology and generalized them to fuzzy topological spaces. In his definition, fuzziness in the concept of openness of a fuzzy subset, is absent. In consequence of the development of fuzzy topology, many authors like Wong [19], Lowen [10] introduced various concepts of fuzzy topology. R. Lowen [11] suggested that the properties should be considered fuzzy, that is, one should be able to measure a degree to which a property holds. E. Lowen and R. Lowen [12] considered compactness degrees, and in [20], investigated measures of separation in $[0,1]$ -topological spaces. In 1985, Shostak [15] gave a new definition of fuzzy topology by introducing a concept of gradation of openness of fuzzy subsets of X . Later, Chattopadhyay [2] et al. attempted to introduce a concept of gradation of openness of a fuzzy set of X by a map $\tau : I^X \rightarrow I$ satisfying three weaker conditions than [15] and later in [3] made a slight modification in their definition and rediscovered the Shostak's concept of fuzzy topology. Gregori [7] proved that each gradation of openness δ is the supremum (infimum) of a strictly increasing (decreasing) sequence of gradations of openness which are equivalent to δ . Stadler and Vicente [13] have introduced a new concept of fuzzy topological subspace over each fuzzy subset from the fuzzy topology δ , which coincides with the usual definition in the case that $\mu = X^Y$, $Y \subset X$. In [16], Shostak developed a theory of compactness degrees and connectedness degrees in $[0,1]$ -fuzzy topological spaces, and in [17], brought up a theory of degrees of precompactness and completeness in the so-called Hutton fuzzy uniform spaces. In 2016 Ibedou [9] discussed graded fuzzy topological spaces. While all of the researches about the C^1 or C^∞ fuzzy manifolds, focused on a crisp set, in [14] and in this paper, we demonstrate the possibility of improving current definitions using a new method. In [14], we investigated some properties of a novel fuzzy topological space (X, τ) , where X is itself a fuzzy subset of a crisp set M . Perhaps the most important generalization of the aforementioned structures in [14], is the consideration of lattice L beyond the unit interval $I = [0, 1]$. Let $L = \langle L, \leq, \wedge, \vee, ' \rangle$ be a complete distributive lattice set with at least 2 elements; 0 is the bottom element and 1 is the top element of L . An L -fuzzy

subset D of the crisp set M , in Goguen's sense [6], is a function $D : M \rightarrow L$ and is denoted by $D \in L^M$. In this manuscript, we define the concept of L -fuzzy topological space (X, \mathfrak{T}) with the L -gradation of openness, where X is an L -fuzzy subset of a crisp set M . We introduce C^∞ L -fuzzy manifolds (X, \mathfrak{T}) with L -gradation of openness, called C^∞ LG -fuzzy manifolds, with a different perception from [5] and [4] and obtain C^∞ n -premanifolds of them. We define C^∞ LG -fuzzy mappings of C^∞ LG -fuzzy manifolds and prove the LG -fuzzy rank theorem. Then we define and discuss LG -fuzzy immersions and LGP -fuzzy imbedding functions. We proceed to define the LG -fuzzy immersed, imbedded submanifolds as well as LG -fuzzy regular submanifolds, and then some theorems about the relations between them are deduced.

2 Preliminaries

Definition 2.1. Let X be an L -fuzzy subset of M . Then any L -fuzzy subset of M which is less than or equal to X is called an L -fuzzy subset of X . We denote the set of all L -fuzzy subsets of X by L_X^M . If τ as a collection of L -fuzzy subsets of X , satisfies the following conditions, then (X, τ) is called an L -fuzzy topological space (L -fts):

- 1) $X, \phi \in \tau$,
- 2) $\{A_i\}_{i \in I} \subseteq \tau \Rightarrow \bigcup_{i \in I} A_i \in \tau$,
- 3) $A, B \in \tau \Rightarrow A \cap B \in \tau$.

Example 2.2. Let $M = \mathbb{R}^n$ and $X = 1$ be a constant L -fuzzy subset of M . Let $B(a, r, b)$ be an L -fuzzy subset that is equal to zero outside or on the sphere $B(a, r)$ and equal to the function b with values in L , inside $B(a, r)$. We call the L -fuzzy topology induced by

$$\beta_{L_n} = \{B(a, r, b), a \in \mathbb{R}^n, r \in \mathbb{R}^+, b : B(a, r) \rightarrow L, \text{ is a function}\},$$

the L -fuzzy Euclidean topology of dimension n and denote it by τ_{L_n} . Therefore we have the L -fuzzy Euclidean topological space $(1_{\mathbb{R}^n}, \tau_{L_n})$.

Definition 2.3. Let $\mathfrak{T} : L_X^M \rightarrow L$, be a mapping satisfying:

- i) $\mathfrak{T}(X) = \mathfrak{T}(\tilde{0}) = 1$,
- ii) $\mathfrak{T}(A \cap B) \geq \mathfrak{T}(A) \wedge \mathfrak{T}(B)$,
- iii) $\mathfrak{T}(\bigcup_{j \in J} A_j) \geq \bigwedge_{j \in J} \mathfrak{T}(A_j)$.

Then \mathfrak{T} is called an L -gradation of openness on X and (X, \mathfrak{T}) is called an LG -fuzzy topological space (L -gfts). Let $x \in M$ and $A \in L_X^M$. When we write $x \in A$, we mean $x \in \text{supp}A$.

Example 2.4. Let $M = \mathbb{R}^n$ and $X = 1$ be a constant L -fuzzy subset of M . As three useful examples, we define

$$\mathfrak{T}_{L_n} : L_X^M \rightarrow L, \quad \mathfrak{T}_{L_n}(B) = \begin{cases} 1 & B \in \tau_{L_n}, \\ 0 & \text{elsewhere.} \end{cases} \quad (1)$$

and

$$\mathfrak{T}_{L_{\text{sup}}} : L_X^M \rightarrow L, \quad \mathfrak{T}_{L_{\text{sup}}}(B) = \begin{cases} 1 & B = \tilde{0}, \\ \sup\{B(x) : x \in M\} & \tilde{0} \neq B \in \tau_{L_n}, \\ 0 & \text{elsewhere,} \end{cases} \quad (2)$$

If we set "inf" instead of "sup" in the above definition, then we have L -gradation of openness $\mathfrak{T}_{L_{\text{inf}}}$.

Let \mathfrak{T}_{L_n} be any L -gradation of openness on $1_{\mathbb{R}^n}$, such that $\text{supp}\mathfrak{T} = \tau_{L_n}$, then we call $(1_{\mathbb{R}^n}, \mathfrak{T}_{L_n})$ the LG -fuzzy Euclidean topological space.

Definition 2.5. Let (X, \mathfrak{T}) be an L -gfts. Set $\text{supp}\mathfrak{T} = \{A \in L_X^M : \mathfrak{T}(A) > 0\}$, then A is called an LG -open subset of X if $A \in \text{supp}\mathfrak{T}$. Furthermore

- 1) Suppose $x \in X$ and $V \in L_X^M$. If there exists an LG -open subset U of X such that $U(x) = V(x)$ and $U \leq V$, then V is called an LG -neighborhood of x in X . We denote the set of all LG -neighborhoods of x in X by $LGN(x)$.

2) If for all $x, y \in X, x \neq y$, there exist two LG -neighborhoods $U_x \in LGN(x), U_y \in LGN(y)$ such that $U_x \cap U_y = \emptyset$. Then (X, \mathfrak{T}) is called a Hausdorff L -gfts.

3) For each L -fuzzy subset A of X and any $U \subset \text{supp}X$, we define the L -fuzzy subset $\chi_{U,A}$ of X by:

$$\chi_{U,A}(z) = \begin{cases} A(z) & z \in U, \\ 0 & \text{elsewhere.} \end{cases}$$

From now on, we write χ_U instead of $\chi_{U,X}$.

4) A is called an LG -closed subset of X if $X - A \in \text{supp}\mathfrak{T}$.

5) Let Z be an LG -open subset of X . Define $\mathfrak{T}_Z : L_X^M \rightarrow L$, by $\mathfrak{T}_Z(A) = \mathfrak{T}(A)$. Then (Z, \mathfrak{T}_Z) is called an LG -fuzzy topological subspace of X (L -gtfss).

Definition 2.6. If $\mathfrak{C} : L_X^M \rightarrow L$, satisfies the following conditions:

- i) $\mathfrak{C}(X) = \mathfrak{C}(\tilde{0}) = 1$.
- ii) $\mathfrak{C}(A \cup B) \geq \mathfrak{C}(A) \wedge \mathfrak{C}(B)$.
- iii) $\mathfrak{C}(\bigcap_{j \in J} A_j) \geq \bigwedge_{j \in J} \mathfrak{C}(A_j)$.

Then \mathfrak{C} is called an L -gradation of closedness on X .

Proposition 2.7. Let \mathfrak{C} and \mathfrak{T} be L -gradations of closedness and openness respectively on X . Then

- i) The mapping $\mathfrak{T}_{\mathfrak{C}} : L_X^M \rightarrow L$, defined by $\mathfrak{T}_{\mathfrak{C}}(A) = \mathfrak{C}(X - A)$, is an L -gradation of openness on X , where $(X - A)$ is an L -fuzzy subset of M defined by $(X - A)(p) = X(p) - A(p)$.
- ii) The mapping $\mathfrak{C}_{\mathfrak{T}} : L_X^M \rightarrow L$, defined by $\mathfrak{C}_{\mathfrak{T}}(A) = \mathfrak{T}(X - A)$, is an L -gradation of closedness on X .
- iii) We have $\mathfrak{C}_{\mathfrak{T}_{\mathfrak{C}}} = \mathfrak{C}, \mathfrak{T}_{\mathfrak{C}_{\mathfrak{T}}} = \mathfrak{T}$.

The proof is straightforward.

Proposition 2.8. Let $\mathfrak{M}_{\mathfrak{T}}(X)$ be the set of all L -gradations of openness on X . We write $\mathfrak{T}_1 \leq \mathfrak{T}_2$, if we have $\mathfrak{T}_1(A) \leq \mathfrak{T}_2(A), \forall A \in L_X^M$. Then $(\mathfrak{M}_{\mathfrak{T}}(X), \leq)$ is a complete lattice.

Proof. It is clear that the relation \leq between the functions from L_X^M to L , is an equivalence relation. Therefore $(\mathfrak{M}_{\mathfrak{T}}(X), \leq)$ is a partially ordered set. Further we define two mappings $\mathfrak{T}_0, \mathfrak{T}_1 : L_X^M \rightarrow L$, by

$$\mathfrak{T}_0(\tilde{0}) = \mathfrak{T}_0(X) = 1, \mathfrak{T}_0(A) = 0, \forall A \in L_X^M - \{\tilde{0}, X\}, \mathfrak{T}_1(A) = 1, \forall A \in L_X^M.$$

Then $\mathfrak{T}_0, \mathfrak{T}_1$ are two L -gradations of openness on X and we have:

$$\mathfrak{T}_0(A) \leq \mathfrak{T}(A) \leq \mathfrak{T}_1(A), \forall A \in L_X^M.$$

Hence $\mathfrak{T}_0, \mathfrak{T}_1$ are minimal and maximal elements of $\mathfrak{M}_{\mathfrak{T}}(X)$, respectively.

An arbitrary intersection of gradations of openness on X , is a gradation of openness. Thus any subset of $\mathfrak{M}_{\mathfrak{T}}(X)$, has a lower bound in it. To prove this, let $\{\mathfrak{T}_k, k \in K\}$, be an arbitrary family of L -gradations of openness on X . We show that $\mathfrak{T} = \bigwedge_{k \in K} \mathfrak{T}_k$ is an L -gradation of openness on X . Obviously, $\mathfrak{T}(X) = \mathfrak{T}(\tilde{0}) = 1$. Also,

$$\mathfrak{T}\left(\bigcup_j A_j\right) = \bigwedge_k \mathfrak{T}_k\left(\bigcup_j A_j\right) \geq \bigwedge_k \left(\bigwedge_j \mathfrak{T}_k(A_j)\right) = \bigwedge_j \left(\bigwedge_k \mathfrak{T}_k(A_j)\right) = \bigwedge_j (\mathfrak{T}(A_j)),$$

and

$$\mathfrak{T}(A \cap B) = \bigwedge_k \mathfrak{T}_k(A \cap B) \geq \bigwedge_k (\mathfrak{T}_k(A) \wedge \mathfrak{T}_k(B)) \geq \bigwedge_k \mathfrak{T}_k(A) \wedge \bigwedge_k \mathfrak{T}_k(B) \geq \mathfrak{T}(A) \wedge \mathfrak{T}(B).$$

This completes the proof. □

Archive of SID

Example 2.9. Consider $(1_{\mathbb{R}^n}, \mathfrak{T}_{In})$ and $0 \in \mathbb{R}^n$. We show that the fuzzy point 0_1 is an IG-closed subset of $1_{\mathbb{R}^n}$: The fuzzy point $0_1 = \chi_{\{0\}}$ is an I-fuzzy subset of \mathbb{R}^n . So,

$$(1_{\mathbb{R}^n} - 0_1)(x) = 1 - \chi_{\{0\}}(x) = \begin{cases} 1 & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Therefore,

$$(1_{\mathbb{R}^n} - 0_1)(x) = \bigcup_{0 \neq k \in \mathbb{Z}} B(k, 1, 1)(x).$$

Hence, $(1_{\mathbb{R}^n} - 0_1) \in \tau_{In}$. Thus $\mathfrak{T}_{In}(1_{\mathbb{R}^n} - 0_1) \geq 0$. So, $1_{\mathbb{R}^n} - 0_1$ is an IG-open set. Hence, 0_1 is an IG-closed subset.

Definition 2.10. Let (X, \mathfrak{T}) be a fuzzy topological space and A, B be any fuzzy subsets of X ,

- 1) A fuzzy subset V of X is called an LG-neighborhood of A if there exists an LG-open subset U such that $A \leq U \leq V$. We denote the set of all LG-neighborhoods of A by $LGN(A)$.
- 2) Let $B \leq A$. Then B is called an LG-interior set of A if $A \in LGN(B)$. The union of all LG-interior sets of A is denoted by LGA° .
- 3) The intersection of all LG-closed subsets containing A is called an LG-closure of A and is denoted by $LG\bar{A}$.
- 4) x is called an LG-boundary point of A if for every LG-neighborhood V of x , we have $V \not\leq A$. The set of these points is called an LG-boundary of A and is denoted by $LG\partial A$.
- 5) If x belongs to the LG-closure of $A - \chi_{\{x, A\}}$, then x is called an LG-limited point of A and the set of these points is denoted by LGA' .
- 6) A is said to be an LG-dense subset of X , if $LG\bar{A} = X$.

From now on, we suppose that M_1, M_2 are two crisp sets, $X \in L^{M_1}, Y \in L^{M_2}$ and $(X, \mathfrak{T}), (Y, \mathfrak{R})$ are two LG-fuzzy topological spaces.

Definition 2.11. Let $f : M_1 \rightarrow M_2$ be a function and $f[X]$ be an L-fuzzy subset of M_2 , defined by

$$f[X](y) = \bigvee \{X(x) \mid x \in f^{-1}(y)\}.$$

If we have $f[X] \leq Y$, then f is called an LG-related function from X to Y and the set of all such functions is denoted by $LGRf(X, Y)$. Furthermore, if we have $\mathfrak{R}(H) \leq \mathfrak{T}(f^{-1}[H])$ for all LG-fuzzy subset H of Y , then f is an L-gradation-preserving LG-related function so it is called an LGP-related function or LGP-fuzzy mapping from X to Y , $f \in LGRf(X, Y)$.

- i) f is called a one-to-one LG-related (LGP-related) function if $f|_{\text{supp}X} : \text{supp}X \rightarrow \text{supp}Y$ is a one-to-one function.
- ii) f is called an onto LG-related (LGP-related) function if $f[X] = Y$.

Remark 2.12. Let $A \in \text{supp}\mathfrak{T}$ and $B \in \text{supp}\mathfrak{R}$. Let f be an LGP-fuzzy mapping from X to Y such that $f[A] \leq B$. Then we have $\mathfrak{R}(H) \leq \mathfrak{T}(f^{-1}[H])$ for each LG-fuzzy subset of Y and in particular $H \leq B$. Thus $\mathfrak{R}_B(H) \leq \mathfrak{T}_A(f^{-1}[H])$ for each LG-fuzzy subset H of Y with $H \leq B$. Therefore f can be considered as an LGP-fuzzy mapping of two L-gfts's, (A, \mathfrak{T}_A) and (B, \mathfrak{R}_B) . So we can write $f \in LGPRf(A, B)$.

Definition 2.13. Let $f \in LGRf(X, Y)$, then

- i) f is called LG-open if $f[A] \in \text{supp}\mathfrak{R} - \{\tilde{0}, Y\}$, $\forall A \in \text{supp}\mathfrak{T} - \{\tilde{0}, X\}$ and $f[X] \in \text{supp}\mathfrak{R}$.
- ii) f is called LG-continuous if $f^{-1}[H] \in \text{supp}\mathfrak{T} - \{\tilde{0}, X\}$, $\forall H \in \text{supp}\mathfrak{R} - \{\tilde{0}, Y\}$ and $f^{-1}[Y] \in \text{supp}\mathfrak{T}$.
- iii) f is called an LG-homeomorphism if it is one-to-one, onto, LG-continuous, LG-open and $f^{-1} \in LGRf(Y, X)$.
- iv) f is called an LGP-homeomorphism if it is bijective and f, f^{-1} are LGP-fuzzy mapping.

Proposition 2.14. Let A, B be LG-open subsets of X, Y respectively. Let $\psi : M_1 \rightarrow M_2$ be a function. Then ψ is an LGP-homeomorphism from A to B if and only if ψ satisfies the two following conditions:

- i) $A(p) = B(\psi(p))$ for all $p \in A$ or $B(q) = A(\psi^{-1}(q))$ for all $q \in B$
- ii) $\mathfrak{R}(H) = \mathfrak{T}(\psi^{-1}[H])$ for all LG -fuzzy subset H of B .

Proof. Let ψ satisfies conditions (i) and (ii), then by Definition 2.11 and Remark 2.12 we have $\psi \in LGPRf(A, B)$ and ψ is an LGP -homeomorphism from A to B . Conversely suppose ψ is an LGP -homeomorphism from A to B . We prove that ψ satisfies (i) and (ii).

i) Since ψ is bijective, for any $q \in B$, there is exactly one element $p \in A$, such that $\psi^{-1}(q) = \{p\}$. So we have $\psi[A](q) = \sup\{A(a) \mid a \in \psi^{-1}(q)\} = A(p)$. On the other hand by Definition 2.11, we have $\psi[A] \leq B$. Hence $A(p) \leq B(q)$. We see $\psi^{-1}[B](p) = B(\psi(p)) = B(q)$. Since by Definition 2.13 (iv), we have $\psi^{-1} \in LGPRf(Y, X)$, then $\psi^{-1}[B] \leq A$. Hence $B(q) \leq A(p)$. Therefore $A(p) = B(q)$. Therefore $A(p) = B(\psi(p))$, for all $p \in A$ and $B(q) = A(\psi^{-1}(q))$, for all $q \in B$. Thus $A = (\psi^{-1}[B])$ and $\psi[A] = B$.

ii) Since $\psi \in LGPRf(A, B)$, we have $\mathfrak{R}(H) \leq \mathfrak{T}(\psi^{-1}[H])$ for all LG -fuzzy subset H of Y , and since $\psi^{-1} \in LGPRf(B, A)$, we have $\mathfrak{T}(D) \leq \mathfrak{R}(\psi[D])$. Set $\psi[D] = H$. Then $D = \psi^{-1}[H]$ by injectivity of ψ . So $\mathfrak{T}(D) \leq \mathfrak{R}(H)$. Hence we have $\mathfrak{T}(\psi^{-1}[H]) = \mathfrak{R}(H)$. \square

Proposition 2.15. *Every LGP -fuzzy mapping from X to Y is an LG -continuous related function, but the converse is not true.*

Proof. Let f be an LGP -fuzzy mapping from X to Y , then $\forall H \in \text{supp}\mathfrak{R} - \{\tilde{0}, Y\}$, we have $0 < \mathfrak{R}(H) \leq \mathfrak{T}(f^{-1}[H])$. Hence $f^{-1}[H] \in \text{supp}\mathfrak{T} - \{\tilde{0}, X\}$. Therefore f is LG -continuous.

Conversely, we define an LG -continuous function which is not an LGP -fuzzy mapping:

Following Example 2.4, consider $f = id : (1_{\mathbb{R}^n}, \mathfrak{T}_{Ln}) \rightarrow (1_{\mathbb{R}^n}, \mathfrak{T}_{Lsup})$. Since $f[1_{\mathbb{R}^n}] = 1_{\mathbb{R}^n}$ and we have

$$f^{-1}[H] = H \in \text{supp}\mathfrak{T}_{Ln} - \{\tilde{0}, X\} = \tau_{Ln} - \{\tilde{0}, 1_{\mathbb{R}^n}\}, \quad \forall H \in \text{supp}\mathfrak{T}_{Lsup} - \{\tilde{0}, Y\} = \tau_{Ln} - \{\tilde{0}, 1_{\mathbb{R}^n}\},$$

and $f^{-1}[1_{\mathbb{R}^n}] \in \text{supp}\mathfrak{T}_{Ln}$. Therefore f is LG -continuous. Now Let $A = \begin{cases} x^2 & x \in (0, \frac{1}{2}), \\ 0 & \text{elsewhere.} \end{cases}$ Then $A \in \tau_{Ln}$ and $\mathfrak{T}_{Lsup}(f[A]) = \frac{1}{4}$. But $\mathfrak{T}_{Ln}(A) = 1$. Hence the condition $\mathfrak{T}_{Ln}(A) \leq \mathfrak{T}_{Lsup}(f[A])$ dose not hold. Hence f is not an LGP -fuzzy mapping. \square

3 L -fuzzy topological manifolds with L -gradation of openness

Definition 3.1. *Let \mathfrak{T} be an L -gradation of openness on X . Then (X, \mathfrak{T}) is an LG -fuzzy topological space of dimension n , if for any $x \in X$, there exists an LG -open subset A of X containing x and an LG -open subset B of $(1_{\mathbb{R}^n}, \mathfrak{T}_{Ln})$, together with an LGP -homeomorphism $\psi \in LGPRf(A, B)$. The pair (A, ψ) is called an LG -local coordinate neighborhood of each $q \in A$ and we assign to q the n LG -local coordinates $x_1(q), x_2(q), \dots, x_n(q)$ of its image $\psi(q)$ in \mathbb{R}^n .*

Definition 3.2. *Let $\mathfrak{A} = \{(A_i, \psi_i) \mid i \in J\}$ be a collection of LG -local coordinate neighborhoods. Since ψ_i is an LGP -homeomorphism for all $i \in J$, then for all $i, j \in J$ whenever $A_i \cap A_j \neq \phi$,*

$$\psi_j \circ \psi_i^{-1} : \psi_i(\text{supp}(A_i \cap A_j)) \rightarrow \psi_j(\text{supp}(A_i \cap A_j)),$$

is an LGP -homeomorphism, that is called an LG -transition function. -

$$\psi_j \circ \psi_i^{-1}(x_1^i, x_2^i, \dots, x_n^i) = (x_1^j, x_2^j, \dots, x_n^j).$$

If $\psi_i \circ \psi_j^{-1}$ and $\psi_j \circ \psi_i^{-1}$ changing the LG -local coordinates are infinitely differentiable or C^∞ , we shall say that (A_i, ψ_i) is C^∞ compatible with (A_j, ψ_j) whenever $A_i \cap A_j \neq \phi$.

Definition 3.3. *An LG -fuzzy topological space (X, \mathfrak{T}) is called an LG -fuzzy topological manifold of dimension n , if it satisfies the two following conditions:*

- i) X is an LG -fuzzy topological space of dimension n ,
- ii) X is a Hausdorff L -gfts.

Definition 3.4. *A differentiable or C^∞ LG -fuzzy structure on an LG -fuzzy topological manifold (X, \mathfrak{T}) , is a family $\mathfrak{A} = \{(A_\alpha, \psi_\alpha), \alpha \in J\}$ of LG -local coordinate neighborhoods such that*

Archive of SID

- 1) $X = \bigcup_{\alpha \in J} A_\alpha$;
- 2) Each pair (A_α, ψ_α) and (A_β, ψ_β) are compatible for all $\alpha, \beta \in J$.
- 3) Any LG-local coordinate neighborhood (V, φ) that is compatible with every $(A_\alpha, \psi_\alpha), \alpha \in J$ is in \mathfrak{A} itself.

A C^∞ LG-fuzzy manifold (X, \mathfrak{T}) is an LG-fuzzy topological manifold with a C^∞ LG-fuzzy structure on it. In what follows, for convenience, "LG-fuzzy manifold with LG-fuzzy structure" will mean C^∞ LG-fuzzy manifold with C^∞ LG-fuzzy structure,

Example 3.5. Let $M = \mathbb{R}^3, X : \mathbb{R}^3 \rightarrow I, X(x) = \begin{cases} 1 & \|x\| = 1, \\ 0 & \|x\| \neq 1. \end{cases}$ Then $suppX = S^2$, the unit sphere. Set

$$\mathfrak{T} : I_X^M \rightarrow I, \quad \mathfrak{T}(A) = \begin{cases} sup\{A(x) \mid x \in X\} & A \in \tau_{In}, A \leq X, \\ 0 & \text{elsewhere.} \end{cases}$$

Then (X, \mathfrak{T}) is an IG-fuzzy manifold of dimension 2.

Proof. Let $J = \{1, 2, 3\}$. We define six IG-open subsets covering X by:

$$\forall x = (x_1, x_2, x_3), \quad A_j^\pm(x) = \begin{cases} \pm x_j & \pm x_j > 0, \|x\| = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Then we show that all A_j^\pm are diffeomorphic to IG-open subset $B : \mathbb{R}^2 \rightarrow I$, defined by:

$$\forall y = (y_1, y_2), \quad B(y) = \begin{cases} \sqrt{1 - y_1^2 - y_2^2} & \|y\| < 1, \\ 0 & \text{otherwise.} \end{cases}$$

Since $suppB = B(0, 1)$, so $B \in \tau_{I2}$. We define six bijections ψ_j^\pm from $suppA_j^\pm = \{(x_1, x_2, x_3) \mid \pm x_j > 0, \|x\| = 1\}$ to $suppB = \{(y_1, y_2) \mid \|y\| < 1\}$, for all $j \in J$ by:

$$\psi_1^\pm(x_1, x_2, x_3) = (x_2, x_3), \quad (\psi_1^\pm)^{-1}(y_1, y_2) = (\pm\sqrt{1 - y_1^2 - y_2^2}, y_1, y_2).$$

$$\psi_2^\pm(x_1, x_2, x_3) = (x_1, x_3), \quad (\psi_2^\pm)^{-1}(y_1, y_2) = (y_1, \pm\sqrt{1 - y_1^2 - y_2^2}, y_2).$$

$$\psi_3^\pm(x_1, x_2, x_3) = (x_1, x_2), \quad (\psi_3^\pm)^{-1}(y_1, y_2) = (y_1, y_2, \pm\sqrt{1 - y_1^2 - y_2^2}).$$

Also, it is seen that $\psi_j^\pm \circ (\psi_i^\pm)^{-1}$ is infinitely differentiable for all $i, j \in J$. For example:

$$\psi_2^\pm \circ (\psi_1^\pm)^{-1}(y_1, y_2) = \psi_2^\pm(\pm\sqrt{1 - y_1^2 - y_2^2}, y_1, y_2) = (\pm\sqrt{1 - y_1^2 - y_2^2}, y_2).$$

Therefore, each pair (A_i^\pm, ψ_i^\pm) and (A_j^\pm, ψ_j^\pm) are compatible, for all $i, j \in J$. We see

$$\forall j \in J, \quad A_j^\pm(x) = \pm x_j = B_j(\psi_j^\pm(x)), \quad \forall x \in A_j^\pm.$$

Let H be an IG-fuzzy subset of $1_{\mathbb{R}^2}$ with $H \leq B$. We show that $\mathfrak{T}((\psi_j^\pm)^{-1}[H]) = \mathfrak{T}_{I\text{sup}}(H)$. Using (2), we have $\mathfrak{T}_{I\text{sup}}(H) = sup\{H(a) \mid a \in \mathbb{R}^2\}$. Since ψ_j^\pm is bijective, for each $a \in \mathbb{R}^2$, there exists one and only one element $p \in suppA_j^\pm$ such that $\psi_j^\pm(p) = a$ or $(\psi_j^\pm)^{-1}(a) = p$. Hence

$$\mathfrak{T}_{I\text{sup}}(H) = sup\{H(\psi_j^\pm(p)) \mid p \in suppA_j^\pm\} = sup\{((\psi_j^\pm)^{-1}[H])(p) \mid p \in suppA_j^\pm\} = \mathfrak{T}((\psi_j^\pm)^{-1}[H]).$$

Hence $\psi_j^\pm \in IGPRf(A_j^\pm, B)$ is an IGP-homeomorphism for all $j \in J$ and this completes the proof. □

Example 3.6. The set of natural numbers, \mathbb{N} , partially ordered by divisibility, is a distributive lattice set, for which the unique supremum is the least common multiple and the unique infimum is the greatest common divisor. Let $L = \mathbb{N} \cup \{\infty\}$. Then L is a complete lattice. Notice that we denote the top element of any lattice by 1, but in this example, ∞ is the top element of $\mathbb{N} \cup \{\infty\}$. We define the LG-fuzzy Euclidean topological space $(1_{\mathbb{R}^{mn}}, \mathfrak{T}_{L\text{sup}})$ by

$$1_{\mathbb{R}^{mn}} : \mathbb{R}^{mn} \rightarrow L, \quad 1_{\mathbb{R}^{mn}}((a_1, a_2, \dots, a_{mn})) = \infty,$$

$$\mathfrak{T}_{Lmn} : L_{\mathbb{R}^{mn}}^{\mathbb{R}^{mn}} \rightarrow L, \quad \mathfrak{T}_{Lmn}(D) = \begin{cases} \infty & D \in \tau_{Lmn}, \\ 0 & \text{elsewhere.} \end{cases}$$

Let $M = \mathcal{M}_{m \times n}(\mathbb{R})$ and $X \in L^M$ be defined by

$$X((a_{ij})) = 2 + \max\{\lfloor |a_{ij}| \rfloor \mid 1 \leq i \leq m, 1 \leq j \leq n\},$$

where $\lfloor |x| \rfloor$ is equal to the greatest integer less than or equal to $|x|$. There is a bijection ψ from M to \mathbb{R}^{mn} :

$$\psi(a_{ij}) = (a_{11}, \dots, a_{1n}, \dots, a_{m1}, \dots, a_{mn}).$$

Hence using ψ and (1), we define

$$\mathfrak{T} : L_X^M \rightarrow L, \quad \mathfrak{T}(A) = \begin{cases} \infty & \psi[A] \in \tau_{Lmn}, \\ 0 & \text{elsewhere.} \end{cases}$$

We show that (X, \mathfrak{T}) is an C^∞ LG -fuzzy mn -manifold. Let B is an L -fuzzy subset of \mathbb{R}^{mn} defined by

$$B((a_1, a_2, \dots, a_{mn})) = 2 + \max\{\lfloor |a_k| \rfloor \mid 1 \leq k \leq mn\}.$$

We see $B = \psi[X]$. Since $\text{supp} B = \mathbb{R}^{mn} = \bigcup_{k=1}^\infty B(0, k)$, Hence $B \in \tau_{Lmn}$. Therefore using (1), for each LG -open subset H of \mathbb{R}^{mn} , with $H \leq B$, we have $\mathfrak{T}_{Lmn}(H) = \infty = \mathfrak{T}(\psi^{-1}[H])$. So by Proposition 2.14, $\psi \in LGPRf(X, B)$ is an LG -homeomorphism. We can cover (X, \mathfrak{T}) by the single LG -coordinate neighborhood (X, ψ) . Hence (X, \mathfrak{T}) is an C^∞ LG -fuzzy mn -manifold.

Definition 3.7. (*LG-open submanifolds*) Let Z be an LG -open subset of the LG -fuzzy manifold (X, \mathfrak{T}) . If $\mathfrak{A} = \{(A_\alpha, \psi_\alpha), \alpha \in J\}$ is an LG -fuzzy structure on X , then (Z, \mathfrak{T}_Z) is an LG -fuzzy topology with LG -fuzzy structure consisting of the LG -coordinate neighborhoods $(A_\alpha \cap Z, \psi_\alpha|_{A_\alpha \cap Z})$.

Example 3.8. Let (X, \mathfrak{T}) be as Example 3.6. We define $Z : \mathcal{M}_{m \times n}(\mathbb{R}) \rightarrow L$, $Z(A) = \begin{cases} X(A) & \det A \neq 0, \\ 0 & \det A = 0. \end{cases}$

We have $Z \leq X$ and $U = \text{supp} Z = Gl(n, \mathbb{R})$ is an open subset of $\mathcal{M}_{m \times n}$. Hence we can prove that Z is an LG -open subset of X . Therefore (Z, \mathfrak{T}_Z) is an LG -fuzzy submanifold of (X, \mathfrak{T}) with the single LG -local coordinate neighborhood $(Z, \psi|_Z)$ where ψ is a bijection defined in Example 3.6.

Example 3.9. Let $L = \mathbb{N} \cup \{\infty\}$ and $M = \mathbb{R}^{n+1}$. Define an L -fuzzy subset

$$X : M \rightarrow L, \quad X(x) = \begin{cases} n + 2 & \|x\| = 1, \\ 0 & \|x\| \neq 1, \end{cases}$$

$$\mathfrak{T} : L_X^M \rightarrow L, \quad \mathfrak{T}(A) = \begin{cases} \infty & A \in \tau_{L(n+1)}, A \leq X, \\ 0 & \text{elsewhere.} \end{cases}$$

Then $\text{supp} X = S^n$. Then (X, \mathfrak{T}) is an LG -fuzzy manifold of dimension n :

Proof. Let $J = \{1, \dots, n + 1\}$. We define $2(n + 1)$ LG -open subsets covering X , $A_j^\pm : M \rightarrow L$, $j \in J$ by:

$$\forall x = (x_1, \dots, x_{n+1}), \quad A_j^\pm(x) = \begin{cases} j & \pm x_j > 0, \|x\| = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Then we show that each A_j^\pm is LG -homeomorphic to the LG -open subset $B_j : \mathbb{R}^n \rightarrow L$ defined by:

$$\forall y = (y_1, \dots, y_n), \quad B_j(y) = \begin{cases} j & \|y\| < 1, \\ 0 & \text{otherwise.} \end{cases}$$

with $2(n + 1)$ LG -maps $\psi_i^\pm : A_j^\pm \rightarrow B_j$ defined by:

$$\psi_j^\pm(x_1, \dots, x_{n+1}) = (x_1, \dots, \hat{x}_j, \dots, x_{n+1}).$$

$$(\psi_j^\pm)^{-1}(y_1, \dots, y_n) = (y_1, \dots, \pm \sqrt{1 - (y_1^2 + \dots + y_n^2)}, \dots, y_n),$$

Archive of SID

where \hat{x}_j means omit x_j . Also it is seen that $\psi_j^\pm \circ (\psi_i^\pm)^{-1}$ is infinitely differentiable for all $i, j \in J$ and we have

$$A_j^\pm(x) = j = B_j(\psi_j^\pm(x)), \quad \forall x \in A_j^\pm \text{ and } \forall j \in J.$$

Also using (1), for each LG -open subset H_j of $1_{\mathbb{R}^n}$, with $H_j \leq B_j$, we have

$$\mathfrak{T}_{L_n}(H_j) = \infty = \mathfrak{T}(\psi_j^\pm)^{-1}(H_j) \quad \forall j \in J.$$

Therefore by Proposition 2.14, $\psi_i^\pm \in LGPRf(A_i^\pm, B_j)$, is an LGP -homeomorphism for all $j \in J$. \square

Theorem 3.10. *Let (M, τ) be a C^∞ ordinary n -manifold with the C^∞ structure $\mathfrak{W} = \{(U_k, \psi_k), k \in K\}$. Consider $X = \tilde{1}$, the constant L -fuzzy subset of M . Let δ be the L -fuzzy topology on X generated by $\{\chi_{U_k}, k \in K\}$. Define*

$$\mathfrak{T}_\tau : L_X^M \rightarrow L, \quad \mathfrak{T}_\tau(A) = \begin{cases} 1 & A \in \delta, \\ 0 & \text{elsewhere.} \end{cases}$$

Then (X, \mathfrak{T}_τ) is an LG -fuzzy manifold of dimension n .

Proof. First we show that \mathfrak{T}_τ is an L -gradation of openness on X :

- 1) Since $\tilde{0}, X \in \delta$, therefore $\mathfrak{T}_\tau(\tilde{0}) = \mathfrak{T}_\tau(X) = 1$.
- 2) For each family of fuzzy subsets $\{A_j\} \subseteq L_X^M$, we have two cases:
 - i) $\{A_j\}_{j \in J} \subseteq \delta \Rightarrow \bigcup_{j \in J} A_j \in \delta$ and $\mathfrak{T}_\tau(A_j) = 1, \forall j \in J \Rightarrow 1 = \mathfrak{T}_\tau(\bigcup_{j \in J} A_j) \geq \bigwedge_{j \in J} \mathfrak{T}_\tau(A_j) = 1$
 - ii) $A_j \subseteq (L_X^M - \delta)$, for some $j \in J \Rightarrow \mathfrak{T}_\tau(A_j) = 0$, for some $j \in J \Rightarrow \mathfrak{T}_\tau(\bigcup_{j \in J} A_j) \geq \bigwedge_{j \in J} \mathfrak{T}_\tau(A_j) = 0$
- 3) For every two fuzzy subsets $A, B \in L_X^M$, we have three cases:
 - i) $A, B \in \delta \Rightarrow A \cap B \in \delta \Rightarrow 1 = \mathfrak{T}_\tau(A \cap B) \geq \mathfrak{T}_\tau(A) \wedge \mathfrak{T}_\tau(B) = 1$;
 - ii) $A \in (L_X^M - \delta), B \in \delta \Rightarrow \mathfrak{T}_\tau(A) = 0, \mathfrak{T}_\tau(B) = 1 \Rightarrow \mathfrak{T}_\tau(A \cap B) \geq 0 = \mathfrak{T}_\tau(A) \wedge \mathfrak{T}_\tau(B)$;
 - iii) $A, B \in (L_X^M - \delta) \Rightarrow \mathfrak{T}_\tau(A) = \mathfrak{T}_\tau(B) = 0 \Rightarrow \mathfrak{T}_\tau(A \cap B) \geq 0 = \mathfrak{T}_\tau(A) \wedge \mathfrak{T}_\tau(B)$.

Next we prove that (X, \mathfrak{T}_τ) is an LG -fuzzy manifold:

Let $p \in M$. Then there exists an open subset $U_k, k \in K$ s.t. $p \in U_k$ and an open set V_k of \mathbb{R}^n along with a homeomorphism $\psi_k : U_k \rightarrow V_k$. Let τ^\flat be the L -fuzzy topology on $1_{\mathbb{R}^n}$ generated by $\{\chi_{V_k}, k \in K\}$. Then $\tau^\flat \subseteq \tau_{L_n}$. Hence we can consider the restriction of \mathfrak{T}_{L_n} on τ^\flat . We see

$$\mathfrak{T}_\tau(\chi_{U_k}) = \mathfrak{T}_{L_n}|_{\tau^\flat}(\chi_{V_k}).$$

Define LGP -homeomorphisms

$$\psi_k^\flat : M \rightarrow \mathbb{R}^n, \quad \psi_k^\flat(p) = \psi_k(p)\chi_{U_k} = \begin{cases} \psi_k(p) & \text{if } p \in U_k, \\ 0 & \text{elsewhere.} \end{cases}$$

Therefore $\psi_k^\flat \in LGPRf(\chi_{U_k}, \chi_{V_k})$. Thus $\mathfrak{W}_\tau = \{(\chi_{U_k}, \psi_k^\flat), k \in K\}$ is an LG -fuzzy structure on X . \square

Theorem 3.11. *Let (X, \mathfrak{T}) be an LG -fuzzy n -manifold with LG -fuzzy structure $\mathfrak{A} = \{(A_j, \psi_j), j \in J\}$.*

If $\mathfrak{T}^\flat = \{\text{supp}A \mid \mathfrak{T}(A) > 0\}$, then $(\text{supp}X, \mathfrak{T}^\flat)$ is a topological manifold of dimension n called a premanifold with the structure $\mathfrak{A}^\flat = \{(\text{supp}A_j, \psi_j|_{\text{supp}A_j}), j \in J\}$ called a prestructure.

Proof. Since $\text{Dom}\mathfrak{T} = L_X^M$, then for all $A \in \text{supp}\mathfrak{T}$, we have A is less than X . Hence $\text{supp}A \subseteq \text{supp}X$.

- i) $\mathfrak{T}(\tilde{0}) = \mathfrak{T}(X) = 1 \Rightarrow \phi = \text{supp}\tilde{0} \in \text{supp}\mathfrak{T}^\flat$ and $\text{supp}X \in \text{supp}\mathfrak{T}^\flat$.
- ii) Let $\{A_j, j \in J\} \subseteq \delta$, Then $\mathfrak{T}(A_j) > 0, \forall j \in J$. Hence $\mathfrak{T}(\bigcup_{j \in J} A_j) \geq \bigwedge_{j \in J} \mathfrak{T}(A_j) > 0$. Thus $\bigcup_{j \in J} A_j \in \text{supp}\mathfrak{T}^\flat$.
- iii) Since $\mathfrak{T}(A \cap B) \geq \mathfrak{T}(A) \wedge \mathfrak{T}(B)$. So $A, B \in \mathfrak{T}^\flat$ implies that $\mathfrak{T}(A \cap B) > 0$. Thus $A \cap B \in \text{supp}\mathfrak{T}^\flat$.

Therefore \mathfrak{T}^\flat is a topology on $\text{supp}X$. Let $p \in X$. Then there exists an LG -open subset $A_k, k \in K$ such that $p \in A_k$ and there exists an LG -open subset $B_k, k \in K$ with an LGP -homeomorphism $\psi_k \in LGPRf(A_k, B_k)$. Hence $p \in \text{supp}X, A_k(p) = B_k(\psi_k(p))$ and $\psi_k|_{\text{supp}A_k} : \text{supp}A_k \rightarrow \text{supp}B_k$ is one-to-one and onto. Therefore $(\text{supp}A_k, \psi_k|_{\text{supp}A_k})$ is a coordinate neighborhood of p . Also $\psi_j \circ \psi_i^{-1}$ is infinitely differentiable for all $i, j \in K$, thus $(\psi_j|_{\text{supp}(A_i \cap A_j)}) \circ (\psi_i^{-1}|_{\text{supp}(B_i \cap B_j)})$ is C^∞ for all $i, j \in K$. Hence $(\text{supp}X, \mathfrak{T}^\flat)$ is a premanifold. www.SID.ir

Theorem 3.12. Let (M, τ) be an n -manifold with structure $\mathfrak{W} = \{(U_j, \varphi_j), j \in J\}$. Let $X = \tilde{1}$. Then $(suppX, (\tau^\triangleright)^\triangleleft) = (M, \tau)$.

Proof. It is clear that $suppX = M$ and for every $U \in \tau$ we have $supp\chi_U = U$. Hence $(\tau^\triangleright)^\triangleleft = \tau$. Since we have $\mathfrak{W}^\triangleright = \{(\chi_{U_j}, \varphi_j^\triangleright), j \in J\}$, then

$$(\mathfrak{W}^\triangleright)^\triangleleft = \{(supp\chi_{U_j}, \varphi_j^\triangleright|_{supp\chi_{U_j}}), j \in J\} = \{(U_j, \varphi_j^\triangleright|_{U_j}), j \in J\} = \{(U_j, \varphi_j), j \in J\} = \mathfrak{W}.$$

□

Remark 3.13. Let (X, \mathfrak{T}) be an L-fuzzy topological space with L-gradation of openness, then $\mathfrak{T}_{\mathfrak{T}^\triangleleft}$ does not necessarily equal \mathfrak{T} . We show it by the following example.

Example 3.14. Let $M = \mathbb{R}$. We define

$$X : M \rightarrow I, \quad X(x) = \begin{cases} \frac{1}{[x]} & x \in (2, +\infty), \\ 0 & \text{elsewhere.} \end{cases}$$

and $\mathfrak{T} : I_X^M \rightarrow I$, by $\mathfrak{T}(A) = \begin{cases} 1 & A \in \tau_{11}, A \leq X \\ 0 & \text{elsewhere.} \end{cases}$.

Then clearly (X, \mathfrak{T}) is an IG-fuzzy manifold of dimension 1. Then by Theorem 3.11, $(suppX, \mathfrak{T}^\triangleleft)$ is a manifold, where $\mathfrak{T}^\triangleleft = \{suppA \mid A \in supp\mathfrak{T}\}$. Hence by Theorem 3.10 we have $\mathfrak{T}_{\mathfrak{T}^\triangleleft} = \{\chi_{suppA} \mid A \in supp\mathfrak{T}\}$. We have $A(x) \leq X(x) \leq \frac{1}{2}, \forall x \in M$ and $\forall A \in \tau_{11}$. Hence $1 = \chi_{suppA} \neq A$. Therefore $\mathfrak{T}_{\mathfrak{T}^\triangleleft} \neq \mathfrak{T}$.

4 LG-fuzzy quotient manifolds

Definition 4.1. Let M be a crisp set and \sim be an equivalence relation on it. If A is an L-fuzzy subset of M such that $A(y) = A(x)$ whenever $y \sim x$, then we define the L-fuzzy subset:

$$\frac{A}{\sim} : \frac{M}{\sim} \rightarrow L, \quad \frac{A}{\sim}([x]) = A(x), \quad \forall x \in M,$$

where $[x] = \{y \mid x \sim y\}$. Since $A \leq X$, thus $\frac{A}{\sim} \leq \frac{X}{\sim}$ and hence $\frac{A}{\sim} \in L_{\frac{X}{\sim}}^{\frac{M}{\sim}}$.

Theorem 4.2. Let (X, \mathfrak{T}) be an LG-fuzzy topological space, such that $X(y) = X(x)$ whenever $y \sim x$, then $(\frac{X}{\sim}, \frac{\mathfrak{T}}{\sim})$ is an LG-fuzzy topological space, called the LG-fuzzy quotient space, where

$$\frac{\mathfrak{T}}{\sim} : L_{\frac{X}{\sim}}^{\frac{M}{\sim}} \rightarrow L, \quad \frac{\mathfrak{T}}{\sim}(\frac{A}{\sim}) = \mathfrak{T}(A).$$

Proof. We show that all elements of $L_{\frac{X}{\sim}}^{\frac{M}{\sim}}$ are in the form $\frac{A}{\sim}$ for some $A \in L_X^M$. Let B be an L-fuzzy subset of $\frac{M}{\sim}$ less than $\frac{X}{\sim}$. We define L-fuzzy subset A of M by $A(x) = B([x]), \forall x \in M$. Let $x \sim y$ so $[x] = [y]$, then $A(x) = B([x]) = B([y]) = A(y)$ and thus $\frac{A}{\sim} = B$. Also,

1) $\frac{\mathfrak{T}}{\sim}(\frac{\tilde{0}}{\sim}) = \mathfrak{T}(\tilde{0}) = 1, \quad \frac{\mathfrak{T}}{\sim}(\frac{X}{\sim}) = \mathfrak{T}(X) = 1.$

2) $\frac{\mathfrak{T}}{\sim}(\frac{A_1}{\sim} \cap \frac{A_2}{\sim}) = \frac{\mathfrak{T}}{\sim}(\frac{A_1 \cap A_2}{\sim}) = \mathfrak{T}(A_1 \cap A_2) \geq \mathfrak{T}(A_1) \wedge \mathfrak{T}(A_2) = \frac{\mathfrak{T}}{\sim}(\frac{A_1}{\sim}) \wedge \frac{\mathfrak{T}}{\sim}(\frac{A_2}{\sim}).$

3) Let $\{A_j\}_{j \in J}$ be a sequence of L-fuzzy subsets of X , such that $\forall j \in J, A_j(y) = A_j(x)$, whenever $y \sim x$, then

$$\bigcup_{j \in J} \frac{A_j}{\sim} [y] = sup\{ \frac{A_j}{\sim} [y], j \in J \} = sup\{ A_j(y), j \in J \} = sup\{ A_j(x), j \in J \} = \bigcup_{j \in J} \frac{A_j}{\sim} [x],$$

$$\frac{\mathfrak{T}}{\sim}(\bigcup_{j \in J} \frac{A_j}{\sim}) = \frac{\mathfrak{T}}{\sim}(\frac{\bigcup_{j \in J} A_j}{\sim}) = \mathfrak{T}(\bigcup_{j \in J} A_j) \geq \bigwedge_{j \in J} \mathfrak{T}(A_j) = \bigwedge_{j \in J} \frac{\mathfrak{T}}{\sim}(\frac{A_j}{\sim}).$$

Hence $\frac{\mathfrak{I}}{\sim}$ is a gradation of openness on $\frac{X}{\sim}$. □

This LG-fuzzy topology is nontrivial when $L = I$, because for each $\alpha \in I$, $\alpha X(x) = \alpha X(y)$ whenever $x \sim y$. Hence $\alpha X \in \text{supp} \frac{\mathfrak{I}}{\sim}$.

Definition 4.3. Consider an LG-fuzzy quotient space $(\frac{X}{\sim}, \frac{\mathfrak{I}}{\sim})$. The equivalence relation \sim is called an LG-open relation if for each fuzzy subset $A \in \text{supp} \mathfrak{I}$ we have $\frac{A}{\sim} \in \text{supp} \frac{\mathfrak{I}}{\sim}$.

Theorem 4.4. Let (X, \mathfrak{I}) be an LG-fuzzy manifold and \sim be an LG-open relation. Then $(\frac{X}{\sim}, \frac{\mathfrak{I}}{\sim})$ is an LG-fuzzy topological space of dimension n called LG-fuzzy quotient topological space of dimension n . If $\text{supp} \mathfrak{I}$ has a countable basis, then $\frac{\mathfrak{I}}{\sim}$ has a countable basis.

Proof. Let (A, ψ) be an LG-locally coordinate neighborhood of $p \in X$ and $\psi \in \text{LGPRf}(A, B)$. Since \sim is an LG-open relation, then we have $\frac{\mathfrak{I}}{\sim}(\frac{A}{\sim}) > 0$. We define a corresponding relation \sim^* on $\text{supp} B$ as follows:

$$a \sim^* b \iff \psi^{-1}(a) \sim \psi^{-1}(b) \text{ for all } a, b \in \text{supp} B.$$

Clearly \sim^* is a reflexive and symmetric relation. Let $a \sim^* b$ and $b \sim^* c$. Then we have $\psi^{-1}(a) \sim \psi^{-1}(b)$, and $\psi^{-1}(b) \sim \psi^{-1}(c)$. Since \sim is transitive, so $\psi^{-1}(a) \sim \psi^{-1}(c)$. Hence $a \sim^* c$. Therefore \sim^* is transitive and so it is an equivalence relation. Since we have

$$a \sim^* b \implies \psi^{-1}(a) \sim \psi^{-1}(b) \implies A(\psi^{-1}(a)) = A(\psi^{-1}(b)) \implies B(a) = B(b).$$

Therefore, $\frac{B}{\sim^*}$ is well-defined. Now we define

$$\frac{\psi}{\sim} : \text{supp}(\frac{A}{\sim}) \rightarrow \text{supp}(\frac{B}{\sim^*}), \quad \frac{\psi}{\sim}([p]) = [\psi(p)].$$

We see

$$[p] = [q] \iff p \sim q \iff \psi(p) \sim^* \psi(q) \iff [\psi(p)] = [\psi(q)] \iff \frac{\psi}{\sim}([p]) = \frac{\psi}{\sim}([q]).$$

Therefore, $\frac{\psi}{\sim}$ is a well-defined and one to one function. Since ψ is onto, we see

$$\forall a \in \text{supp} B, \exists p \in \text{supp} A \text{ s.t. } a = \psi(p) \implies \frac{\psi}{\sim}([p]) = [\psi(p)] = [a].$$

Hence, $\frac{\psi}{\sim}$ is onto. We have

$$\frac{B}{\sim^*}([a]) = B(a) = A(\psi^{-1}(a)) = \frac{A}{\sim}([\psi^{-1}(a)]) = \frac{A}{\sim}((\frac{\psi}{\sim})^{-1}([a])).$$

On the other hand for each LG-open subset H of $1_{\mathbb{R}^n}$ which $H \leq B$, we have

$$\frac{\mathfrak{I}_{Ln}}{\sim}(\frac{H}{\sim}) = \mathfrak{I}_{Ln}(H) = 1 = \mathfrak{I}(\psi^{-1}(H)) = \frac{\mathfrak{I}}{\sim}(\frac{\psi^{-1}(H)}{\sim}).$$

Hence, $\frac{\psi}{\sim} \in \text{LGPRf}(\frac{A}{\sim}, \frac{B}{\sim^*})$. Therefore, if $\mathfrak{A} = \{(A_j, \psi_j), j \in J\}$ be a C^∞ LG-structure of X . Then $\frac{\mathfrak{A}}{\sim} = \{(\frac{A_j}{\sim}, \frac{\psi_j}{\sim}), j \in J\}$ is a C^∞ LG-fuzzy structure of $(\frac{X}{\sim}, \frac{\mathfrak{I}}{\sim})$. Now, suppose that $\text{supp} \mathfrak{I}$ has a countable basis $\beta = \{A_i, i \in \mathbb{N}\}$. Let $\frac{A}{\sim} \in \text{supp} \frac{\mathfrak{I}}{\sim}$ and $A = \bigcup_{j \in K} A_j$. Since $K \subseteq \mathbb{N}$. Hence for all $y \in X$ we have:

$$\frac{A}{\sim}([y]) = A(y) = \bigcup_{j \in J} A_j(y) = \text{sup}\{A_j(y), j \in K\} = \text{sup}\{\frac{A_j}{\sim}([y]), j \in K\} = \bigcup_{j \in K} \frac{A_j}{\sim}([y]).$$

Therefore, $\frac{\beta}{\sim} = \{\frac{A_i}{\sim}, i \in K\}$ is a countable basis for $\text{supp} \frac{\mathfrak{I}}{\sim}$.

Archive of SID

Example 4.5. Consider the IG-fuzzy Euclidean topological space $(1_{\mathbb{R}}, \mathfrak{T}_{I1})$. We define a relation on \mathbb{R} as follows:
 $\forall x, y \in 1_{\mathbb{R}}, x \sim y$ if $x - y \in \mathbb{Z}$. Since $1_{\mathbb{R}}(y) = 1_{\mathbb{R}}(x) = 1$ whenever $y \sim x$. Hence $\frac{1_{\mathbb{R}}}{\sim}$ is well-defined and hence by Theorem 4.2 we have the IG-fuzzy quotient topological space $(\frac{1_{\mathbb{R}}}{\sim}, \frac{\mathfrak{T}_{I1}}{\sim})$. We show that it is an IG-fuzzy topological manifold:

Let $A : \mathbb{R} \rightarrow I, A(x) = x - [x]$. Since

$$suppA = \mathbb{R} - \mathbb{Z} = \bigcup_{k \in \mathbb{Z}} (k, k + 1) = \bigcup_{k \in \mathbb{Z}} B(k + \frac{1}{2}, \frac{1}{2}).$$

Then we see:

$$A = \bigcup_{k \in \mathbb{Z}} B(k + \frac{1}{2}, \frac{1}{2}, b_k), \text{ where } b_k : B(k + \frac{1}{2}, \frac{1}{2}) \rightarrow I, b_k(x) = x - [x].$$

Therefore by Example 2.2, $A \in \tau_{I1}$ and hence by Example 2.4, $\mathfrak{T}_{I1}(A) = 1$. Since for all $x, y \in \mathbb{R}$ we have

$$x \sim y \Rightarrow y = x + k, k \in \mathbb{Z} \Rightarrow [y] = k + [x].$$

$$y - [y] = k + x - (k + [x]) = x - [x] \Rightarrow A(y) = A(x). \tag{3}$$

Hence $\frac{A}{\sim}$ is well-defined. So $\frac{\mathfrak{T}_{I1}}{\sim}(\frac{A}{\sim}) = \mathfrak{T}_{I1}(A) = 1$. Define

$$B : \mathbb{R} \rightarrow I, B(y) = \begin{cases} y & y \in (0, 1), \\ 0 & \text{otherwise.} \end{cases}$$

We can write $B = B(\frac{1}{2}, \frac{1}{2}, b)$, where $b : B(\frac{1}{2}, \frac{1}{2}) \rightarrow I, b(y) = y$. So $B \in \tau_{I1}$. Hence $\mathfrak{T}_{I1}(B) = 1 = \frac{\mathfrak{T}_{I1}}{\sim}(\frac{A}{\sim})$.

We define $\psi : supp\frac{A}{\sim} \rightarrow suppB$ by $\psi[x] = x - [x]$. If $[x] = [y]$, then it means that $x \sim y$, so by (3), we have $x - [x] = y - [y]$. Hence $\psi([x]) = \psi([y])$. Therefore ψ is well-defined. We show that ψ is injective:

$$\psi([x]) = \psi([y]) \Rightarrow x - [x] = y - [y] \Rightarrow y - x = [y] - [x] \in \mathbb{Z} \Rightarrow x \sim y \Rightarrow [x] = [y].$$

Since $\forall t \in (0, 1), [t] = 0$, so $\psi([t]) = t - [t] = t$. Therefore ψ is surjective. So by Definition 2.11 we have $\frac{A}{\sim}([x]) = B(\psi([x]))$. On the other hand for each LG-open subset H of $1_{\mathbb{R}^n}$ with $H \leq B$, we have

$$\frac{\mathfrak{T}_{I1}}{\sim}(\psi^{-1}(H)) = \frac{\mathfrak{T}_{I1}}{\sim}(\frac{H}{\sim}) = \mathfrak{T}_{I1}(H) = 1.$$

Hence $\psi \in IGPRf(\frac{A}{\sim}, B)$. Thus we have a single IG-local coordinate neighborhood $(\frac{A}{\sim}, \psi)$ for all points of $\frac{1_{\mathbb{R}}}{\sim}$.

Proposition 4.6. Consider all of the hypotheses of Theorem 4.4. Let $R = \{(x, y) \mid x \sim y\}$. Then $(\frac{X}{\sim}, \frac{\mathfrak{T}}{\sim})$ is a Hausdorff L-gfts if and only if $\chi_{R, X \times X}$ is an LG-closed subset of $X \times X$.

Proof. Let $(\frac{X}{\sim}, \frac{\mathfrak{T}}{\sim})$ be a Hausdorff L-gfts, then for each $[x], [y] \in \frac{X}{\sim}$ where $[x] \neq [y]$, there exist $\frac{U}{\sim}, \frac{V}{\sim} \in supp\frac{\mathfrak{T}}{\sim}$ such that $[x] \in \frac{U}{\sim}, [y] \in \frac{V}{\sim}$ and $\frac{U}{\sim} \cap \frac{V}{\sim} = \phi$. So for each $(x, y) \in (X \times X)$ where $(x, y) \notin R$ there exist $U, V \in supp\mathfrak{T}$, such that $U \cap V = \phi$ and $(x, y) \in (U \times V)$. we show that $supp(U \times V) \cap R = \phi$.

$$(a, b) \in supp(U \times V) \cap R \Rightarrow a \in U, b \in V, a \sim b \Rightarrow [a] \in \frac{U}{\sim}, [b] \in \frac{V}{\sim}, [a] = [b] \Rightarrow \frac{U}{\sim} \cap \frac{V}{\sim} \neq \phi,$$

that is a contradiction. It means that for each $(x, y) \in supp(X \times X - \chi_{R, X \times X})$ there exists $U \times V \in supp(\mathfrak{T} \times \mathfrak{T})$ such that $(x, y) \in U \times V \leq (X \times X - \chi_{R, X \times X})$. Therefore, $(\mathfrak{T} \times \mathfrak{T})(X \times X - \chi_{R, X \times X}) > \mathfrak{T}(U) \wedge \mathfrak{T}(V) > 0$. Hence $\chi_{R, X \times X}$ is an LG-closed subset of $X \times X$.

Conversely, suppose that $\chi_{R, X \times X}$ is an LG -closed subset of $X \times X$. Then $(X \times X - \chi_{R, X \times X})$ is an LG -open subset. By Theorem 3.11,

$$\text{supp}(X \times X - \chi_{R, X \times X}) \in (\mathfrak{T} \times \mathfrak{T})^\triangleleft.$$

Hence $\text{supp}(X \times X) - R$ is an ordinary open subset. So for each $(x, y) \in (\text{supp}(X \times X) - R)$, there exists an open subsets $U \times V \in (\mathfrak{T} \times \mathfrak{T})^\triangleleft$ such that $(x, y) \in U \times V \subseteq (\text{supp}(X \times X) - R)$. We show that $U \cap V = \phi$

$$a \in U \cap V \Rightarrow a \sim a \text{ and } (a, a) \in U \times V \Rightarrow (a, a) \in (U \times V) \cap R,$$

that is a contradiction. It means that for each $(x, y) \in \text{supp}(X \times X)$ where $(x, y) \notin R$, there exist $U, V \in \text{supp}\mathfrak{T}$, such that $U \cap V = \phi$ and $(x, y) \in U \times V$. Since \sim is an LG -open relation, then we have $\mathfrak{T}(\frac{U}{\sim}) \geq 0$ and $\mathfrak{T}(\frac{V}{\sim}) \geq 0$. Therefore for each $[x], [y] \in \frac{X}{\sim}$ where $[x] \neq [y]$, there exist $\frac{U}{\sim}, \frac{V}{\sim} \in \text{supp}\frac{\mathfrak{T}}{\sim}$ such that $[x] \in \frac{U}{\sim}, [y] \in \frac{V}{\sim}$ and $\frac{U}{\sim} \cap \frac{V}{\sim} = \phi$. So $(\frac{X}{\sim}, \frac{\mathfrak{T}}{\sim})$ is a Hausdorff L-gfts. □

5 LG -fuzzy product manifolds

The concept of the product of fuzzy topological spaces was introduced by C. K. Wong [19] and later by Hutton [8]. We define and investigate LG -fuzzy product manifolds by the following theorem:

Theorem 5.1. *Let $X \in L^{M_1}, X_2 \in L^{M_2}$ and $(X_1, \mathfrak{T}_1), (X_2, \mathfrak{T}_2)$ be two LG -fuzzy manifolds of dimensions m, n and with the LG -fuzzy structures $\mathfrak{A}_i = \{(A_{\alpha_i}, \psi_{\alpha_i}) \mid \alpha_i \in K_i\}, i = 1, 2$ respectively. Then $(X_1 \times X_2, \mathfrak{T}_1 \times \mathfrak{T}_2)$ is an LG -fuzzy manifold of dimension $m + n$.*

Proof. We define for all $A_1 \in L_{X_1}^{M_1}, A_2 \in L_{X_2}^{M_2}$:

$$A_1 \times A_2 \in L_{X_1 \times X_2}^{M_1 \times M_2}, \quad (A_1 \times A_2)(x, y) = A_1(x) \wedge A_2(y),$$

and

$$(\mathfrak{T}_1 \times \mathfrak{T}_2)(A_1 \times A_2) = \mathfrak{T}_1(A_1) \wedge \mathfrak{T}_2(A_2).$$

It can be verified that $\mathfrak{T}_1 \times \mathfrak{T}_2$ is an L -gradation of openness on $X_1 \times X_2$. Now let $p_1 \in X_1, p_2 \in X_2$. Then there exist two LG -open subsets A_i of X_i containing p_i , for $i = 1, 2$ and two LG -open subsets B_i of LG -fuzzy Euclidean spaces of dimension m, n , respectively together with two LGP -homeomorphisms $\psi_i \in (A_i, B_i)$. Therefore for any $(p_1, p_2) \in X_1 \times X_2$, there exists an LG -open subset $A_1 \times A_2$ of $X_1 \times X_2$ containing (p_1, p_2) and an LG -open subset $B_1 \times B_2$ of LG -fuzzy Euclidean space of dimension $m + n$, together with an LGP -homeomorphism $\psi_1 \times \psi_2 \in (A_1 \times A_2, B_1 \times B_2)$ such that for each LG -open subsets $H_1 \leq B_1, H_2 \leq B_2$ we have

$$\mathfrak{T}_{L(m+n)}(H_1 \times H_2) = (\mathfrak{T}_1 \times \mathfrak{T}_2)(\psi_1^{-1}(H_1) \times \psi_2^{-1}(H_2)),$$

where $(\psi_1 \times \psi_2)(x_1, x_2) = (\psi_1(x_1), \psi_2(x_2)) \in \mathbb{R}^{m+n}$. One can prove easily that

$$\mathfrak{A}_1 \times \mathfrak{A}_2 = \{((A_{\alpha_1} \times A_{\alpha_2}), (\psi_{\alpha_1} \times \psi_{\alpha_2})) \mid \alpha_1 \in K_1, \alpha_2 \in K_2\},$$

is an LG -fuzzy structure on $X_1 \times X_2$. □

Example 5.2. *Let $M = \mathbb{R}^2, X = \chi_{S^1}$. One can easily prove that (X, \mathfrak{T}_{I_3}) is an LG -fuzzy manifold of dimension 1, similarly to Example 3.6. Then $(X \times X, \mathfrak{T}_{I_3} \times \mathfrak{T}_{I_3})$ is an LG -fuzzy manifold of dimensions 2 and $\text{supp}(X \times X) = S^1 \times S^1$.*

6 C^∞ LG -fuzzy mappings of LG -fuzzy manifolds

The concept of the fuzzy vector space (V, η) over a field F was defined in [18]. We extend this definition by L -fuzzification:

Definition 6.1. *An L -fuzzy vector space (V, η) or ηV over a field F is an ordinary vector space V over the field F , with a map $\eta : V \rightarrow L$ satisfying the following conditions for all $a, b \in V$ and $r \in F$.*

- 1) $\eta(a + b) \geq \min\{\eta(a), \eta(b)\}$,
- 2) $\eta(-a) = \eta(a)$,

- 3) $\eta(0) = 1$,
- 4) $\eta(ra) \geq \eta(a)$,

Definition 6.2. Let (X, \mathfrak{T}) be an LG-fuzzy manifold of dimension n , $U \in \text{supp}\mathfrak{T}$ and $V \in \text{supp}\mathfrak{T}_{L1}$. The LG-related function f from U to V , is called a C^∞ LG-fuzzy mapping, if for every $p \in U$,

$$\hat{f} = f \circ \psi^{-1} : \psi(\text{supp}(A \cap U)) \rightarrow \text{supp}V,$$

is C^∞ where (A, ψ) is an LG-local coordinate neighborhood of p .

We denote the set of all C^∞ LG-fuzzy mappings from an LG-open subset U of X , containing p to $1_{\mathbb{R}}$, by $C_L^\infty(p)$. If we define $\eta : C_L^\infty(p) \rightarrow L$, $\eta(f) = A(p)$, where (A, ψ) is an LG-coordinate neighborhood of p , then $C_L^\infty(p)$ may be considered as an L-fuzzy vector space $(C_L^\infty(p), \eta)$. Let $\psi(q) = (x_1, \dots, x_n)$, $\forall q \in \text{supp}(A \cap U)$. Then $\hat{f}(x_1, \dots, x_n) = y(q)$, and since \hat{f} is C^∞ , there exist all partial derivatives of any order of y .

Example 6.3. In Example 3.8, if we define $f : \mathcal{M}_{m \times n} \rightarrow \mathbb{R}$, $f((a_{ij})) = \det((a_{ij}))$, then using the single IG-local coordinate neighborhood $(Z, \psi|_Z)$, we have $\hat{f} = f \circ \psi$ is C^∞ . Hence $f \in C_I^\infty(p)$ for all $p \in Z$.

From now on, we suppose that M_1, M_2 are two crisp sets, $X \in L^{M_1}, Y \in L^{M_2}$ such that $(X, \mathfrak{T}), (Y, \mathfrak{R})$ are two LG-fuzzy manifolds of dimension n, m and LG-fuzzy structures $\mathfrak{A} = \{(A_i, \psi_i), i \in K\}$ and $\mathfrak{D} = \{(D_j, \varphi_j), j \in J\}$ respectively and $U \in \text{supp}\mathfrak{T}, V \in \text{supp}\mathfrak{D}$.

Definition 6.4. An LG-fuzzy function $F \in \text{LGR}f(U, V)$ is a C^∞ LG-fuzzy mapping if for every $p \in U$,

$$\hat{F} = \varphi \circ F \circ \psi^{-1} : \psi(\text{supp}(A \cap U)) \rightarrow \varphi(\text{supp}(B \cap V)),$$

is C^∞ where $(A, \psi), (B, \varphi)$ are LG-local coordinate neighborhoods of p and $F(p)$ respectively. $F \in \text{LGR}f(U, V)$ is called a LG-diffeomorphism if it is an LG-homeomorphism and F, F^{-1} are C^∞ .

More precisely, if $\psi(q) = (x_1, \dots, x_n)$, $\forall q \in \text{supp}(A \cap U)$ and $\varphi(w) = (y_1, \dots, y_m)$, $\forall w \in B$, then

$$\hat{F}(x_1, \dots, x_n) = (f_1(x_1, \dots, x_n), \dots, f_n(x_1, \dots, x_n)),$$

and each $y_i = f_i(x_1, \dots, x_n)$ is C^∞ on $\psi(A)$.

Definition 6.5. The rank of $F \in \text{LGR}f(X, Y)$ at p is equal to the rank at $x = \psi(q)$ of the Jacobian matrix:

$$\begin{pmatrix} \frac{\partial f_1}{\partial x_1} & \dots & \frac{\partial f_1}{\partial x_n} \\ \vdots & & \vdots \\ \frac{\partial f_m}{\partial x_1} & \dots & \frac{\partial f_m}{\partial x_n} \end{pmatrix}_x.$$

Example 6.6. Let $M_1 = M_2 = \mathbb{R}^2$, $L = I$ and $X : M_1 \rightarrow I, Y : M_2 \rightarrow I$ be defined by:

$$X(x_1, x_2) = \begin{cases} 1 & \|x\| = 1, \\ 0 & \|x\| \neq 1, \end{cases} \quad \text{and} \quad Y(y_1, y_2) = \begin{cases} 1 & \|y\| = 1, \\ 0 & \|y\| \neq 1. \end{cases}$$

If we define $\mathfrak{T} : I_X^{M_1} \rightarrow I$, and $\mathfrak{R} : I_Y^{M_2} \rightarrow I$, by

$$\mathfrak{T}(A) = \begin{cases} 1 & A \in \text{supp}\mathfrak{T}_{I2}, A \leq X, \\ 0 & \text{elsewhere.} \end{cases} \quad \text{and} \quad \mathfrak{R}(D) = \begin{cases} 1 & D \in \text{supp}\mathfrak{T}_{I2}, D \leq Y, \\ 0 & \text{elsewhere.} \end{cases}$$

In a similar manner to the Example 3.5, we can prove that $(X, \mathfrak{T}), (Y, \mathfrak{R})$ are IG-fuzzy manifolds. Let

$$F : M_1 \rightarrow M_2, \quad F(x_1, x_2) = (x_1 - x_2, \sqrt{2x_1x_2}).$$

We prove that $F \in \text{IGR}f(X, Y)$ is a C^∞ IG-fuzzy mapping. First we show $F|_{\text{supp}X} : \text{supp}X \rightarrow \text{supp}Y$ is well-defined and $F[X] = Y$:

$$(x_1, x_2) \in S^1 \Rightarrow x_1^2 + x_2^2 = 1 \Rightarrow (x_1 - x_2)^2 + (\sqrt{2x_1x_2})^2 = 1 \Rightarrow F(x_1, x_2) \in S^1,$$

$$F[X](y_1, y_2) = \sqrt{\{X(x_1, x_2) : (x_1, x_2) \in F^{-1}(y_1, y_2)\}} = 1 = Y(y_1, y_2).$$

Let $(A_1^+, \psi_1^+), (D_2^+, \varphi_2^+)$ be IG-local coordinate neighborhoods on X, Y respectively, then we see:

$$\varphi_2^+ \circ F \circ (\psi_1^+)^{-1}(y) = \varphi_2^+ \circ F(\sqrt{1-y^2}, y) = \varphi_2^+(\sqrt{1-y^2} - y, \sqrt{2y\sqrt{1-y^2}}) = \sqrt{1-y^2} - y,$$

is C^∞ . Similarly, one can show that $\varphi_i^\pm \circ F \circ (\psi_j^\pm)^{-1}$ is C^∞ for all $i, j = 1, 2$.

Archive of SID

Example 6.7. Let $F : \mathbb{R} \rightarrow \mathbb{R}^2$, $F(t) = (\cos(t - \frac{\pi}{2}), \sin(t - \frac{\pi}{2}))$. Then $F \in IGPRf(1_{\mathbb{R}}, 1_{\mathbb{R}^2})$ and $\text{rank } F = 1$ at every point of X .

Theorem 6.8. (*LG-fuzzy rank theorem*) Let $F \in LGRf(U, Y)$ be a C^∞ fuzzy mapping and $\text{rank } F = k$ at every point of X . If $p \in X$, then there exist LG-local coordinate neighborhoods (A, ψ) , (B, φ) such that

$$\psi(p) = (0, \dots, 0) \in \mathbb{R}^n, \quad \varphi(F(p)) = (0, \dots, 0) \in \mathbb{R}^m,$$

and $\hat{F} = \varphi \circ F \circ \psi^{-1}$ is given by:

$$\hat{F}(x_1, \dots, x_n) = (x_1, \dots, x_k, 0, \dots, 0). \quad (4)$$

Proof. Using Theorem 3.11, we see that $(\text{supp}X, \mathfrak{A}^\triangleleft)$ and $(\text{supp}Y, \mathfrak{R}^\triangleleft)$ are two topological manifolds of dimension n, m with the structures $\mathfrak{A}^\triangleleft = \{(\text{supp}A_i, \psi_i|_{\text{supp}A_i}), i \in K\}$ and $\mathfrak{D}^\triangleleft = \{(\text{supp}D_j, \varphi_j|_{\text{supp}D_j}), j \in J\}$ respectively. Also $F|_{\text{supp}X} : \text{supp}X \rightarrow \text{supp}Y$ is a C^∞ mapping and $\text{rank } F = k$ at every point of X . Fix $p \in \text{supp}X$, then by the rank theorem, there exist coordinate neighborhoods $(\text{supp}A, \psi|_{\text{supp}A})$, $(\text{supp}D, \varphi|_{\text{supp}D})$ of p and $F(p)$ respectively such that

$$\psi|_{\text{supp}A}(p) = (0, \dots, 0) \in \mathbb{R}^n, \quad \varphi|_{\text{supp}D}(F|_{\text{supp}X}(p)) = (0, \dots, 0) \in \mathbb{R}^m,$$

and $\hat{F}|_{\psi(\text{supp}A)} = \varphi|_{\text{supp}D} \circ F|_{\text{supp}A} \circ \psi^{-1}|_{\psi(\text{supp}A)}$ is given by:

$$\hat{F}|_{\psi(\text{supp}A)}(x_1, \dots, x_n) = (x_1, \dots, x_k, 0, \dots, 0).$$

Therefore the LG-fuzzy rank theorem holds for LG-fuzzy manifolds. \square

Remark 6.9. We can cover X and $\tilde{X} = F[X]$ by these LG-local coordinate neighborhoods $\mathfrak{A} = \{(A_s, \psi_s) \mid s \in S\}$, and $\mathfrak{D} = \{(D_s, \varphi_s) \mid s \in S\}$ respectively where $S \subseteq K$. Since \mathfrak{A} is an LG-structure of X , one can show that \mathfrak{D} is an LG-structure of $F[X]$. If F is an LG-diffeomorphism, then we have $\text{rank } F = \dim X = \dim Y$.

Definition 6.10. The C^∞ L-related function $F \in LGRf(X, Y)$ is an LG-fuzzy immersion (submersion) if $\text{rank } F = \dim X (= \dim Y)$. at every point of X .

Theorem 6.11. Let $F \in LGRf(X, Y)$ be a C^∞ LG-fuzzy mapping. If F is an injective LG-fuzzy immersion, then $(\tilde{X}, \tilde{\mathfrak{A}})$ is an LG-fuzzy submanifold of dimension n , called an LG-fuzzy immersed submanifold and $F \in LGRf(X, \tilde{X})$ is an LG-diffeomorphism.

Proof. F establishes a one-to-one correspondence between $\text{supp}X$ and $F(\text{supp}X)$. Thus, $F \in LGRf(X, \tilde{X})$ is one-to-one and onto. Since for each $q \in F(\text{supp}X)$, there exists only one $p \in \text{supp}X$ such that $F^{-1}(q) = \{p\}$, hence

$$\tilde{X}(q) = F[X](q) = \text{sup}\{X(a) \mid F(a) = q\} = X(p).$$

Since $F \in LGRf(X, Y)$, we have $F[X] \leq Y$; therefore $\tilde{X} \leq Y$. Hence \tilde{X} is an LG-fuzzy subset of Y . We use F to endow \tilde{X} with an LG-structure

$$\tilde{\mathfrak{D}} = \{(D_s|_{\tilde{X}}, \pi \circ \varphi_s|_{\tilde{X}}) \mid (D_s, \varphi_s) \in \mathfrak{D}, \forall s \in S\},$$

where $\pi(y_1, \dots, y_m) = (y_1, \dots, y_n)$ is the projection and an LG-fuzzy topology

$$\tilde{\mathfrak{A}} : L_{\tilde{X}}^{M_2} \rightarrow L, \quad \tilde{\mathfrak{A}}(H) = \mathfrak{A}(F^{-1}[H]).$$

Then $(\tilde{X}, \tilde{\mathfrak{A}})$ is an LG-fuzzy manifold of dimension n , called an LG-fuzzy immersed submanifold and $F \in LGRf(X, \tilde{X})$ is an L-gradation-preserving. Therefore F is an LGP-diffeomorphism. \square

In general, gradation of openness $\tilde{\mathfrak{A}}$ and the LG-fuzzy structure on \tilde{X} depend on F as well as X , i.e. $(\tilde{X}, \tilde{\mathfrak{A}})$ is not a submanifold of (Y, \mathfrak{R}) . So we add the condition of LG-continuity of F , F^{-1} in the following definition:

Definition 6.12. An LG-fuzzy imbedding is a one-to-one LG-fuzzy immersion $F \in LGRf(X, Y)$ with $F \in LGRf(X, \tilde{X})$ is an LG-homeomorphism from X to $\tilde{X} = F[X]$ as an LG-fuzzy subspace of Y . The image of an LG-fuzzy imbedding is called an LG-fuzzy imbedded submanifold.

Theorem 6.13. Let $F \in LGRf(X, Y)$ be an LG-fuzzy immersion. Then for each $p \in X$, exists an LG-neighborhood A of p such that $F|_{\text{supp}A}$ is an LG-fuzzy imbedding.

Proof. According to Theorem 6.8, we may choose (A, ψ) and (D, φ) , the LG -local coordinate neighborhoods of p and $F(p)$, respectively, such that (4) holds. Since $F[A] = D$ and D is an LG -open subset of Y , hence L -gradation of openness $\tilde{\mathfrak{T}}$ of $F[A]$, is the same as its L -gradation of openness $\mathfrak{D}|_D$ as an L -gftss of Y , i.e. $\tilde{\mathfrak{T}}(H) = \mathfrak{T}(F^{-1}[H]) = \mathfrak{D}(H)$, for all LG -open subset H of D . On the other hand, ψ and φ are LGP -homeomorphisms, hence \hat{F} is an LGP -homeomorphism of $\psi[A]$ and $\varphi[D]$. Therefore $F|_{suppA}$ is a homeomorphism, and thus the theorem holds. \square

Example 6.14. Let $Z = \chi_{(1,+\infty)}$. Then Z is an LG -open subset of $1_{\mathbb{R}}$. If $\mathfrak{T}_Z = \mathfrak{T}_{I_1}|_z$, then (Z, \mathfrak{T}_Z) is an LG -fuzzy submanifold of $(1_{\mathbb{R}}, \mathfrak{T}_{I_1})$. Let $W = B((0,0), 1, 1)$, then W is an LG -open subset of $1_{\mathbb{R}^2}$. Consider $F : \mathbb{R} \rightarrow \mathbb{R}^2$, $F(t) = (\frac{1}{t} \cos 2\pi t, \frac{1}{t} \sin 2\pi t)$. Then $F \in IGRf(Z, W)$ and $\text{rank } F = 1$ at every point of Z . We see $F^{-1}(x, y) = \frac{1}{\sqrt{x^2 + y^2}}$, so F is a one-to-one LG -fuzzy immersion. Since $F \in IGRf(Z, \tilde{Z})$ is an LG -homeomorphism, F is an LG -fuzzy imbedding.

7 LG -fuzzy submanifolds of LG -fuzzy manifolds

Definition 7.1. An LG -fuzzy subset N of an LG -fuzzy manifold (X, \mathfrak{T}) , is said to have the LG -fuzzy k -submanifold property if each $p \in N$ has an LG -local coordinate neighborhood (A, ψ) on X with LG -local coordinates x_1, x_2, \dots, x_n such that $\psi(p) = (0, \dots, 0) \in \mathbb{R}^n$, and

$$\psi(suppA \cap suppN) = \{(x_1, x_2, \dots, x_n) \in \psi(A) \mid x_{k+1} = \dots = x_n = 0\}.$$

If N has this property, LG -coordinate neighborhoods of this type are called preferred LG -local coordinates.

Denote by $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^k$, $k \leq n$, the projection to the first k coordinates. Using the notation above, we may state the following proposition:

Theorem 7.2. Let $N \leq X$ have the LG -fuzzy k -submanifold property. Then each preferred LG -local coordinate system (A, ψ) of X defines an LG -local coordinate neighborhood (A', ψ') on N where $A' = A \cap N$, $\psi' = \pi \circ \psi|_{A'}$. Therefore the inclusion $i \in LGRf(N, X)$ is an LG -fuzzy imbedding.

Proof. Since N is an LG -open subset of X , thus (N, \mathfrak{T}_N) is an LG -fuzzy topological subspace of X . Then (A', ψ') are LG -coordinate neighborhoods covering N , where $A' = A \cap N$ is an LG -open subset of N and $\psi' = \pi \circ \psi|_{A'}$ is an LGP -homeomorphism. Suppose that for two preferred neighborhoods (A'_1, ψ'_1) and (A'_2, ψ'_2) , A'_1, A'_2 have a nonempty intersection. We know that the change of LG -local coordinates is given by LGP -homeomorphisms $\psi'_1 \circ \psi'^{-1}_2$ and $\psi'_2 \circ \psi'^{-1}_1$ which we must show to be C^∞ . Let

$$\gamma(x_1, \dots, x_k) = (x_1, \dots, x_k, 0, \dots, 0) \in \mathbb{R}^n,$$

so that $\pi \circ \gamma$ is the identity on \mathbb{R}^k . This map γ is C^∞ . Hence its restriction to $\psi'(A')$, an LG -open subset of \mathbb{R}^k , is C^∞ ; thus $\psi'^{-1} = \psi \circ \gamma$ is C^∞ , since it is a composition of C^∞ maps. On the other hand, $\psi' = \pi \circ \psi$ so ψ' is a C^∞ map on A' . Hence $\psi'_1 \circ \psi'^{-1}_2$ is C^∞ . If $y_i = f_i(x_1, \dots, x_k)$, $i = 1, \dots, k$, are the functions giving $\psi'_1 \circ \psi'^{-1}_2$, which we know to be C^∞ , then it can easily be checked that $\psi_1 \circ \psi_2^{-1}$ is given by $y_i = f_i(x_1, \dots, x_k, 0, \dots, 0)$, $i = 1, \dots, k$. Therefore $\psi'_1 \circ \psi'^{-1}_2$ is C^∞ by Definition 3.2. Thus the totality of these LG -neighborhoods define a unique differentiable structure on N . In preferred LG -local coordinates (A', ψ') , $i \in LGRf(N, X)$ is given on V by

$$\psi \circ i \circ \psi'^{-1}(x_1, \dots, x_k) = (x_1, \dots, x_k, 0, \dots, 0).$$

So the map i is clearly an LG -fuzzy immersion. Because we have taken the relative LG -fuzzy topology on N , the fuzzy map i is by Definition 2.13 (iii) an LG -homeomorphism to its image $i(N)$, with the LG -fuzzy subspace topology, that is, i is an LG -fuzzy imbedding. \square

Definition 7.3. A regular LG -fuzzy submanifold of an LG -fuzzy manifold (X, \mathfrak{T}) is any LG -fuzzy topological subspace N with the LG -fuzzy submanifold property and with the structure that the corresponding preferred LG -local coordinate neighborhoods determine on it.

Example 7.4. Let $M = \mathbb{R}^3$, $X : \mathbb{R}^3 \rightarrow I$, $X(x) = \begin{cases} 1 & \|x\| = 1, \\ 0 & \|x\| \neq 1. \end{cases}$. Then $suppX = S^2$, the unit sphere. Let

$\mathfrak{T} : I_X^M \rightarrow I$, $\mathfrak{T}(A) = \begin{cases} 1 & A \in \tau_{I_3}, A \leq X \\ 0 & \text{elsewhere.} \end{cases}$. We shall see that X is an LG -fuzzy submanifold of $(1_{\mathbb{R}^3}, \mathfrak{T}_{I_3})$. If

$q = (x_1, x_2, x_3)$ is an arbitrary point in $\text{supp}X$, it cannot lie on more than one coordinate axis. For convenience, we assume that it does not lie on the x_3 -axis. We introduce the spherical LG-local coordinates (ρ, θ, φ) ; they are defined on $1_{\mathbb{R}^3 - \{x_3\text{-axis}\}}$ and if $(1, \theta_0, \varphi_0)$ are the LG-coordinates of q , we may change them a little so that it is replaced by $\tilde{\rho} = \rho - 1$, $\tilde{\theta} = \theta - \theta_0$, and $\tilde{\varphi} = \varphi - \varphi_0$. Then it defines an LG-coordinate neighborhood of q , with q having LG-coordinates $(0, 0, 0)$ and with the LG-open subset V of X .

Remark 7.5. So far, we have defined three classes of LG-fuzzy submanifolds of an LG-fuzzy n -manifold (X, \mathfrak{T}) . The first of these, which we usually simply call an LG-fuzzy submanifold, was defined (in 6.11) as the image $N = F[N']$ of an LG-fuzzy immersion F of N' into X . Since $F : N' \rightarrow N \leq X$ is one-to-one and onto, we conduct (as part of the definition) carry over to N the LG-fuzzy topology and LG-fuzzy structure of N' ; LG-open subsets of N are the images of LG-open sets of N' and LG-coordinate neighborhoods (A, ψ) of N are of the form $A = F[A']$, $\psi = \psi' \circ F^{-1}$, where (A', ψ') is an LG-local coordinate neighborhood of N' . The fact that F is LG-continuous shows that the LG-fuzzy topology of N gained in this way is in general finer than its relative LG-fuzzy topology as an LG-fuzzy subspace of X , that is, if D is LG-open subset of X , then $D \cap N$ is LG-open subset of N , but there may be LG-open subsets of N which are not of this form.

An LG-fuzzy imbedding is a particular type of LG-fuzzy immersion, one in which A is LG-open subset of N if and only if $A = F[U'] = D \cap N$ for some LG-open subset D of X so that the LG-fuzzy topology of the submanifold $N = F[N']$ is exactly its relative LG-fuzzy topology as an LG-fuzzy topological subspace of X . An LG-fuzzy imbedded submanifold is so a special type of (immersed) LG-fuzzy submanifold.

Ultimately, if $N \leq X$ is an LG-fuzzy regular submanifold, then it is also an LG-fuzzy imbedded submanifold since the inclusion $i : N \rightarrow X$ is an LG-fuzzy imbedding as we proved in 7.2.

Theorem 7.6. Let $F \in \text{LGRf}(N', X)$ be an LG-fuzzy imbedding of an LG-fuzzy manifold N' of dimension k in an LG-fuzzy manifold of dimension n . Then $N = F[N']$ has the LG-fuzzy k -submanifold property and thus N is an LG-fuzzy regular submanifold. As such, it is LG-diffeomorphic to N' with respect to the LG-fuzzy mapping $F \in \text{LGRf}(N', N)$.

Proof. Let $q = F(p)$ be any point of N . According to Theorem 7.2 (and its proof), there are (A, ψ) and (B, φ) , LG-local coordinate neighborhoods of p and $F(p)$, respectively, such that (4) holds. If $F[A] = V \leq N$, then the LG-neighborhood V would be a preferred LG-local coordinate neighborhood relative to N . To deduce this result, we should use the fact that F is an LG-imbedding. This denotes at least that $F[A]$ is a relatively LG-open subset of N , that is, $F[A] = W \cap N$, where W is LG-open subset of X . Since $F[A] \leq V$, we can suppose $W \leq V$. Thus $\varphi[W]$ is an LG-open subset of $\varphi[B]$ containing the origin in \mathbb{R}^n and $\varphi[F[A]] \leq \varphi[W]$, which is a slice S of $\varphi[V]$, $S = \{x \in \varphi[V] \mid x_{k+l} = \dots = x_m = 0\}$. Hence we may select an (smaller) LG-open subset $\varphi[V'] \leq \varphi[W]$ and $\varphi' = \varphi|_{\text{supp}V'}$. This is an LG-local coordinate neighborhood of q for which $F[A] \cap V' = V' \cap N$; furthermore, taking $A' = F^{-1}[V']$, we see that (A', ψ') , with $\psi' = \psi|_{\text{supp}A'}$, is an LG-local coordinate neighborhood of p and the pair (A', ψ') and (V', φ') have exactly the properties needed in 7.1 and $F[A'] = V' \leq N$. This proves at the same time, that N has the LG-fuzzy k -submanifold property. This is true since the inverse of $F \in \text{LGRf}(N', N)$ is given in the preferred LG-local coordinates $(V', \pi \circ \varphi')$ and (A', ψ') by $\hat{F}^{-1}(x_1, \dots, x_k) = (x_1, \dots, x_k)$, which is C^∞ . \square

Remark 7.7. Suppose that $N \leq X$ is an LG-fuzzy immersed submanifold and that $q \in N$. Then there is an LG-neighborhood (V, ψ) of q , with $\psi(p) = (0, \dots, 0)$ such that the slice $S' \subseteq \text{supp}V$, consisting of all points of V whose last $n - k$ coordinates vanish, is an LG-open set and an LG-local coordinate neighborhood of the LG-fuzzy submanifold structure of N is given by LG-local coordinate map

$$\psi'(q) = \pi \circ \psi(q) = (x_1(q), \dots, x_k(q)).$$

Theorem 7.8. If $F \in \text{LGRf}(N, X)$ is a one-to-one LG-fuzzy immersion and N is a compact L-gfts, then F is an LG-fuzzy imbedding and $\tilde{N} = F[N]$ an LG-fuzzy regular submanifold.

Proof. Since F is LG-continuous and both N and \tilde{N} are Hausdorff L-gfts's, we have an LG-continuous (one-to-one) mapping from a compact L-gfts to a Hausdorff L-gfts. Since an LG-closed subset K of N is compact, so $F(K)$ is compact and therefore LG-closed. Thus F takes LG-closed subsets of N to LG-closed subsets of X , and since F is one-to-one and onto, it takes LG-open subsets to LG-open subsets as well. It follows that F^{-1} is LG-continuous, so $F \in \text{LGRf}(N, \tilde{N})$ is an LG-homeomorphism and therefore an LG-imbedding. \square

Theorem 7.9. Let $F \in \text{LGRf}(X, Y)$ be a C^∞ LG-fuzzy mapping. Suppose that F has constant rank k on X and that $q \in F(X)$. Let D denotes $F^{-1}(q)$; then χ_D is an LG-closed, LG-fuzzy regular submanifold of X of dimension $n - k$.

Proof. Let $p \in D$; since F has constant rank k on an LG -neighborhood of p , we may find LG -local coordinate neighborhoods (A, ψ) , (B, φ) such that (4) holds. By Example 2.9, the fuzzy point 0_1 is an IG -closed subset of \mathbb{R}^n , then $\chi_{\{q\}}$, is an LG -closed subset of Y . Hence χ_D is an LG -closed subset since the inverse image of $\chi_{\{q\}}$, under a continuous map, is LG -closed. We shall show that χ_D has the LG -fuzzy $n - k$ submanifold property. This means that the only points of D mapped onto q are those whose first k coordinates are zero, that is,

$$\text{supp}A \cap D = \psi^{-1}(\psi \circ F^{-1} \circ \varphi^{-1}(0)) = \psi^{-1}(\hat{F}^{-1}(0)) = \psi^{-1}\{x \in \psi(A) \mid x_1 = \dots = x_k = 0\}.$$

Hence χ_D is a regular LG -fuzzy $(n - k)$ -submanifold since it has the LG -fuzzy submanifold property. □

Corollary 7.10. *If $F \in LGRf(X, Y)$ is a C^∞ LG -fuzzy mapping of LG -fuzzy manifolds, $\dim X = n \leq m = \dim Y$, and $\text{rank } F = n$ at every point of $D = F^{-1}(q)$, then χ_D is an LG -closed, regular LG -fuzzy submanifold of X . The corollary holds because at $p \in A$, F has the maximum rank possible, namely m . It follows from the independence of rank on LG -local coordinates that, in some LG -neighborhood of p in N , F also has this rank; thus the rank of F is m on an LG -open subset of N containing A . But such an LG -fuzzy subset is itself an LG -fuzzy n -manifold (an LG -open submanifold) to which we may apply the theorem.*

8 Conclusion

In this paper, we generalize all of the fuzzy structures which we have discussed in [14] to L -fuzzy set theory, where $L = \langle L, \leq, \wedge, \vee, ' \rangle$ denotes a complete distributive lattice with at least two elements. We define the concept of an LG -fuzzy topological space (X, \mathfrak{T}) which X is itself an L -fuzzy subset of a crisp set M and \mathfrak{T} is an L -gradation of openness of L -fuzzy subsets of M which are less than or equal to X . Then we define C^∞ L -fuzzy manifolds with L -gradation of openness and C^∞ LG -fuzzy mappings of them such as LG -fuzzy immersions and LG -fuzzy imbeddings. We fuzzify the concept of the product manifolds with L -gradation of openness and define LG -fuzzy quotient manifolds when we have an equivalence relation on M and investigate the conditions of the existence of the quotient manifolds. We also introduce LG -fuzzy immersed, imbedded and regular submanifolds.

References

- [1] E. L. Chang, *Fuzzy topological spaces*, Journal of Mathematical Analysis and Applications, **24** (1968), 182-190.
- [2] K. C. Chattopadhyay, R. N. Hazra, S. K. Samanta, *Gradation of openness: Fuzzy topology*, Fuzzy Sets and Systems, **49**(2) (1992), 237-242.
- [3] K. C. Chattopadhyay, S. K. Samanta, *Fuzzy topology: Fuzzy closure operator, fuzzy compactness and fuzzy connectedness*, Fuzzy Sets and Systems, **54**(2) (1993), 207-212.
- [4] M. K. EL-Gayyar, E. E. Kerre, A. A. Ramadan, *On smooth topological spaces II: Separation axioms*, Fuzzy Sets and Systems, **119**(3) (2001), 495-504.
- [5] M. Ferraro, D. H. Foster, *C^1 fuzzy manifolds*, Fuzzy Sets and Systems, **54**(1) (1993), 99-106.
- [6] J. A. Goguen, *L -fuzzy sets*, Journal of Mathematical Analysis and Applications, **18** (1967), 145-174.
- [7] V. Gregori, A. Vidal, *Fuzziness in Chang's fuzzy topological spaces*, Journal Rendiconti dell'Istituto di Matematica dell'Universit di Trieste, **XXX** (1999), 111-121.
- [8] B. Hutton, *Products of fuzzy topological spaces*, Topology and its Application, **11** (1980), 59-67.
- [9] I. Ibedou, *Graded fuzzy topological spaces*, Cogent Mathematics and Statistics, **3**(1) (2016), 1-13.
- [10] R. Lowen, *Fuzzy topological spaces and fuzzy compactness*, Journal of Mathematical Analysis and Applications, **56** (1979), 621-633.
- [11] R. Lowen, *Mathematics and fuzziness, some personal reflections*, Information Sciences, **36**(1-2) (1985), 17-27.
- [12] E. Lowen, R. Lowen, *On measures of compactness in fuzzy topological spaces*, Journal of Mathematical Analysis and Applications, **131**(2) (1988), 329-340.

- [13] M. Macho Stadler, M. A. de Prada Vicente, *On fuzzy subspaces*, Fuzzy Sets and Systems, **58** (1993), 365-373.
- [14] M. Mostafavi, *An approach to some properties of fuzzy topological spaces*, Cankaya University Journal of Science and Engineering, **14**(1) (2017), 1-9.
- [15] A. P. Šostak, *On a fuzzy topological structure*, Rendiconti del Circolo Matematico di Palermo Serie II, **11** (1985), 89-103.
- [16] A. P. Šostak, *On compactness and connectedness degrees of fuzzy sets in fuzzy topological spaces*, General Topology and its Relations to Modern Analysis and Algebra, VI, **16** (1988), 519-532.
- [17] A. P. Šostak, D. Dzhajanbajev, *On completeness and precompactness spectra of fuzzy sets in fuzzy uniform spaces*, Commentationes Mathematicae Universitatis Carolinae, **31**(1) (1990), 149-157.
- [18] W. B. Vasantha Kandasamy, F. Smarandache, I. Kandasamy, *Set linear algebra and set fuzzy linear algebra*, Infinite Study, (2008), 344 pages.
- [19] C. K. Wong, *Fuzzy topology, product and quotient theorems*, Journal of Mathematical Analysis and Applications, **45**(2) (1974), 512-521.
- [20] P. Wuyts, R. Lowen, *On local and global measures of separation in fuzzy topological spaces*, Fuzzy Sets and Systems, **19**(1) (1986), 51-80.
- [21] L. A. Zadeh, *Fuzzy sets*, Information and Control, **8** (1965), 338-353.

C^∞ L -fuzzy manifolds with L -gradation of openness and C^∞ LG -fuzzy mappings of them

M. Mostafavi

خمینه‌های L -فازی C^∞ با L -درجه‌بندی بازبودن و نگاشت‌های L -فازی C^∞ آن‌ها

چکیده. در این مقاله، ما تمام ساختارهای فازی که در [۱۴] مورد بحث قرار داده‌ایم را با نظریه مجموعه‌های L -فازی تعمیم می‌دهیم که $L = \langle L, \leq, \wedge, \vee, ' \rangle$ نشانگر یک شبکه توزیع‌پذیر با حداقل با دو عنصر می‌باشد. ما مفهوم فضای توپولوژی LG -فازی (X, τ) ، را معرفی می‌کنیم که X خود یک زیرمجموعه L -فازی از مجموعه مرجع M است و τ یک L -درجه‌بندی بازبودن از زیر مجموعه‌های L -فازی از M است که کمتر یا مساوی M می‌باشند. سپس ما خمینه‌های L -فازی C^∞ با L -درجه‌بندی بازبودن و نگاشت‌های L -فازی C^∞ آن‌ها را تعریف می‌کنیم مانند غوطه-ورسازی‌های L -فازی و نشانده‌های L -فازی. در ادامه ما مفهوم خمینه‌های حاصل ضربی را با L -درجه‌بندی بازبودن فازی می‌کنیم و مفهوم خمینه‌های خارج قسمتی LG -فازی را هنگامی که ما یک رابطه هم ارزی روی M داریم، ارائه می‌نماییم و شرایط وجود چنین خمینه‌های خارج قسمتی LG -فازی را بررسی می‌کنیم. همچنین ما زیرخمینه‌های غوطه‌ور شده و نشانده شده و معمولی LG -فازی را معرفی می‌کنیم.