Effect of two methods of training, Small Sided Game (SSG) and resistanceplyometric, on plasma level of Interleukin-6 and tumor necrosis factor-alpha in youth soccer players

S. Rostami, MSc¹ M. Faramarzi, PhD² G. Arbab, PhD Student³ L. Ahmadi, MSc⁴

Instructor Department of Physical Education¹, Islamic Azad University, Mobarake Branch, Isfahan, Iran. Associate Professor Department of Physical Education², Shahrekord University, Shahekord, Iran. Instructor Department of Physical Education³, PhD Student of Physical Education⁴, Islamic Azad University, Bandar Abbas Branch, Bandar Abbas, Iran.

(Received 19 May, 2013 Accepted 27 Aug, 2013)

ABSTRACT

Introduction: The aim of this study was to examine the effect of two concurrent resistance-plyometric and small sided game methods on pro-inflammatory in youth soccer players.

Methods: 30 youth soccer players with a range of 14 to 17 years of age were divided randomly in three groups, SSG (n=8), resistance-plyometric (n=11) and control (n=11). Height, weight, BMI and VO2max of the subjects were measured. The next day, blood samples from the anterior brachial vein of subjects were taken. The experimental groups participated in their exercise program (8 week, 2 sessions per week plus three current sessions). All variables were measured again after 8 weeks. In order to compare data within groups' Dependent t-test was used and for comparison between groups Independent t-test was used by SPSS 17.

Results: The findings showed that after 8 weeks of SSG training plasma level of IL-6 and TNF- α increased significantly. The results in other groups did not show significant change (P ≥ 0.05).

Conclusion: SSG training can induce a catabolic environment in youth soccer players' body.

Key words: Interluecin-6 - Cytokine - Soccer Player

Introduction:

Soccer special exercises usually include exercise games done in a smaller field with a few number of people. This special type of exercise is called small-sided game (SSG) (13). The SSG exercises is often as a part of exercise programs for adult players in various forms depending on exercise goal (8). The interval exercises by using SSG exercise is effective for improving aerobic preparation and soccer special resistence. Nemeth et.al (2007) surveyed the effectiveness of one hour and a half wrestling exercise on inflammatory cytokines and growth factors in highschool adolescents (aged 14 to 18), and the result showed that the exercise stages for one hour and a half (12 weeks) with intensity leads to decrease anabolic insulin-like mediators, 1GF-1, GH and to increase inflammatory cytokines (16). Also, Anderson et al (2009) in their research, analyzed cytokine responses after two soccer matches in 72 hours. The blood specimen was taken from elite female players between two

Correspondence: S. Rostami, MSc. Department of Management and Accounting Islamic Azad University, Mobarkeh Branch, Isfahan, Iran Tel: +98 913 1088601 Email: srostami.20@gmail.com halves of the first match. 20min after the first match, 69h after the first match and 20min after the second match. The results showed that interleukin-6 amounts, necrotuzing, α tumor, interferons, leukocytes, interleukin-12 increased. The return way to the former position had no impact on cytokine secretion (19). On the other side, soccer is a sport based on explosive movements like kicking, jumping and fast replacing. The players traversed about 10Km and needed frequent and fast replacements in regular time intervals during the match. So, one of the most important parts in players exercise programs should be improving the soccer special power defined as a player ability for using the muscle power and ability effectively and continuously in soccer special activity during the match (14).

Rian et al (2004), also studied effectiveness of training 6 week combined (resistenceplyometric) of physical function of 27 boys aged 12-15, and showed that this combined training method has a considerable effect on physical performance of youth-athletes (11). Rubley et.al (2009) surveyed the vertical jump and kick powers in youngster female players. 16 players perfomed plyometric exercises for 14 weeks, and the above tests were given before and after exercises. The results showed that plyometric exercises lead to increase kick and jump power (19). So regarding to developmentimprovement of physical performance by each one of resistence and plyometric exercises by itself and investigation. according to recent the combination of two exercises leads to more development of physical performance (7).

Therefore, nowadays in exercise programs of most soccer players specially young players, there are both soccer special resistance exercises like SSG strength-plyometric together. Although, various studies have been done about these exercises effects on physical functions, a few studies have been done about hormonal and biochemical effects of these exercises in youth separately or compoundly. Cytokines are one of the biochemical factors as well. They are an important intercellular massage-transfer molecules which regulate inflammatory reaction. The number of leukocytes and IL-6 level increase and α tumor necrosing factors α (TNF α) has been shown in male soccer players in the previous research (13). Also, it was shown that the preinflammatory response can be balanced through anti-inflammatory cytokines production after resistence exercises (10).

Today, the complication of anabolic, catabolic response to exercises is unclear. So, in this study, we focused on new mediatory factors studied previously separately which can intervene in this process. It means whether a period of mixed exercises (resistence-plyometric) and SSG in young soccer players can lead to change of resting level of inflammatory factors or not?

Methods:

In this study based on contrastive analysis, the effectiveness of 2 kinds of special soccer training and resistance-plyometric, or modifications of some pre-inflammatory in young boy players, was compared. In this way that two exercise protocols after 8 week training on IL-6 and TNF- α changes in young players were compared.

The related data for measuring the research dependant variables was collected by pre-test and post-test and the effectiveness of independent variable on dependant variable was surveyed in experimental examinees. The studied group including 30 young players with continuous participation in Giti Pasand soccer team were selected for two protocols, soccer specialized and resistance-plyometric. In regard to the topic, the technique and method of study meaning the existence of two experimental and control group, pre-test and post-test and the presentation of independent variable from experimental group, this research is a quasi-experimental study done in the field, and is also applicable due to using the acquired results.

The research examinees included 50 individuals of young soccer players in Giti Pasand club checking by the coach of the mentioned team, 30 subjects participating constantly in exercises (at least for past 1 year) with full physical health and without chronic disease or injury, were selected by, purposeful sampling technique. Then the subjects Randomized in 3 experimental, 1 (the Soccer Specialized exercise group including 8 people), 2 (the resistance-plyometric exercise group including 8 people) and control group (11 people). The selection method was in a way that at first the questionnaires containing medical history of the selected players by initial checking with the club coaches and managers were distributed among the players.

In this questionnaire, there was emphasis on some entrance conditions in the study including lack of disease history and following an appropriate diet, lack of smoking, alcohol, caffeine and drug.

Then, we selected 30 qualified subjects who had participated continuously in the exercises in order to do two exercise protocols. It is worth mentioning that the players were distributed randomly in both groups and all examinees pronounced that they have had full physical health and haven't had any diseases and infection history. After the necessary checking with authorities and written consent from the parents and players, the planning began to do the investigation. In table1, the central indexes related dispersion to general characterization of examinees including age, height, weight, body mass index and Vo2max have been presented.

The examinees were asked to attend the club in order to measure the primary anthropometric indexes such as height, weight, body structure.

The height based on cm and using tape measure and the weight based on Kg and using Seca balance and when they had the fewest clothes on without shoes, were measured.

Also, body mass index using an applicant for measuring body structure was measured in the club office.

After measuring these indexes, running test for 12 min was done to measure the maximum consumed oxygen. For measuring the maximum consumed oxygen of the examinees, walking and running test for 12 minutes was used, in this way that the examinees ran around the field for 12 minutes and the traversed distance by every examinee was calculated and then using the following formula, their Vo₂max were calculated.

 $Vo_2(max) = \left(\frac{mili\ Litire}{Kg\ \times\ min}\right) = [D\ \times\ 0/0268] - 11/3$

Which in this formula, D for traversed distance was in meter. (9)

The exercise program was done during 8 weeks divided into two stages in 4 week to adjust the work intensity. The first stage included SSG exercise for 4 weeks with 2 sets for 11 minutes and 120 seconds break between the sets which was performed in 4×4 method in a 40×20 m field in the second 4 weeks, SSG protocol included 3 sets for 7 minutes and 60 seconds break which is performed in 2×2 method and a 20×15 m field.

This program was used previously by Hill Has et al (2009) and its effective properties was proved (5). These exercises were done twice a week along with current soccer exercises.

The second exercise protocol (plyometricresistance exercise) accomplished by 11 players included mined exercise for 8 weeks (4).

The examinees performed the plyometricresistance exercise along with the current exercise. The intensity of exercise was slow in the first two weeks and medium in next six weeks. The resistance exercises includinganterior of the pesterort of the tigh, chest press food press and tigh movements and plyometric exercises were the movement with box and medicine ball. The rest of examinees (11 people) as control group will do ordinary soccer exercises under the supervision of their coaches during 8 weeks.

For analyzing the biochemical variables, the blood taking was done after 12-14 hours of fasting and in two stages (before exercises and after 8 weeks of the exercise). In the first stage, the examinees were asked to do on exercise in two days before the test. Then, they attended the medical diagnosis laboratory. The temperature and time of the test were recorded to maintain the conditions in the next stage. 7 mL clogged blood was taken from the vein of left hand in sitting and rest states and immediately the serums were separated. The serum obtained from the lab was kept in temperature -80 °C to be used in case of necessity. After this stage, the examinees were under specialized exercise for 8 weeks and they were invited again to the laboratory to give blood like the first stage after passing the desirable time and 24 hours after the last session of exercise. 1L-6 and TNF- α were tested. Using method ELISA and respectively company kits, Bender med systems and bioscience.

To analyze the data, descriptive statistics. To evaluate the changes in every index from pre-test to post-test and to compare the result between the groups t-test and ANOVA analysis tests were used SPSS 17 software was used to analyze and a significant level ($P \le 0.05$) in this study was considered statistical significant.

Results:

The result of the recent study showed that interleukirn-6 (P=0.001) increased considerably in SSG group which wasn't significant in control and resistance – plyometric groups (P=0.052 and P=0.628 respectively). An exercise period of SSG in this study raised the level of necrotizing factor in tumor α (P=0.001), but this change is not significant in the resistance-plyometric exercise and control group (repectively P=0.602 and P=0.846). The data related to the comparison between the groups show that there is a significant difference between difference mean of three groups in posttest (P=0.005).

The pursuing test (LSD) is used to determine the difference between groups. There is a significant discrepancy between SSG and control group and also SSG and resistance-plyometric (interleukin-6 is P=18% SSG between mixed groups and is P=0.0001 SSG between control groups, necrotizing factor of tumor α is P=0.003 SSG between mixed groups and is P=0.013 SSG) between control groups that show a significant increase in the average amount of inteleukin-6 and necrotizing factor of α in SSG group.

 Table 1. Physical and psychological characteristics in enemaines

Statistics Group	Number	Age (year)	Height (cm)	Weghit (kg)	Body Mass Index (kg/m2)	VO2MAX (milt/kg)
SSG Group	8	14.62 ± 3.1	1.81 ± 0.26	69.5 ± 10.5	23.66 ± 2.14	39.22 ± 2.84
Ressistance – Plyometric Group	11	14.57 ± 0.9	1.71 ± 0.58	62.29 ± 15.1	21.63 ± 2.95	40.54 ± 3.04
Control Group	11	$15.67{\pm}0.6$	1.74 ± 0.42	68.19 ± 14.7	21.31 ± 2.11	40.89 ± 2.76

Table 2. Descriptive changes of data								
Statistics Variable	Group	Pretest	Posttest	Intra Group P	Inter Group P			
	SSG	1.64 ± 0.42	5.75 ± 1.94	* 0.001				
IL-6	Resistance-Plyometric	2.26 ± 1.86	3.44 ± 2.24	0.0527	* 0.005			
	Control	2.40 ± 1.28	2.55 ± 0.98	0.628				
	SSG	7.47 ± 0.85	11.68 ± 3.26	* 0.001				
TF-ą (pg/ml)	Resistance-Plyometric	1.8 ± 0.3	1.2 ± 0.3	0.602	* 0.009			
	Control	10.47 ± 3.40	1017 ± 2.78	0.846				
* Simificant Ch	29.92							

* Significant Change



Diagram 1- Changes of IL-6 and TNF during 8 Weeks of Training in Three Group

Conclusion:

The results in the recent research showed that interleukin-6 and necrosis factor of tumor α rose considerably in SSG group which was insignificant in resistance-plyometric control groups. Regular and long-term exercises remarkably change the secretion of interleukin-6 and necrosis factor of tumor α . Respecting the involvement of most lower muscles in research in method SSG, including running pattern, the kind of spasmodic muscle may have an important role to release interleukin-6 and necrosis factor of tumor α . On the other side, each regular and lognterm physical activity causing the increase of spending energy leads to increase the level in preinflammatory cytokines cycle. In general, it can be said that long-term stress raises the level of the stress hormones and leads to positive changes in muscular synthesis. Also, the significant increase in IL-6 and TNF- α may be due to increasing in intensity of this exercise method followed by the probable decrease of glycogen storage. In other words, when glycogen in type - 2 muscular fibers during long-term exercise is discharged, type-2 fibers are called respectively to preserve the energy. When these fibers are called, they are duplicated and immediately produce interleukin-6 (15).

A research by Elya Kim et al (2009) showed that the effectiveness of volleyball current exercises on male and female players in national level during 8 weeks with significant increase, is compatible with our research (1). In this research, the investigators declared jump, frequent hits and rapid replacements cause tiny injuries in tissues leading to increase inflammatory cytokines. Also in soccer, the frequent hits, jumps and rapid replacements in long-term may lead to such mechanism and be one of the reasons for the rise of interleukin-6 and TNF- α level in this research. Timoei et al (2009), analyzed the impact of endurance exercise during 5 weeks on prepubertal and early pubertal youths. Also, in this research interleukin-6 and necrosis factor of tumor α rise significantly which is associated with the recent research result (12). In this investigation, it is stated that the increase of interleukin-6, necrosis factor of tumor α can be due to increasing the energy spending by severe physical activity. In regard to much dependence of soccer exercises to muscular glycogen and the probable discharge at the end of each exercise session and relatively high spent energy of these exercises may be the probable reason of the increase of IL-6 and TNF- α in this study. The study by Ensilly et al (2008) about the resistance exercises in professional cyclists (10) and the study by Anderson et al (2009) about cytokine response after two soccer matches with 72h interval showed a significant rise in interleukin-6 which corresponds with the recent research (6).

In a study by Farzanegi et al (2010), profile changes of cytokines in male adolescent gymnasts during 8 weeks exercise after vaccination against influenza was analyzed and a decrease in interleukin-6 concentration and necrosis factor of tumor α was observed which does not correspond with our study. It can be said that regarding to similarity in exercise duration, perhaps vaccine impact or exercise intensity is the reason of difference in the result with the recent research result (8). In a research by Akbari et al (2009), interleukin-6 and necrosis factor of tumor α had a significant decrease which contradicts out result this difference may be because of perceiving supplemental L-carnitine and observing improvement of immune system function (7). In a study by Rian Kord (2004), interleukin-6 and necrosis factor of tumor α showed a significant decrease which is opposed to our result. This difference may be due to hyper-activity of the examinee during increasing age (3).

It can be concluded that in SSG due to high severe exercise and high level of stress hormones, evacuation of glycogen storage existing in muscle and tiny injuries in tissue emerging from rapid replacements and frequent hits in soccer, the inflammatory cytokines increase significantly. Our findings based on inteleukin-6 increase after 8 weeks of SSG show that this interleukin is an inflammatory cytokine sensitive to sport activity. Running and current hits in this training method with intra-tissue injuries might lead to increase interleukin-6 in long term which is one of the reasons for increasing hypertrophy. On the other side, regular and long-term physical activity which causes an increase in spending energy leads to increase the levels in pre-inflammatory cytokines cycle. Generally, it can be said that long stress can increase the level of stress hormones and lead to positive changes in muscle synthesis. Also, the significant IL-6, TNF- α may be due to high intensity of these exercises method followed by the decrease of glycogen storage. The high level of interleukin-6 can cause a decrease in the function of growth-hormone axis - the growth hormoneinsulin-like facor 1.

Numerous researches have shown that the preinflammatory cytokines have catabolic role in response to long-term exercise. Such changes were not seen in resistance-plyometric method. More analysis to study the coordination and relationship between these variables by growthbiochemical mediators in plasma can illustrate the role of these changes. In this research, the efforts focus on deciding the limitations as much as possible to draw a realistic conclusion from the findings. The controlled limitation included age, gender, time and taking blood condition and uncontrolled limitations, individual differences like genetic, family and psychological factors, lack of control for examines diet and impossibility of disease control.

Acknowledgment:

It needs to appreciate physical education group in Shahre Kord University and the players and technical personnel in Gitipasand club in Shahin Shahr who helped us in implementation of research sketch.

Hormozgan Medical Journal, Vol 18, No.4, Oct-Nov 2014 www.SID.ir

References:

- Dellal A, Chamari C, Wong DP, Ahmaidi S, Keller D, Barros MLR, et al. Comparison of physical and technical performance in European professional soccer match-play: The FA Premier League and La LIGA. *Eur J Sport Sci.* 2010;25:93-100.
- Dellal A, Keller D, Carling C, Chaouachi A, Wong DP, Chamari K. Physiological effects of directional changes in intermittent exercise in soccer players. J Strength Cond Res. 2010;32:3219-3226.
- Di Salvo V, Baron R, Tschan H, Calderon Montero FJ, Bachl N, Pigozzi F. Performance characteristics according to playing position in elite soccer. *Int J Sports Med.* 2012;28:222-227.
- 4. Duthie GM, Young WB, Warren B, Aitken DA. The acute effects of heavy loads on jump squat performance: an elevation of the complex and contrast methods of power development. *J Strength Cond Res.* 2007;16:530-538.
- 5. Eduardo EJ, Janeria MA. Effects of Complex Training on Explosive Strength in Adolescent Male Basketball Players. *J Strength Cond Res.* 2008;22:903-909.
- Fanchini M, Azzalin A, Castagna C, Schena F, McCall A, Impellizzeri FM. Effect of bout duration on exercise intensity and technical performance of small-sided games in soccer. J Strength Cond Res. 2009;25:453-458.
- Faigenbaun AD, McFarland JE, Keiper FB, Tevlin W, Ratamess NA, Kang J, et al. Effect of a short-term plyometric and resistance training program of fitness performance in boys age 12 to 15 years. *J Sports Sci Med*. 2007;6:519-525.
- Febbraio MA, Pedersen BK. Muscle-derived interleukin-6: mechanisms for activation and possible biological roles. *FASEB Journal*. 2002;16:1335-1347.
- Helgerud J, Engen LC, Wisloff U, Hoff J. Aerobic endurance training improves soccer performance. *Med Sci Sports Exerc.* 2011;33:1925-1931.

- Hill-hass S, coutts G, Rowsell B, Dawson T. Generic versus small- sided game training in soccer. *Int Sport Med.* 2009;30:639-642.
- Hoff J, Wisloff U, Engen LC, Kemi OJ, Helgerud J. Soccer specific aerobic endurance training. *Br J Sports Med.* 2012;36:218-221.
- 12. Katis A, Kellis E. Effects of small-sided games on physical conditioning and performance in young soccer players. *Journal of Sports Science and Medicine*. 2009;8:374-380.
- Matavulj D, Kukolj M, Ugarkovic D, Tihanyi J, Saric S. Effects of plyometric training on jumping performance in junior basketball players. *J Sports Med Phys Fitness*. 2001;10:159-164.
- 14. Pedersen BK, Febbraio MA. Muscle as an Endocrine Organ: Focus on Muscle-Derived Interleukin-6. *Physiol Rev.* 2008;88:1379-1406.
- 15. Pederson BK, Fischer CP. Beneficial health effect of exercise- the role of IL-6 as a myokine. *Trends in Pharmacological Sciences*. 2007;28:149-156.
- 16. Rampinini E, Impellizzeri F, Castagna C, Abt G, Chamari K, Sassi A, et al. Factors influencing physiological responses to small sided games. J Sport Sci. 2006;25:659-666.
- 17. Sedano Campo S, Vaeyens R, Philippaerts RM, Redondo JC, De Benito AM, Cuadrado C. Effects of lower-limb plyometric training on body composition, explosive strength, and kicking speed in female soccer players. *J Strength Cond Res*. 2009;23:1714-1722.
- Steensberg A, Febbraio MA, Osada T, Schjerling P, van Hall G. Interleukin-6 production in contracting human skeletal muscle is influenced by pre-exercise muscle glycogen content. *J Physiol*. 2001;537:633–639.
- Steensberg A, van HG, Osada T. Production of interleukin-6 in contracting human skeletal muscles can account for the exercise-induced increase in plasma interleukin-6. *J Physiol*. 2000;529:237-242.

تاثیر دو شیوه تمرینی بازی در ابعاد کوچک (SSG) و مقاومتی - پلایومتریک بر سطوح استراحتی اینترلوکین -6 و عامل نکروز تومور آلفا در بازیکنان نوجوان فوتبال

سعيد رستمى (دكتر محمد فرامرزى) كلناز ارباب] ليلا احمدى أ

^۱ مربی گروه تربیت بدنی، دانشگاه آزاد اسلامی، واحد مبارکه، ^۲ دانشیار گروه تربیت بدنی، دانشگاه شهرکرد ^۲ دانشجوی دکترای تربیت بدنی، ^٤ کارشناس ارشد تربیت بدنی، دانشگاه آزاد اسلامی، واحد بندرعباس

مجله پزشکی هرمزگان سال هجدهم شماره چهارم مهر و آبان ۹۳ صفحات ۳۴۶–۳۳۹

حكنده

مقدمه: هدف پژوهش حاضر، بررسی تأثیر ۸ هفته تمرین مقاومتی – پلایومتریک و بازی در ابعاد کوچک (SSG) بر غلظت اینترلوکین -٦ و عامل نکروز تومور آلفا بازیکنان پسر نوجوان فوتبال بود.

روش کار: بدین منظور ۲۰ بازیکن پسر فوتبال با دامنه سنی ۱۶–۱۷ سال (میانگین سن ۲۰/±۱۶ ۹/۰ سال، قد ۲۰/۰±۱۷/۱ متر، وزن ۱/۱۰±۲۲/۲ کیلوگرم) برای شرکت در این تحقیق انتخاب شدند و به طور تصادفی در سه گروه بازی در ابعاد کوچک (۸ نفر)، مقاومتی-پلایومتریک (۱۱ نفر) و کنترل (۱۱ نفر) تقسیم شدند. قبل و بعد از ۸ هفته تمرین و در حالت استراحتی 6-11 و ۲۰-۲۲ اندازه گیری شد. سپس آزمودنیها به مدت ۸ هفته در گروههای مربوطه تمرینات خود را انجام دادند. برنامه تمرین گروه SSG شامل ۸ هفته تمرین بازی در ابعاد کوچک بود که به دو مرحله ۶ هفتهای برای تنظیم شدت کار تقسیم شد و توسط ۸ نفر صورت میگرفت. برنامه تمرین مقاومتی – پلایومتریک نیز توسط ۱۱بازیکن اجرا شد. این برنامه شامل ۸ هفته تمرین ترکیبی مقاومتی – پلایومتریک بود که آزمودنیها تمرینات مقاومتی – پلایومتریک را دو بار برای مقایسه بین گروهی از آزمون ۲۵۷۸ و MOVA

فتاییج: نتایج این تحقیق نشان داد اجرای ۸ هفته تمرین بازی در ابعاد کوچک به افزایش β-II و TNF-α منجر شد (برای هر سه گروه (P=۰/۰۰۱) و در گروه مقاومتی – پلایومتریک تغییر معنی داری دیده نشد (به ترتیب P=۰/۰٥۲ و P=۰/۱۲۲.

نتیجهگیری: به طور کلی، میتوان نتیجه گرفت که در شیوه تمرینی SSG به علت شدت بالای تمرین و سطوح بالای هورمونهای استرسی، تخلیه ذخایر گلیکوژن عضله و همچنین آسیبهای ریز درون بافتی که به واسطه جایجاییهای سریع و ضربههای مکرر در فوتبال به وجود میآید، سایتوکینهای التهایی افزایش معنی داری مییابد. **کلیدواژ دها:** اینترلوکین ۲ – سایتوکین – بازیکن فوتبال نویسنده مسئول: سعید رستمی انشکه مدیریت و حسابباری دانشگاه اصفهان – ایران تلفن: ۱۰۸۸۲۲۱۱ ۹۸۰+ پست الکترونیکی: Srostami.20@gmail.com

دریافت مقاله: ۹۲/۲/۲۹ اصلاح نهایی: ۹۲/۵/۲۹ پذیرش مقاله: ۹۲/۶/۵