

*Original Article***Prevalence of Dental Occlusal Patterns and Their Association with Obstructive Upper Airway Diseases in Primary School Children, Isfahan, Iran**

SM. Sonbolestan MD*, M. Mirmohammadsadeghi MD**,
M. Rogha MD***, SH. Nemati MD****

ABSTRACT

Background: Teeth, apart from their physiologic function, play an important role in general appearance of individuals. Therefore any disorder in their growth and evolution, will cause psychologic, social and even economic problems for the person. This observational cross-sectional study aimed to investigate relationship between dental occlusal patterns and obstructive nasal-upper airway diseases.

Methods: This study was performed in schools of No.4 Education at district of Isfahan in educational year 1381-82 between 607 students (277 male, 330 female) aged 9-12 years old. Results of accurate ENT and dentistry physical examination were registered and analyzed using SPSS software and Chi-square and Mantel - Hanzel test.

Results: With increasing age, The frequency of abnormal occlusal patterns increases (from 45.5% to 68.2% in males and 25.8% to 48.5% in females, $p=0.015$). The increasing in degree of palatal tonsillar hypertrophy was related to higher frequency of abnormal occlusal patterns (36.7% in +1 tonsillar hypertrophy, and 70% in +4 tonsillar hypertrophy, $p=0.02$). Also, history of frequent common colds, and history of previous nasal fractures were related with abnormal patterns [58.9% ($p=0.032$), and 83.4% ($p=0.043$), respectively]. Five other parameters including sinusitis, hypertrophy of nasal turbinates, rhinitis, nasal polyposis and nasal septal deviation were not associated with abnormal occlusal patterns ($p>0.1$).

Conclusion: Some of obstructive upper airway diseases are related with abnormal dental occlusal patterns. These parameters can be simply diagnosed, treated or prevented.

Key words: dental occlusion, malocclusion, obstructive nasal disorders, nasal turbinates, adenoid, rhinitis, tonsillectomy, open mouth breathing

JRMS 2005; 10(6): 349-354

Because of the important role of teeth in individual general appearance, existence of any disorder in growth and evolution of these structures will cause some psychosocial problems for the involved persons. The term "malocclusion" means malposition and abnormal contact of maxillary and mandibular teeth that interferes with exclusive movements of the jaws that are essential for mastication and also bear a special poor appearance for the patient. Dental malocclusion causes some oral and periodontal disorders that their manage-

ment strategies generally are very complex and expensive for the individual and society. Time consumption for treating such a problem may interfere with social-economic function of the affected persons. Also, there are some obvious legal obstacles for employment of these individuals in some governmental organizations (such as airlines and so on).

Mouth-breathing secondary to relative or complex obstruction of nose or upper airway leads to abnormal bone growth and underdevelopment of craniofacial structures, and

*Associated Professor, Otolaryngology and Head and Neck Surgery Department, Isfahan University of Medical Sciences, Isfahan, Iran.

**Resident, Otolaryngology and Head and Neck Surgery Department, Isfahan University of Medical Sciences, Isfahan, Iran.

***Assistant Professor, Otolaryngology and Head and Neck Surgery Department, Isfahan University of Medical Sciences, Isfahan, Iran.

****Resident, Otolaryngology and Head and Neck Surgery Department, Isfahan University of Medical Sciences, Isfahan, Iran.

Correspondence to: Dr. SM Sonbolestan, Otolaryngology and Head and Neck Surgery Department, Isfahan University of Medical Sciences, Isfahan, Iran. E-mail: sonbolestan@med.mui.ac.ir

therefore, can adversely affect dental occlusion and natural interaction of upper and lower jaws¹. This relationship has been shown in numerous studies, such as those performed by Hannuksela², Freg², Linder Aronson³ and in several other studies⁴⁻⁹. There are some other studies that are less conclusive about this relationship, such as those performed by Koski², Woodshed DG.¹⁰ and Shanker S.¹¹ who had not confirmed such a relationship between obstructive nasal diseases and dental malocclusion. Because of these paradoxical findings in this subject and because diagnosis of etiologic factors play a major role in prevention and treatment of such abnormalities, we decided to study the effects of nasal and upper airway obstruction on dental occlusal changes.

Subjects and Methods

This was an observational, cross-sectional study, performed in schools of No.4 Educational district of Isfahan, Iran in educational year 2002-3. By stratified random sampling method, 607 students of these schools (277 male and 330 female) between 9-12 years old were included in our study. These students generally belong to middle socio-economic class and also were in a range of age which is between two growth peaks; a matter that is very important in development and evolution of deciduous and permanent teeth and growth of jaws. The children with syndromic faces or with previous history of adenotonsillectomy or uvulopalatopharyngoplasty were excluded from our study. In each student, an accurate history taking (from children and their parents) and initial physical examination of ear, nose and throat and also dental occlusion patterns were performed in their schools by two well educated interns and two ENT residents, and each suspicious case of malocclusion were referred to the dentist for further advanced physical examination. Palatal tonsillar hypertrophy was examined by inspection and scored based on Brodsky, Moore, and Stanievich scoring system⁽¹⁹⁾. Also, diagnoses of sinusitis and rhinitis were made clinically, based on history taking and direct inspection of anterior nasal

cavity (anterior rhinoscopy) and oropharynx for PND and mucosal changes, by means of appropriate nasal speculums, tongue blades, and headlight. Meanwhile, we inspected evidences of hypertrophic nasal turbinates, nasal polypsis, and septal deviation in our cases. The registered data were computerized and analyzed, using

SPSS software and chi-square test, chi-square Mantel-Haenszel test, and variance analysis.

Results

In this study, the relationship between dental occlusal patterns and several parameters in nose and throat examination of our cases was studied. The results of our study are summarized in tables 1 and 2.

In all age groups, occlusion class I (normal) had the highest frequency, and then, the class II and III (abnormal patterns) were more frequent, respectively. With increasing the age, the frequency of abnormal occlusal patterns increased (from 45.5% to 68.2% in males and 25.8% to 48.5% in females, $P=0.015$). In male subjects, abnormal occlusal patterns, were more frequent, relative to females. Palatal tonsillar hypertrophy was examined and scored, based on Brodsky, Moore, and Stanievich scoring system¹⁹, in which, +1 hypertrophy means that the tonsils occupy less than 25% of distance from midline to tonsillar pili, and +2, +3, +4 means that 25-50%, 50-75%, and more than 75% of this distance are occupied by the tonsils, respectively¹⁹. According to our study, the increase in degree of palatal tonsillar hypertrophy were related with higher frequency of abnormal occlusal patterns (36.7% in +1, and 70% in +4 palatal tonsillar hypertrophy; $P=0.02$). Also, history of frequent common colds, that means more than 6 episodes of common cold in each year, and history of nasal bone fracture, were statistically related to abnormal occlusal patterns [relative frequency in each group was 59.5%, ($P=0.006$), 58.9% ($P=0.032$), and 83.4%, ($P=0.047$), respectively]. Five other parameters such as sinusitis ($P=0.49$), nasal turbinate hypertrophy

($P=0.219$), rhinitis ($P=0.422$), nasal polyposis ($P=0.408$) and nasal septal deviation ($P=0.306$) were not significantly related to dental occlusal abnormal patterns. One parameter, (i.e. nasal tumor), was not found in our study sample and its relationship with dental occlusal pattern was impossible to be determined.

In this study, we have also determined the relationship between these different occlusal patterns and mean number of abnormal parameters in upper airway physical examination. ($P= 0.173$). This is summerized in Table 2.

Table 1. Different dental Occlusal patterns and their relationship with some parameters in upper airway physical examination, observed in 607 students, aged 9-12 yearsold, Isfahan, Iran, 1382.

Parameter	N (frequency:%)	Occlusion class I (%)	Occlusion Class II (%)	Occlusion class III (%)
Age and gender *	9 yr (M)	44(7.2)	24(54.5)	19(43.2)
	10yr (M)	94(15.5)	47(50)	43(45.7)
	11yr (M)	95(15.6)	54(56.8)	35(36.8)
	12yr (M)	44(7.2)	14(31.8)	27(61.4)
	9yr (F)	31(5.1)	23(74.2)	7(22.6)
	10yr (F)	57(9.4)	45(78.9)	12(21.1)
	11yr (F)	145(23.9)	89(61.4)	52(35.9)
	12yr (F)	97(15.9)	50(51.5)	46(47.4)
Relative size of palatal tonsils	+1	452(74.5)	237(60.4)	166(36.7)
	+2	90 (15)	46(51.1)	40(44.4)
	+3	55(9.1)	24(43.6)	28(50.9)
	+4	10(1.6)	3(30)	7(70)
Frequent common colds	51(8.4)	21(41.2)	29(56.9)	1(2)
Positive Hx. of nasal fracture	6(1)	1(16.7)	4(66.7)	1(16.7)
Sinusitis	20(3.3)	10(50)	10(50)	0(0)
Hypertrophied nasal turbinates	237(39)	144(60.8)	84(35.4)	9(3.8)
Rhinitis	142(23.4)	81(57)	54(38)	7(4.9)
Polyposis	9(1.5)	5(55.6)	3(33.3)	1(11.1)
Different variants of septal deviation	220(36.2)	118(53.6)	96(43.6)	6(2.7)
total	607(100)	346(57)	241(39.7)	20(3.3)

M: male, F: female, yr: years old, Hx: history.

Table 2. Different dental occlusal patterns and their relationship with mean number of abnormal parameters in upper airway physical examination.

Class of Occlusion	N	Mean number of abnormal parameters
I	346	1.3324(SD=0.72)
II	241	1.3324(SD=0.47)
III	20	1.4979(SD=0.4)
total	607	1.5500(SD=0.6)

Discussion

According to classification of Angle in 1899, three major types of dental occlusion have been defined as ¹²:

Occlusion class I (normal pattern): In this pattern, mesial buccal casps of the first molars of upper jaw, lie just on the buccal sulcus of their lower jaw equivalents.

Occlusion class II (abnormal pattern): In this pattern, mesial buccal casps of the first molars of upper jaw, lie more anteriorly than the buccal sulcus of their lower jaw equivalents and even may lie on sulci between lower jaw molars and second premolars.

Occlusion class III (abnormal pattern): In this pattern, opposite to the class II pattern, mesial buccal casps of the first molars of upper jaw, lie on distal casps or on sulci between first and second molars of lower jaw.

Chronic mouth breathing secondary to relative or complete nasal or upper airway obstruction, leads to abnormal craniofacial growth, and subsequently affects evolution of dental occlusion and interaction of upper and lower jaws. In human beings, there are two peaks in growth curve: the first in ages of 7-8 years old, and another in ages of 13-14 years old and puberty period ¹. If causes of upper airway obstruction and chronic mouth breathing are not obviated before these ages, their stigmata and sequelae will be established and their management in future will be very unlikely or even impossible.

We can classify the etiologic factors of nasal and upper airway obstruction as follow ²⁰:

1- Structural disorders:

- a: Deformities: external, internal, congenital malformations, injuries,
- b: Neoplasms and masses,
- c: Foreign bodies,

2- Inflammatory disorders:

- a: Rhinitis/sinusitis: bacterial, viral, fungal,
- b: Nasal and sinus polyposis,
- c: Ozena, atrophic rhinitis,
- d: Immunologic diseases: sarcoidosis, wegener's granulomatosis,

3- Allergic rhinitis,

4- Vasomotor rhinitis.

Adenotonsillar hypertrophy is one of the most common etiologic factors of upper airway obstruction, especially in children ¹⁹. Postural changes secondary to chronic mouth breathing lead to disequilibrium in normal forces that affect teeth, and also, soft and hard tissues of orofacial structures, and there after may cause some sequelae such as open bite, protrusion of maxilla, posterior cross bite, and so on ¹⁹. It is interesting to mention that some of these aspects may resolve after medical or surgical interventions such as adenotonsillectomy ¹⁹. Meanwhile, some of etiologic factors may be simply preventable.

Several studies have shown some aspects of relationship between causes of upper respiratory tract obstruction and dental malocclusion. For example, Linder-Anderson and Oulis C.J. Showed the relationship between enlarged tonsils and craniofacial dysmorphology ¹³, and Lopatien K. in 2002, found some causative relations between upper respiratory tract obstruction and malocclusion ¹⁴, but these studies are not thorough and have not considered all of causes of upper airway obstruction, and on the other hand, some other studies have not met to similar results and there are many controversies in this regard. At last, we did not have any documented study about prevalence of dental occlusal patterns in our patients, especially pediatric-age groups, in whom some causes of upper airway obstruction such as adenotonsillar hypertrophy are common. Therefore, we decided to study the relative frequency of different occlusal patterns in a group of students aged between 9-12 years old, and also show the role of nasal and upper airway obstructive disorders in creation of abnormal occlusal patterns. This population of cases has passed their first growth peak, and meanwhile, and we can find some upper airway obstructive diseases such as adenotonsillar hypertrophy in them that can be managed to prevent further sequelae in their craniofacial growth through their second growth peak.

• According to our results (tables 1,2), occlusion class I (normal pattern) had the highest frequency in all age groups, except in 12 years old boys, which was related to this age and gender of these subgroup. After class I occlusion, the classes II and III occlusions (abnormal patterns) were more frequent, respectively, which is concordant with reference books ¹². It is obvious that with increasing the age, the frequency of occlusion class I reduces and substitutely, the frequency of occlusion classes II and III increase. Therefore, the frequency of abnormal occlusions increases with age and this relationship is statistically significant.

• In male subjects, relative to females, normal occlusion had lower and abnormal occlusions had higher frequencies. This is similar to ob-

servations of Woodside D.G. in 1991, who found that, abnormal occlusal patterns are more frequent in boys who underwent adenoidectomy than in girls underwent this surgery¹⁰.

- ♦ Hypertrophy of palatal tonsils was related statistically with frequency of abnormal dental occlusions ($p=0.02$). This is compatible with findings of other studies, mentioned previously^{13, 14}.

- ♦ Frequent common colds and history of nasal bone fracture, were meaningfully related with the type of dental occlusal pattern ($p=0.032$ and 0.047 , respectively). We could not find anything in this regard in our review of textbooks and articles.

- ♦ Our observations suggest that in students with sinusitis and nasal septal deviation, there were relatively lower frequency of normal occlusion and higher frequency of occlusion class II, compared with healthy students; however, chi-square test did not show any significant relationship between them. Also, there isn't any similar study in this regard in medical references.

- ♦ Nasal polyposis and hypertrophy of turbinates had no valid relationship with occlusal patterns. Also in this regard we could not find any similar study in references.

- ♦ Relative frequency of occlusion class I in any of two groups of students, with and without rhinitis, was similar; also, frequency of class II occlusion was approximately equal (even a little lower in rhinitis group). Abnormal occlusion class III was more frequent in rhinitis group, but this is not statistically significant, according to chi-square test. Because of discordance between these results and results of other studies such as that of Martinez Estenous Ji. In 1998¹⁵, and Bertolani MF in 2004¹⁶, we suggest complementary and advanced studies in this regard, especially, when we consider that in the first study, results of radiologic criteria were concordant with results of this study.

- ♦ As we demonstrate in table 3, individuals who had abnormal occlusal patterns, had averagely further obstructive problems com-

pared to individuals who had normal occlusal patterns; however, this is not statistically significant. Numerous studies have been done in this regard, and some of them are concordant with our results, and some others are not. For example, the results of observations of Kluemper GT. in 1995¹⁷, Shanker S. in 1999¹¹, and Faria in 2002¹⁸ are similar to our study results, while, those of Lofstrand in 1999⁸ and other studies are not. Therefore we suggest further studies in this regard.

Conclusion

According to results of this study and some other ones about influence of obstructive upper airway diseases on craniofacial growth and their role as etiologic, or at least, as comorbid factors in generation of abnormal dental occlusion patterns, we can conclude that early diagnosis and treatment of such etiologic factors can prevent aberrant changes in craniofacial and dentofacial region and abnormal sequelae in appearance of individuals, and reduce necessity of advanced and very expensive orthodontic treatments. This is achievable simply by two ENT and dentistry visits, and like other screening tests, such as thalassemia test before marriage, or thyroid function test in newborns, can achieve to its important place in preventive medicine. Also, these diagnostic-therapeutic measures can reduce relapses of orthodontic treatments which are done for correction of these malformed dento-facial appearances. To achieving these purposes, a thorough coordination between otolaryngologist, pediatrician, orthodontist, and pediatric dentist is mandatory, and also, utilization of advanced instruments and equipments, such as rhinomanometry, rhinoscopy, platysmography and so on is suggested.

Acknowledgement

We thank Dr. Sadegh Roghaee (DDS), A.H. Moghbel (intern), A. Nasr (intern) a lot, for their contribution in this research.

References

1. Proffit WR, Field HW. *Contemporary-orthodontics*. 3rd-ed. Mosby; 2000.
2. Naspitz CK, Tickelman DG. *Childhood Rhinitis and Sinusitis*. Marcel Dekker WC; 1990.pp 170-3.
3. Akhavan Niaki E. *Cross-bite, Side effects and treatments*. Tehran University Publication 1996; 2270:237-43.
4. Harvold EP, Tomer BS, Vargervik K, Chierici G. *Primate experiments on oral respiration*. Am J Orthod 1981 Apr;79(4):359-72.
5. Bresolin D, Shapiro PA, Shapiro GG, Chapko MK, Dassel S. *Mouth breathing in allergic children: its relationship to dentofacial development*. Am J Orthod 1983 Apr;83(4):334-40.
6. Josell SD. *Habits affecting dental and maxillofacial growth and development*. Dent Clin North Am 1995 Oct;39(4):851-60.
7. Baumam I, Plindert PK. *Effect of breathing mode and nose ventilation on growth of the facial bones*. HNO 1996 May;44(5):229-34. German.
8. Lofstrand-Tidestrom B, Thilander B, Ahlqvist-Rastad J, Jakobsson O, Hultcrantz E. *Breathing obstruction in relation to craniofacial and dental arch morphology in 4-year-old children*. Eur J Orthod 1999 Aug;21(4):323-32.
9. Schlenker WL, Jennings BD, Jeiroudi MT, Caruso JM. *The effects of chronic absence of active nasal respiration on the growth of the skull: a pilot study*. Am J Orthod Dentofacial Orthop 2000 Jun;117(6):706-13.
10. Woodside DG, Linder-Aronson S, Lundstrom A, McWilliam J. *Mandibular and maxillary growth after changed mode of breathing*. Am J Orthod Dentofacial Orthop 1991 Jul;100(1):1-18.
11. Shanker S, Vig KW, Beck FM, Allgair E Jr, Vig PS. *Dentofacial morphology and upper respiratory function in 8-10-year-old children*. Clin Orthod Res 1999 Feb;2(1):19-26.
12. Ballenger JJ, Snow JB. *Otorhinolaryngology: Head and Neck Surgery*. Baltimore: Williams and Wilkins; 1996.
13. Linder-Aronson S. *Craniofacial morphology in children with and without enlarged tonsils*. Europ J Orthod 1990;12(3):233-43.
14. Lopatiene K, Babarskas A. *Malocclusion and upper airway obstruction*. Medicina (kaunas) 2002; 38(3):277-83.
15. Martinez Esteinou JL, Omana Vidal E. *Dental malocclusion and bony abnormalities in girls with nasopharyngeal obstruction of allergic origin*. Pract Odontol 1988 Dec;9(12):8, 10, 12 passim.
16. Bertolani MF. *Alternaria spores at different heights from the ground*. Allergy 2004 Jul; 59(7): 746-52.
17. Kluemper GT, Vig PS, Vig KW. *Nasorespiratory characteristics and craniofacial morphology*. Eur J Orthod 1995 Dec;17(6):491-5.
18. Faria PT, de Oliveira Ruellas AC, Matsumoto MA, Anselmo-Lima WT, Pereira FC. *Dentofacial morphology of mouth breathing children*. Braz Dent J 2002; 13(2):129-32.
19. Wiatrak BJ, Wooley AL. *Pharyngitis and adenotonsillar diseases*. In: Johnson JT, Charls W editors. Cummings, Otolaryngology head & neck surgery. 3rd edition, Vol 5. Mosby; 1998.
20. Fairbanks DNF, Kaliner M. *Nonallergic Rhinitis and Infections*. In: Johnson JT, Charls W editors. Cummings, Otolaryngology head & neck surgery. 3rd edition, Vol 5. Mosby; 1998.