

## Techniques for environmental monitoring of predatory fauna on branches of Bramley apple trees in Northern Ireland

<sup>1\*</sup>A. G. S. Cuthbertson and <sup>2</sup>A. K. Murchie

<sup>1</sup>Central Science Laboratory, Sand Hutton, York YO41 1LZ, United Kingdom

<sup>2</sup>Department of Agriculture and Rural Development for Northern Ireland and the Queen's University of Belfast, Newforge Lane, Belfast BT9 5PX, United Kingdom

### Abstract

Ranges of trapping devices were investigated for their ability to sample predatory mites and insects on branches of apple trees. Straw traps caught significantly more predatory mites whereas corrugated cardboard traps proved better for larger predators such as the common earwig, *Forficula auricularia*. The most abundant predatory species sampled on the trees was the 'whirligig' mite *Anystis baccarum*. The efficiency of the various trapping devices and the potential of incorporating *A. baccarum* within environmentally based management strategies for invertebrate pest control within the Bramley apple orchards are discussed.

**Key words:** *Anystis baccarum*, biodiversity, Bramley's Seedling apple, *Forficula auricularia*, pest management, predatory fauna, trapping

\*Corresponding Author, E-mail: [a.cuthbertson@csl.gov.uk](mailto:a.cuthbertson@csl.gov.uk)

### Introduction

A wide range of predators have been exploited, or have the potential to be exploited, as biological control agents against pests of fruit. In the British Isles, these include members of the families Miridae (mirid bugs) (Hemiptera), Anthocoridae (flower bugs) (Hemiptera), Forficulidae (earwigs) (Dermaptera) and several mite families (Solomon, 1992). Leaf-dwelling predators such as the mites *Typhlodromus pyri* and *Zetzellia mali* can be sampled by leaf washings (Zacharda, *et al.*, 1988). However, larger predatory species are generally more mobile and cannot be sampled in this way as they disperse when vegetation is disturbed. Various authors have used artificial shelter traps for trapping *F. auricularia* and other insect fauna (Chant and McLeod, 1952; Campbell, 1978; McAdam, *et al.*, 1994 and Helsen, *et al.*, 1998) and mites (Morgan and Anderson, 1958). Similarly, sticky traps in the form of grease bands have been used to trap mites (Staples and Allington, 1959) and mirids (Stewart and Gaylor, 1991). However, little information is available regarding the effectiveness of different trapping systems in sampling the predatory fauna on branches of apple trees. The aim of this study was to compare several different techniques for trapping and hence sampling the predatory fauna from branches of apple trees, and to determine the

most abundant predatory species within Northern Irish apple orchards.

### Materials and Methods

#### Study orchard

The study site was a commercially active Bramley's seedling apple orchard at Loughgall, County Armagh, Northern Ireland (N 54° 24.3' W 006° 35.8'). This orchard received normal fungicide spray treatment to control apple scab (*Venturia inaequalis*) but no acaricide treatments.

#### Interception traps

One of each trap type (Table 1) was attached to different branches of each of ten randomly chosen trees, from 17 March until 28 August 1997. The traps were changed at fortnightly intervals and placed under Tullgr n funnels for 24 hours where predatory arthropods were expelled into 30% ethanol. All specimens were identified and recorded. Numbers of predatory mites and insects captured in the traps were subjected to analysis of variance (ANOVA).

#### Leaf washing

Over the same time period as the trapping experiment, leaf samples were taken at fortnightly intervals from the apple trees (8 leaves per sample date) and, following the method of Zacharda *et al.*

Table 1: Trapping devices used to capture predatory fauna on branches of apple trees

Trap type	Construction	Dimensions (L= length, W= width, D= depth)
Straw	Nylon mesh bag (5 x 3 mm. rectangular mesh) containing heat-sterilised (48 h. at 55°C) chopped barley straw	L- 50 mm., W- 90 mm.
Corrugated Cardboard	Corrugated cardboard (L-150 mm., W-190 mm., 3 mm ridged cardboard) rolled up inside open-ended plastic cylinders	L- 200 mm., Dia. 50 mm.
Ridged Corriboard (Single sheet)	Single sheet of laminated corrugated Corriboard	L-140 mm., W-90 mm., D-4 mm.
Multi-layer Corriboard trap (Layered)	Three sheets of laminated corrugated Corriboard	L-14 cm. , W-9 cm., D-13 mm.
Sticky	Single-sided yellow sticky continuous trap paper ( <i>Agralan Ltd.</i> Swindon, UK) wrapped around the branch	W-120 mm.

Table 2: Mean number ( $\pm$  SEM) of predatory mites per leaf ( $n=8$ ) averaged over 12 fortnightly samples from 17 March until 28 August 1997 in a non-acaricide sprayed Bramley orchard

Predatory mite	Family	Mean per leaf
<i>Typhlodromus pyri</i>	Phytoseiidae	0.15 $\pm$ 0.11
<i>Phytoseius macropilis</i>	Phytoseiidae	0.11 $\pm$ 0.08
<i>Zetzellia mali</i>	Stigmaeidae	0.10 $\pm$ 0.08

(1988), leaf dwelling predators were sampled. Numbers of predatory species washed from the leaves were identified and recorded.

## Results

### Leaf washing

Only three species of predatory mites were washed from leaves. In order of abundance they were *T. pyri*, *Phytoseius macropilis* and *Z. mali* (Table 2).

### Interception traps

The most numerous predatory species obtained from the traps was the generalist predatory mite, *Anystis baccharum* (Table 3).

Straw traps trapped more predatory mites ( $F=11.38$ , d.f.= 4,36,  $P<0.01$ ), including *A. baccharum*, compared to the other traps. However, corrugated cardboard traps trapped the majority of larger predators ( $F= 2.99$ , d.f.= 4,36,  $P= 0.03$ ) (Table 3); in particular, *F. auricularia*. The sticky traps were impractical as they quickly became covered with flies

which made sorting of small predatory mites extremely difficult.

## Discussion and Conclusion

The presence of the phytoseiid and stigmaeid mites in leaf washings from this non-acaricide sprayed orchard was expected. *Typhlodromus pyri* is widespread and abundant on various trees, shrubs, and herbaceous plants (Collyer, 1964) and is common in both intensively managed and wild orchards. In the present study, the density of *T. pyri* at 0.15 mites per leaf was smaller than that in a dessert orchard in southern England where the population reached 0.25 per leaf (Cross and Berrie, 1994). Similarly, *P. macropilis* is a widespread predator of both eriophyid (rust mites) and tetranychid (red spider mites) mites on several host plants, including unsprayed apple and plum trees (Alford, 1984). *Zetzellia mali* is also considered a common predator of *Aculus schlechtendali* (rust mite) and *Panonychus ulmi* (red spider mite) on fruit trees (Santos, 1976; Alford, 1984).

Table 3: Mean number ( $\pm$  SEM) of predatory fauna captured in ten traps averaged over 12 fortnightly samples from 17 March until 28 August 1997 in a non-acaricide sprayed Bramley apple orchard.

Trap type							
Predatory mites		Family	Straw	Cardboard	Corriboard <sup>1</sup>	Corriboard <sup>2</sup>	Sticky
<i>Typhlodromus pyri</i>		Phytoseiidae	1.20 ±0.54	0.60 ±0.29	0.20 ±0.26	0.10 ±0.13	0.40 ±0.29
<i>Typhlodromus richteri</i>		Phytoseiidae	0.50 ±0.29	0.00	0.00	0.00	0.60 ±0.34
<i>Phytoseius macropilis</i>		Phytoseiidae	0.70 ±0.28	0.30 ±0.20	0.10 ±0.13	0.30 ±0.28	0.20 ±0.17
<i>Zetzellia mali</i>		Phytoseiidae	0.50 ±0.22	0.20 ±0.17	0.10 ±0.13	0.00	0.00
<i>Anystis baccarum</i>		Acaridae	7.60 ±3.48	0.80 ±0.71	1.50 ±0.75	1.10 ±0.83	0.70 ±0.20
<i>Parasitus fimetorum</i>		Parasitidae	0.50 ±0.22	0.00	0.00	0.00	0.00
<i>Gamasodes spiniger</i>		Parasitidae	0.20 ±0.17	0.00	0.00	0.00	0.00
<i>Allothrombium fuliginosum</i>		Trombididae	0.00	0.20 ±0.17	0.10 ±0.13	0.00	0.10 ±0.13
<i>Glycyphagus domesticus</i>		Acaridae	0.00	0.00	0.00	0.10 ±0.13	0.00
Predatory insects							
<i>Anthocoris nemoralis</i>		Anthocoridae	0.20 ±0.17	0.30 ±0.27	0.10 ±0.12	0.10 ±0.12	1.50 ±1.23
<i>Anthocoris nemorum</i>		Anthocoridae	0.30 ±0.27	0.40 ±0.21	0.40 ±0.28	0.10 ±0.12	0.80 ±0.32
<i>Forficula auricularia</i>		Forficulidae	0.60 ±0.28	4.70 ±1.57	0.00 ±0.72	1.70 ±0.47	1.20 ±0.57
<i>Coccinella</i>							
<i>Septempunctata</i>		Coccinellidae	0.00	0.50 ±0.28	0.10 ±0.12	0.00	0.00
<i>Propylea</i>							
<i>Quatuordecimpunctata</i>		Coccinellidae	0.00	0.10 ±0.12	0.10 ±0.12	0.00	0.00

<sup>1</sup>Single sheet of ridged corriboard

<sup>2</sup>Multi-layered corriboard

The most abundant predatory species caught in traps and visible on the trees in this orchard was the predatory mite, *Anystis baccarum* (Linnaeus). *Anystis baccarum* has been little researched in orchards in the British Isles, and their abundance remains relatively unknown (Cuthbertson and Murchie, 2004a). As all members of this family are generalist predators which feed readily on a wide range of arthropods (Baker, 1967), this species could be a valuable natural enemy of orchard pests (Cuthbertson, 2000; Cuthbertson and Murchie, 2000; Cuthbertson and Murchie, 2004b).

The most abundant predatory species caught in traps and visible on the trees in this orchard was the predatory mite, *Anystis baccarum* (Linnaeus). *Anystis baccarum* has been little researched in orchards in the British Isles, and their abundance remains relatively unknown (Cuthbertson and Murchie, 2004a). As all members of this family are generalist predators which feed readily on a wide range of arthropods (Baker, 1967), this species could be a valuable natural enemy of orchard pests (Cuthbertson, 2000; Cuthbertson and Murchie, 2000; Cuthbertson and Murchie, 2004b). Baker (1983) noted that *A. baccarum* preyed on leafrolling caterpillars in apple orchards in New Zealand and Putman and Herne (1966) found *A. baccarum* numbers to increase in orchards during outbreaks of *P. ulmi* in Ontario. Cuthbertson *et al.* (2003a,b) and Cuthbertson and Murchie (2003a; 2004a) went on to show that *A. baccarum* readily preyed upon *A. schlechtendali*, *P. ulmi*, *Rhopalosiphum insertum* (apple-grass aphid) and *Bryobia rubrioculus* (apple brown mite) within the Bramley apple orchards. The former three of these species are economically important pests in Northern Irish apple orchards (Mowat and Clawson, 1996).

A range of other common predatory arthropods were found in the traps. These are mainly generalist predators but will also be instrumental in reducing pest numbers to some extent. The most common large predator was *F. auricularia* which probably makes an important contribution to predation of pest species but they are of dubious value in orchards because they can also cause damage to the apples (Croxall *et al.*, 1951; Sial, 1957). Another predator caught, the ladybird, *Coccinella septempunctata*, is attracted into orchards by outbreaks of aphids (Collyer, 1953). However, if as a result of spraying or predatory activity the aphids become scarce, the larvae of this species will also feed upon *P. ulmi*. In this way a slight infestation of aphids in May and June may

contribute to *P. ulmi* control by attracting adult *C. septempunctata* into the orchards (Collyer, 1953).

Predacious mites and insects are usually sampled by the same method as the prey they feed on (Huffaker *et al.*, 1970). However, this method may not always present a complete picture of the populations of certain species. Putman and Herne (1964) found that a large proportion of the population of *Typhlodromus caudiglans* Schuster, was found on twigs. The difficulty arises because some species move back to sheltering crevices or rough surfaces when fully fed or during daytime. In addition, many insect predators of mites may drop from the leaves or fly away when leaves or twigs are being removed from the plants (Huffaker *et al.*, 1970). The use of traps offers shelter from the climate by mimicking natural crevices and prevents foliage or twigs from being disturbed when sampling.

Each type of trap caught both predatory mites and insects. The straw traps proved the best for trapping mites, especially *A. baccarum*. This may simply be related to the size of the mesh. Larger predators such as *F. auricularia* may have had difficulty trying to squeeze through the mesh openings. When inside the trap, the straw would offer good protection from changing climatic conditions and also a possible shelter for prey. Collembola were commonly found in the straw traps where *A. baccarum* were observed to readily prey upon them (Cuthbertson, AGS, personal observation).

The corrugated cardboard traps were useful for trapping fauna in hedgerow restoration studies (McAdam *et al.*, 1994). In this study, *F. auricularia* were taken more frequently in this trap compared to the others. Campbell (1978) and Helsen *et al.* (1998) found corrugated cardboard traps to work well for trapping *F. auricularia* which aligned themselves in the corrugations during the day and returned to the same roost each night. The corboard traps trapped larger predators in much the same way as the corrugated cardboard traps.

This study has highlighted the need for a range of assessment techniques for sampling predatory fauna on apple trees. Lord (1965) commented that there is no ideal method for monitoring predators on apple trees because of their non-random distribution, variation in foliage growth, and the active nature of many predators. The tendency to sample for beneficial mite predators such as *T. pyri* and *Z. mali* by washing leaves may lead to an underestimation of the role of more mobile mite predators such as *A. baccarum*. Cuthbertson and Murchie (2004a)

investigated the seasonal abundance of *A. baccharum* within Northern Irish orchards and demonstrated how its population peaked when pest species, such as, *P. ulmi*, are beginning to increase in number. *Anystis baccharum*, the most abundant predator within the orchard, has the potential to control various invertebrate pest species (Cuthbertson *et al.*, 2003a,b; Cuthbertson and Murchie, 2004a) and also is compatible with various chemical fungicides (Cuthbertson and Murchie, 2003a). Therefore, encouraging the presence and population build up of *A. baccharum* and other predatory fauna within the orchards will help reduce pest populations. Also, enabling fruit growers to recognise *A. baccharum* as a beneficial species will help ensure its survival within the orchard ecosystem (Cuthbertson and Murchie, 2003b; Cuthbertson, 2004, 2005). The greater the biodiversity of predatory fauna within a relatively stable ecosystem like an orchard, then the more natural control and less reliance on chemical insecticides. This also alleviates consumer concerns, justified or not, regarding the build-up of chemical residues on fruit produce and helps move the apple industry into a more environmentally sustainable production system.

This study has shown that a range of trapping devices are necessary for sampling predatory fauna on branches of apple trees. *Anystis baccharum* is the most abundant predatory species sampled in the orchard. This mite has potential to be incorporated into environmentally based management strategies for the control of invertebrate pests. Fruit growers should be encouraged to conserve, not only *A. baccharum*, but all predatory fauna populations within their orchards.

### Acknowledgements

Dr Andrew G. S. Cuthbertson was funded by a Department of Agriculture and Rural Development (Northern Ireland) Studentship.

### References

- Alford, D. V., *A colour atlas of fruit pests, their recognition, biology and control*. Wolfe Publishing Ltd. London: 320, 1984
- Baker, R. T., *Predation of leafroller larvae by spiders and mites*. *Weta*, **6**: 22-23, 1983
- Baker, W. V., *Some observations on predation in an anystid mite*. *Entomol. Mon. Mag.*, **103**: 58-59, 1967
- Campbell, C. A. M., *Regulation of the damson-hop aphid (Phorodon humuli (Schränk)) on hops (Humulus lupulus L.) by predators*. *J. of Hort. Sci.*, **53**: 235-242, 1978
- Chant, D. A. and J. H. McLeod, *Effects of certain climatic factors on the daily abundance of the European earwig, Forficula auricularia L. (Dermaptera: Forficulidae)*, in Vancouver, British Columbia. *Can. Entomol.*, **84**: 174-180, 1952
- Collyer, E., *Biology of some predatory insects and mites associated with the fruit tree red spider mite (Metatetranychus ulmi (Koch)) in south-eastern England. IV The predator-mite relationship*. *J. of Hort. Sci.*, **28**: 246-259, 1957
- Collyer, E., *Phytobagous mites and their predators in New Zealand orchards*. *New Zeal. J. of AGR. RES.*, **7**: 551-558, 1954
- Cross, J. and A. M. Berrie, *Effects of repeated foliar sprays of insecticides or fungicides on organophosphate-resistant strains of the orchard predatory mite Typhlodromus pyri on apple*. *Crop Protection*, **13**: 39-44, 1994
- Croxall, H. E., Collingwood, C. A. and J. E. E. Jenkins, *Observations on brown rot (Sclerotinia fructigena) of apples in relation to injury caused by earwigs (Forficula auricularia)*. *Ann. of Appl. Biol.*, **38**: 833-843, 1951
- Cuthbertson, A. G. S., *The role of the predatory mite, Anystis baccharum, in Bramley apple orchards*. Ph.D Thesis, The Queen's University of Belfast, UK, 170, 2000
- Cuthbertson, A. G. S., *Unnecessary pesticide applications in Northern Ireland apple orchards due to mis-identification of a beneficial mite species*. *Research Journal of Chemistry and Environment*, **8** (3): 77-78, 2004
- Cuthbertson, A. G. S., *Re-discovery of a predatory mite in Northern Irish apple orchards*. *Biodiversity News*, **30** (1): 29, 2005
- Cuthbertson, A.G.S. and A.K. Murchie, *The prospect of biological control within Northern Ireland Bramley orchards*. *Biology and Environment: Proceedings of the Royal Irish Academy*, **100**: 50, 2000.
- Cuthbertson, A. G. S. and A. K. Murchie, *The impact of fungicides to control apple scab (Venturia inaequalis) on the predatory mite Anystis baccharum and its prey Aculus schlechtendali (apple rust mite) in Northern Ireland Bramley orchards*. *Crop Protection*, **22**: 1125-1130, 2003a



- Cuthbertson, A.G.S. and A.K. Murchie, The need for proper identification of beneficial and pest species in the Bramley apple orchards. Online: [www.ruralni.gov.uk/crops/fruit/top\\_fruit/technical\\_information/proper\\_identification.htm](http://www.ruralni.gov.uk/crops/fruit/top_fruit/technical_information/proper_identification.htm). Accessed January 2005: 3, 2003b
- Cuthbertson, A. G. S. and A. K. Murchie, The phenology, oviposition and feeding rate of *Anystis baccarum*, a predatory mite in Bramley apple orchards in Northern Ireland. *Exp. Appl. Acarol.*, **34**: 367-373, 2004a
- Cuthbertson, A. G. S. and A. K. Murchie, The presence of *Anystis baccarum* (L.) in Northern Ireland Bramley apple orchards. *Ir. Nat. J.*, **27**: 465-467, 2004b
- Cuthbertson, A. G. S., Bell, A. C. and A. K. Murchie, The impact of the predatory mite *Anystis baccarum* on apple rust mite (*Aculus schlechtendali*) populations in Northern Ireland Bramley orchards. *Ann. Appl. Biol.*, **142**: 107-114, 2003a
- Cuthbertson, A. G. S., C. C. Fleming, and A. K. Murchie, Detection of *Rhopalosiphum insertum* (apple-grass aphid) predation by the predatory mite *Anystis baccarum* using molecular gut analysis. *Agric. Forest Entomol.*, **5**: 219-225, 2003b
- Helsen, H., Vaal, F. and L. Blommers, Phenology of the common earwig *Forficula auricularia* L. (Dermaptera:Forficulidae) in an apple orchard. *Int. J. of Pest Manage.*, **44**: 75-79, 1998
- Huffaker, C. B., M. Van de Vrie, and J. A. McMurtry, Ecology of tetranychid mites and their natural enemies: a review II. Tetranychid populations and their possible control by predators: A evaluation. *Hilgardia*, **40**: 391-458, 1970
- Lord, F. T., Sampling predator populations on apple trees in Nova Scotia. *Can. Entomol.*, **97**: 287-298, 1965
- McAdam, J. H., Bell, A. C. and T. Henry, The effect of restoration techniques on flora and microflora of hawthorn-dominated hedges. In: Watt, T. A. and G. P. Buckley (eds): *Hedgerow Management and Nature Conservation*. Wye College Press. Wye: 25-32, 1994
- Morgan, C. V. G. and N. H. Anderson, Techniques for biological studies of tetranychid mites, especially *Bryobia arborea* M. & A. and *B. praetiosa* Koch. (Acarina:Tetranychidae). *Can. Entomol.*, **90**: 212-215, 1958
- Mowat, D. J. and S. Clawson, The need for pest control in Northern Ireland Bramley apple orchards. *Crop Protection in Northern Britain*, **2**: 225-230, 1996
- Putman, W. L. and D. H. C. Herne, Relations between *Typhlodromus caudiglans* Schuster (Acarina:Phytoseiidae) and phytophagous mites in Ontario peach orchards. *Can. Entomol.*, **96**: 925-943, 1964
- Putman, W. L. and D. H. C. Herne, The role of predators and other biotic agents in regulating the population density of phytophagous mites in Ontario peach orchards. *Can. Entomol.*, **98**: 808-820, 1966
- Santolucito, J. Evaluation of *Zetzellia mali* as predator on *Panonychus ulmi* and *Aculus schlechtendali*. *Environ. Entomol.*, **5**: 187-191, 1976
- St. J. N. M., The earwig as an orchard pest, *Fruit Yearbook*. Royal Horticultural Society, London, UK: 140-145, 1957
- Solomon, M. G., Exploitation of predators in UK fruit and hop culture. *Phytoparasitica*, **20**: 51-56, 1992
- Staples, R. and W. B. Allington, The efficiency of sticky traps in sampling epidemic populations of the Eriophyid mite *Aceria tulipae* (K.), vector of wheat streak mosaic virus. *Ann. Entomol. Soc. Am.*, **52**: 159-164, 1959
- Stewart, S. D. and M. J. Gaylor, Age, sex, and reproductive status of the tarnished plant bug (Heteroptera:Miridae) colonizing mustard. *Environ. Entomol.*, **20**: 1387-1392, 1991
- Zacharda, M., O. Pultar, and J. Muska, Washing technique for monitoring mites in apple orchards. *Expe. Appl. Acarol.*, **5**: 181-183, 1988