Effects of wastewater stabilization pond effluent on agricultural crops

^{1*}K. Naddafi, ²N. Jaafarzadeh, ¹M. Mokhtari, ³B. Zakizadeh and ⁴M.R. Sakian

¹Department of Environmental Health Engineering, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

²Department of Environmental Health, Ahwaz University of Medical Science, Ahwaz, Iran ³Department of Environment, Khuzestan Industrial Estate Company, Ahwaz, Iran ⁴Water Divisions, Khuzestan Water and Power Organization, Ahwaz, Iran

Abstract

Wastewater reclamation and reuse is one of the best alternatives for compensating water shortages. Water supply and environmental conservation can be met through wastewater reclamation. Principally, treated wastewater is a reliable water resource, especially for periodic droughts and in arid areas. This study designed and implemented to investigate the full scale application of effluent for irrigation use. A major objective of this study is to assess on health effects and feasibility of crop irrigation by using stabilization ponds effluent of southern Hovaizeh Wastewater Treatment Plant located in Khuzestan Province. Two experimental plots of about 0.5 ha. were constructed. One of the plots irrigated by stabilization pond effluent and the other irrigated by Nisan River water. Basic parameters in two plots such as type of cultivated crops, amount of fertilizer use and lack of soil contamination have been similar in both. The only difference was the type of water applied for irrigation of agricultural crops. It was shown that high salinity of soil reduced the growth rate of agricultural crops growth in two studied plots showed the growth rate and quality of crops were increased by using of stabilization pond effluent in comparison with Nissan River water.

Key words: *Reuse, stabilization pond, effluent, agricultural corps, Hovaizeh WTP* *Corresponding Author, E-mail: <u>K naddafee@yahoo.com</u>

Introduction

During last decade, there is a growing concern about usable water resources decrease. Currently, the world is moving towards water crisis. Water shortage is an important concern in arid areas such as Africa, Southern Asia and Middle East and even in some parts of the World which it may lead to a war crisis (Jaafarzadeh, 1996). On the other hands continued population growth, increased per capital water consumption and increased water requirements for industry and irrigation result in considerable decrease of usable water resources. Therefore, treated wastewater recycling into the hydrological cycle is of significant importance and has many benefits. The major uses of treated wastewater are in agricultural irrigation, industrial activities and groundwater recharge. The amount of water used in the above-mentioned activities depends on the properties of water catchments, meteorological and gehydrological factors and direct and indirect water consumption rate. With respect to public health, principles of engineering economy, aesthetic standards and more importantly public acceptance, wastewater reuse can be developed.

Reuse of wastewater effluent is currently receiving more attention because of water shortages. Use of treated wastewater effluent can be a new source of water (Zadhoosh, 1996). Traditionally, domestic wastewater was used as fertilizer in agriculture or as soil conditioner in some parts of Asia. However, documents on application of effluent in agriculture were found in Bazlau, Germany in 1531 and in Edinburgh in Scotland. As wastewater treatment methods developed more affection was made on treated wastewater reuse in agriculture. Especially, some specialists believe that land treatment of effluent is the best alternative (Alizadeh and Naghibi, 1996). Moreover, infiltration of effluent into soil can cause environmental degradation as well as river pollution (Kaynezhad and Ebrahimi, 1997). Two main reasons that have firstly been encouraged wastewater reuse include: 1. River pollution can be avoided, and 2. Wastewater is an enrich source of nutrients for plants. After that water resource protection and effluent reuse in areas with water shortage have been taken into account. Scientific advances, economic consideration and more

importantly experiments gained up to now, both encourage and discourage propel to use effluent for irrigation although these reasons are still important (Blumental and Peasey, 2000).

At the present time, most areas in Iran are suffering from water shortage and most others will experience this problem. Thus, wastewater can be a reliable source of water and after being collected it will be an available water source (Metcalf and Eddy, 2003). Investigation of potential and importance of treated wastewater reuse will help society to compare the reclaimed water and water extracted from drinking water resources in large scale.

This experiment has been done at Hovaizeh area at Khuzistan Province of Iran on 2003.

Materials and Methods

A field study was carried out to investigate the safety and feasibility of crop irrigation, which were mostly eaten raw, by treated effluent of stabilization pond received municipal wastewater. The farms chosen as study field were located adjacent to Hovaizeh wastewater treatment plant system.

Two experimental plots, each sized 0.5 ha. were chosen to study the treated effluent effects on cross through statistical analysis. One of the farms vas irrigated using the stream that is divided from N_h. River. The flooded irrigation was applied ands. In order to protect the effluent percola. In in to the adjacent farm, insulation we. may be een the farm using nylon sheets. In addit. , t. , farms were fenced to prevent animal's intr. ion 'at might cause secondary pollution. In gen ral, both farms were similar in terms of size ype. . . crop, amount of fertilizers used and lack of llutants in soil; expecting that the water sour . red for irrigation was different to enable us to evalua the crops and chemical (especially heavy mctals) and microbial pollution that might be exist. The samples were taken from the inlet to each plot. Some examinations EC, pH and temperature were measure on-site and to examine the rest parameters DO, BOD, COD, TDS, alkalinity, acidity, chloride, salinity, sodium absorption ration (SAR), Nitrogen and sulfide compounds and metals including Ca, K, Mn, Na, Fe, the samples were preserved at 4 °C and sent to the Khuzestan Central Laboratory. All the examinations were followed Standard Methods (Metcalf and Eddy, 2003). One way ANOVA was used to determine statistically significance of the characteristics of the soils irrigated by either Nissan River or treated effluent.

Results

Table 1 provides the physical and chemical parameters of the treated effluent in the inlet of two plots and those of Nisan river water; microbial quality of each source is provided in Table 2. Heavy metal contents in the crops produced in each plot are given in Table 3. Parasitic quality of the water produced from the crops of each plot is presented in Table 4 and Table 5 gives the crops development and growth, in addition to their observed characteristics. According to Table 1 it can be found that the high salinity in Nisan river as well as soil poor quality influences the permeability of the ground. Water soluble salts increase the osmotic pressure of soil water which in the ist of absorb water.

Table 1: The overlage physical and chemical properies of the the ded effluent and Nisan river water

Parameter	Unit	River	Treated Effluent	
Ţ <u>,</u> C	µmohs/ cm	3905	2780	
Te verature	°C	22	24.2	
pH	-	7.7	7.67	
T.S.S.	mg/l	599	607	
T.D.S.	mg/l	2188	1892	
Total Hardness	mg/l as CaCO ₃	584.5	851	
Ca	mg/l	170	156	
K	mg/l	189	22	
Mg	mg/l	39	112	
Na	mg/l	145	309	
Acidity	mg/l as CaCO ₃	81	18	
Alkalinity	mg/l as CaCO ₃	655	157	
BOD ₅	mg/l	89	17	
COD	mg/l	289	102	
Sulfate	mg/l	420	731	
Nitrate	mg/l	1.38	4.94	
Nitrite	mg/l	0.066	0.14	
Chloride	mg/l	735	533	
Salinity	-	33.74	27.08	
SAR	-	0.912	0.75	

Table 2: The mean total coliform and fecal coliform inNisan river water and the treated effluent

Parameter	Unit	River	Treated Effluent
Total Coliform	MPN/100 ml	88.7×10^{5}	30×10^{6}
Fecal Coliform	MPN/100 ml	4.9×10^4	19×10^4

Therefore, respiration of plant increases and the plant growth will decline as a consequence. Accordingly, it can be underfeed that no growth occurred in most farms. The relative amounts of chemicals in the ground were almost the same amount found in the irrigation water. Increased application of chemical fertilizers affects on the concentration of soil chemical properties. Thus, both kinds of irrigation waters can negatively impact the soil. Heavy metal concentrations had not any adverse effect on crops. Examinations carried out on the edible parts of the parts showed no significant difference in heavy metal contents between their concentrations in plants irrigated with Nisan River water and those irrigated with the treated effluent. The samples taken from both Nisan rivers and the treated effluent indicated high coliform contents so that chlorination process for both irrigation sources should be improved to reduce the risks related to crops contamination by pathogens. Crop growth rate were significantly different according to the irrigation source and ground chemical quality.

The study showed that of overall 36 garden personal farms irrigated by Nisan river water, crop production growth in 7 farms (19.4%) was 20-60%, in 10 farms (27.7%) was 5-30%, and in 19 farms (25.7%) n growth was recorded, while of overall 28 farms irrigated by the treated effluent, growth in 6 farms (21.4%) was 10-100%, in 4 farms (14.2%) was 20-60%, in 4 farms (14.2%) was 5-30%, and in 14 farms (50%) no growth was observed.

Of overall 44 sweet basil farms irrigated by Nisan river water, growth in 21 farms (47.7%) was 30-60%, in 10 farms (22.7%) was 5-30% and in 13 farms (29.5%) no growth was observed, while the 24 farms irrigated by the treated effluent showed no growth and they were completely dry. Of overall

32 radishes farm irrigated by Nisan River water, growth in 10 farms (31.2%) was 30-60%, in 9 farms(28.1%) was 5-30% and in 13 farms (40.6%) no growth was occurred, while the 20 radish farms irrigated by the treated effluent showed no growth in plants.

In several rows of cucumber irrigated by Nisan river water no growth was observed, while of overall 15 cucumber farms irrigated by the treated effluent, growth in 5 farms (33.3%) was 30-50%, in 6 farms (40%) was 15-30% and in 4 farms (33.3%) was 5-15%.

Discussion and Conclusion

Gradually the uncommon water resources such as treated effluents has found an assured place in human consumption cycle, because with urban development and increased public health a lot effort has been put into supply water. Thus, a potential source of water in urban areas can be treated effluents that have continuously been increased.

From the above-mentioned results and the analyses carried out on the water and rivereffluent it can be concluded that low crop production may result from ground poor quality (high salinity). The existing high salinity of the studied area should have been improved before growing crops. With comparison of the results obtained from crops production in both plots, it is revealed that growth was quantitatively and qualitatively higher in the farms irrigated by the treated effluent than that of the farms irrigated by Nisan river water. Numerous works on this subject earlier. For instance, Alizadeh, et al., (1996) showed that vegetables (tomato, lettuce, cucumber and carrot) irrigation by treated effluent increased their development in comparison with irrigation by water.

Metal	Unit	Garden Purslane		Sweet Basil		Radish		Cucumber	
	River	Treated Effluent	River	Treated Effluent	River	Treated Effluent	River	Treated Effluent	
Ni	mg/l	85	270	50	-	45	250	90	85
Hg	mg/l	0.3	0.04	0.06	-	-	0.95	0.12	0.38
Cd	mg/l	30	30	30	-	40	50	20	30
Cr	mg/l	85	50	20	-	40	820	30	50

Table 3: The mean concentraition of heavy metal in Nisan river water and the treated effluent

Crop	Irrigation by River Water		Irrigation by Treated Effluent		
Crop	Sample	Sample 2	Sample	Sample 2	
Garden Purslane	-	-	-	-	
Sweet Basil	-	-	-	Nematode	
Radish		Live larvae	-	Live larvae	
Cucumber	-	-	-		

Table 4: Parasitic quality of the water generated from crops washing by River water and the treated effluent

Table 5: The crops growth irrigated by Nisan water
and the treated effluent

Crop	River	Treated Effluent
Garden Purslane	Growth in 7 Farms: 30-60% Growth in 10 Farms: 5-20% Growth in 19 Farms: no growth Overall Farms: 36	Growth in 6 Farms: 70- 100% Growth in 4 Farms: 30-60% Growth in 4 Farms: 30-50% Growth in 14 Farms: no growth Overall farms: 28
Sweet Basil	Growth 21 Farms: 30-60% Growth in 10 Farms: 30-50% Growth in 12 Farms: no growth Overall Farms: 34	No growth in any Farms Overall Farms: 34
Radish	Growth in 10 Farms: 30-60% Growth in 12 Farms: 5-30% Growth in 13 Farms: no growth Overall Farms: 32	No growth in any Farms Overall Farms: 20
Cucumber	No growth in any farm	Growth in 5 rows: 30-50% Growth in 6 rows: 15-30% Growth in 4 rows: 5-15% Overall rows: 15

They stated that as treated effluent is a rich source of nutrients needed by plants, it would increase plant growth 60 that using treated effluent for irrigation of the studied plants can be equal to 25 tone application of manure on land (Zadhoosh, 1996). Zadhoosh (1996) stated that using raw wastewater for wheat farm irrigation increased grain and wheat weight in comparison with irrigation increased grain and wheat weight in comparison with irrigation by water (Abedi and Najafi, 2001). Apparently, wastewater reuse has some disadvantages. For instance, wastewater disposal into seasonal Shiraz river to be reused can result in adjacent soil pollution and transmission of some heavy metals to plants in long term (AWWA, 1998). In general, care should be exercised for irrigation of raw edible vegetables by wastewater effluent. People may be reluctant to use the vegetables irrigated by wastewater effluent, although they do not have any health effect. Accordingly, most consultant engineers do not recommend using wastewater effluent for plants irrigation such as vegetables. However, there is no scientific evidence to encourage or discourage wastewater effluent for irrigation use.

References

- Abedi, M. J. and F. Najafi, Application of treated domestic effluent in agriculture. Publication of Iranian national irrigation and drainage committee. 1^{st.} Ed., Tehran, Iran, 248, 2001
- Alizadeh, A. and A. Naghibi, Use of treated domestic wastewater in irrigation. Proceeding of the second national water and soil issues congress, Tehran, Iran, 369, 1996
- AWWA, WEF and APHA, Standard methods for the examination of water and wastewater, 14th. Ed., 1998
- Blumental, U. and A. Peasey, Guidelines for wastewater reuse in agriculture and aquaculture. Ult Ed. Loughborough University, 2000
- Jaafarzadeh, N., Effect of wastewater of Shiraz city for crops irrigation on heavy metal increase in soil and plants. Proceedings of the second national water and soil congress, Published by agricultural education and development organization, Tehran, Iran, 369, 1996
- Kaynezhad, M. A. and S. Ebrahimi, Environmental engineering. Sahand university publication, 1^{st.} Ed., Tabriz, Iran, 545, 1997

- Metcalf and Eddy, Wastewater engineering: Treatment and Reuse. 4th. Ed., McGraw Hill Publication, 2003
- *Rowe, D. and I. Abdelmagid, Handbook of wastewater reclamation and reuse.* 2^{nd.} Ed., Lews publisher, CRC *Press,* 420, 1995
- Zadhoosh., Wastewater effluent irrigation effects on soil and plants M.S. thesis, Tehran university, Tehran, Iran, 1996