

Characterization of bioflocculants produced by bacteria isolated from crude petroleum oil

¹S. Zaki; ¹S. Farag; ¹G. Abu Elreesh; ²M. Elkady; ¹M. Nosier; ¹*D. Abd El Haleem

¹Department of Environmental Biotechnology, Genetic Engineering and Biotechnology Research Institute, City for Scientific Research and Technology Applications, Burgelarab, Alexandria, Egypt

²Department of Fabrication Technology, Advanced Technology and New Materials Research Institute, City for Scientific Research and Technology Applications, Alexandria, Egypt

Received 21 October 2010; revised 25 April 2011; accepted 9 August 2011; available online 1 September 2011

ABSTRACT: The aim of this study was to look for high efficient bioflocculant-producing microorganisms. Among 36 bacterial colonies isolated from a crude petroleum oil sample, three of them as *Bacillus subtilis* and *Pseudomonas* spp. exhibited flocculation activity exceeding 90 % after 3 days of cultivation. They were identified by 16 S rDNA sequence analysis as *Bacillus subtilis* and *Pseudomonas* sp. Spectroscopic analysis of the polymers by nuclear magnetic resonance and fourier-transform infrared revealed that the polymers were glycoproteins. These polymers were soluble in water and insoluble in any organic solvents tested. The effects of bioflocculant dosage, temperature and pH on the flocculation activity were evaluated. The maximum bioflocculation activities were observed at an optimum bioflocculant dosage of 3.5 mg/L (strains *Bacillus subtilis* and *Pseudomonas*) and 5.0 mg/L (strain CPO14), respectively. In addition, these biopolymers were able to flocculate kaolin suspension (5 g/L) over a wide range of pH (pH 3–9) and temperature (5–50 °C) tested in the presence of CaCl₂. The highest flocculation activities of strains CPO8, CPO13 and CPO14 were 96.03 %, 92.17 % and 97.59 %, respectively in the early stationary phase (at 24 h), while the cell production reached its maximum in the stationary phase (at 72 h). Their efficient flocculation capabilities suggest potential applications in industries.

Keywords: Bioflocculant; Biopolymer; Flocculation activity; Optimization

INTRODUCTION

Industrial process is a potential source of pollution and requires a specific treatment for the waste produced. Introduction of waste treatment increases the plant cost, therefore an attempt to improve its efficiency is valuable. This applies to all kinds of industries but is particularly important for low or medium added value industries with difficult wastes like tanneries and clay processing industry. Flocculation is the most widely used process for treating the wastewater in these kinds of industries (Rossini *et al.*, 1999). Separation by flocculation is effective and sufficient when the end objective is reduction in sludge volume and rapid separation of solids from liquids (Besra *et al.*, 2003).

The problem of removal of kaolin turbidity from the effluents before discharging into the environment was studied by many researchers. Faust and Aly (1983)

quote extensive application of kaolin suspensions as a model for natural turbidity of raw surface water in Jar Test studies of coagulation and flocculation in the laboratory. Inorganic coagulants such as alum in combination with lime have been conventionally used for the removal of clay particles in the effluents from clay processing industry as well as from raw water. The sludge formed from such treatment poses disposal problems because of its aluminium content and tend to accumulate in the environment (Faust and Aly, 1983). Stauber *et al.* (1999) and Divakaran and Sivasankara, (2001) have reported that although not conclusively proved, increasing concern about the residual aluminium, which may be present in water as a result of alum treatment, is being expressed by the public in connection with Alzheimer's disease. Thus, it has become necessary to develop harmless, more efficient and environment friendly flocculants for removal of turbidity in surface waters and effluents.

*Corresponding Author Email: abdelhaleemm@yahoo.de
Tel.: +2011 586 6113; Fax: +2034 593 420



Biofloculants have received considerable scientific and biotechnological attention in recent years due to their biodegradability, benign nature and lack of secondary pollution of their degradative intermediates (Salehizadeh and Shojaosadati, 2001). They reported that biofloculants are microorganism-produced special natural organic macromolecule substances that can flocculate suspended solids, cells, colloidal solids, etc. In recent years, several kinds of microorganisms, which secrete flocculation biopolymer, have been screened and isolated from activated sludge, soil and wastewater. The species include bacteria, fungi, actinomyces and algae. Generally, soil and activated sludge samples are the major sources for isolating biofloculant-producing microorganisms (Abd-El-Haleem et al., 2008). However, isolation and characterization of biofloculant-producing microorganisms from crude petroleum oil samples were not well documented in literatures. Therefore, present study aims to look for biofloculant-producing microorganisms in crude petroleum oil sample collected on 28 May 2009 from a sews-gulf oil drilling field in Egypt. Screening, isolation and characterizations of both biofloculant-producing microorganisms and their extracellular biofloculants were performed.

MATERIALS AND METHODS

Media and cultivation conditions

Isolation medium

The Sup-plemented nutrient broth (SNB) medium used to isolate bacteria from the oil sample (collected from a sews-gulf oil drilling field in Egypt) was containing per liter of 500 mL of Nutrient Broth and 500 mL mineral salt solution. The 500 mL of salt solution was prepared by adding the various salts such as KH_2PO_4 , 20 g; K_2HPO_4 , 5.0 g, $(\text{NH}_4)_2\text{SO}_4$, 30 g; NaCl , 0.1 g; $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.01 g; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g; $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 0.01 g; $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g; $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 0.01 g; and $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 0.002 g, glucose, 0.03 % (w/v); and yeast extract, 0.03 % (w/v) (Francy et al., 1991). One-half SNB medium with 2 % Bacto Agar (SNA) was used to isolate heterotrophs.

Prescreening medium

The prescreening medium used to select for biofloculant-producing bacteria contained (per liter) 1% glucose, 0.35% yeast extract, 0.5 % K_2HPO_4 , 0.2 % KH_2PO_4 , 0.05 % $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.01% NaCl and agar 1.5 %. The broth of the prescreening medium was used as a seeding medium.

Fermentation medium

The fermentation medium was containing per liter; glucose 10 g, NH_4Cl 7 g, K_2HPO_4 0.5 g, MgSO_4 0.5 g, FeCl_3 40 mg, CaCl_2 150 mg and MnSO_4 140 mg.

Cultivation conditions

The initial pH of all media was adjusted to 7.2 to 7.5 with NaOH (1 M) and HCl (0.5 M). All the media were prepared with distilled water and sterilized at 121 °C for 20 min. All cultivations were done at 30 °C.

Isolation, screening and identification of biofloculant-producing bacteria

Screening process for biofloculant-producing bacteria was performed in two experimental steps. First, all bacteria that might exist in the sample were isolated by adding 1 mL of oil sample to 99 mL of SNB medium in 250 mL flasks. The mixture was placed on a rotary shaker at 200 rpm for 2 h at 30 °C to produce a well-dispersed suspension. Subsequently, the suspension was diluted serially in SNB broth and plated in triplicate on one-half SNB with 2% Bacto Agar to isolate heterotrophs. The plates were incubated at 30 °C. Pure cultures of bacteria were isolated by cycles of re-plating on SNB agar plates. Thirty six pure bacteria with different colony morphologies were isolated and maintained on Nutrient Agar slants until used.

Secondly, to screen among all isolates for biofloculant-producing bacteria, a loop of each isolate was suspended in sterile 1 mL distilled water and an aliquot ~ 100 μL was spread on the prescreening medium and incubated at 30 °C for 48 h (Dermlim et al., 1999; Abd-El-Haleem et al., 2008). Microorganisms were screened for their abilities to produce biofloculant based on colony morphology (mucoid and ropy) (Murray et al., 1994). The positive strains were grown in 50 mL fermentation medium in 250 mL flasks on a rotary shaker at 30 °C for 3 days. At the end of cultivation, the culture broths were measured for flocculation activities (described below).

The strains with the highest flocculating activities (CPO8, CPO13 and CPO14) were selected and identified based on 16 S rDNA sequencing. Their isolated genomic DNA was subjected to PCR amplification of 16 S rDNA using universal primers. Amplified products were gel purified using Qiagen gel extraction kit and purified products were sequenced (Kim et al., 2006). Subsequently, comparative sequence analysis of 16 S

rDNA amplified genes was performed and a phylogenetic tree was generated according to the procedure described previously by Al-Thani *et al.* (2009).

Determination of flocculation activity

Flocculation activities were calculated and measured using a modified Kaolin clay suspension method (Kurane *et al.*, 1986). A mixture of 5 g/L synthetic clay suspension (Sigma–Aldrich, 0.1–4 µm particle size) with 0.5 mL bioflocculant solution (culture broth, cell-free supernatant or extracted biopolymer) in the presence of CaCl₂ (as coagulant) was stirred with rapid mixing at 230 rpm for 2 min, followed by slow mixing at 80 rpm for 3 min using Laboratory Flocculator (Jar Tester Model CZ150) and left standing for 5 min. A sample for Optical densities (OD) measurement was withdrawn using automatic pipette from a height of 3 cm below the surface of clay suspension. Relying on the upper phase OD for clay suspension that was measured at 550 nm with a spectrophotometer (7230G, Shanghai, China) the flocculation efficiency of the different isolated strains was screened. A control was prepared using the same method but the sample was replaced by distilled water. The flocculation activity was calculated according to the equation:

$$\text{Flocculation activity (\%)} = \left[\frac{A - B}{A} \right] \times 100$$

Where A and B are the supernatant OD of the control (clay suspension without any bioflocculant addition) and sample respectively, at 550 nm.

Time course of bioflocculant production

Time courses of the selected strains were carried out by cultivating in fermentation medium with the addition of 5 % seeding culture (24 h and viable cell count of 10⁸ CFU/mL) at 30 °C on a rotary shaker at 200 rpm for 7 days. Samples were taken every 12 h to measure, optical density (OD 600 nm) and flocculation activity (as described above).

Partial purification of the bioflocculants

To produce bioflocculants, the strains were pre-cultured in the seed medium as described above and inoculated into fermentation medium with the addition of 2 % starter culture (24 h) on a rotary shaker (200 rpm) at 30 °C for another 24 h. To partially purify bioflocculants, culture broths were centrifuged to remove cells by centrifugal separation (4000 g, 30 min).

Subsequently, cell-free supernatants were freeze-dried. The lyophilized precipitates were redissolved in an appropriate amount of distilled water then re-precipitated with four volumes of cold 95 % ethanol and left for four hours at 4 °C. After centrifugation at 7500 g, 4 °C for 20 min, above steps were repeated for two additional times and the final precipitates were freeze dried.

Characterization of the partially purified biopolymers

Infra-red spectra (IR) of the biopolymers were measured on a KBr disk with a Perkin-Elmer series 1600 FT-IR to determine the functional groups of the biopolymers. The ¹H NMR spectra of the bioflocculants were recorded on a 500 MHz Bruker a JOEL GSX 500 spectrometer (500 MHz). The bioflocculants were dissolved in 2H₂O. The protein concentrations of the biopolymers were determined according to Bradford method (Bradford, 1976). Total sugars were determined by the phenol-sulfuric acid reaction using the procedure of Chaplin and Kennedy (1994). Additional analysis of the biopolymers was performed, which included the solubility test in distilled water and several solvents such as acetone, carbon tetrachloride, ethanol, isopropanol, hexane, methanol and nitrobenzene (Collins *et al.*, 1973). The effects of polymer concentrations (final concentration of 0–7 mg/L) on the flocculation activity were studied. Subsequently, the effects of pH (3–12) and temperature (5 - 100 °C) on the flocculation activity of the partially purified biopolymers were studied by measuring the flocculation activity of the reaction mixture containing the optimum concentration of biopolymers at the specified ranges of pH and incubated at different temperatures (Kwon *et al.*, 1996).

RESULTS AND DISCUSSION

Screening and identification of bioflocculant-producing bacteria

Despite, so many studies about bioflocculants have been done; flocculation activity and culture cost of bioflocculants are still the major limiting factors with regard to their application (Zhang *et al.*, 2007). Consequently, there is a need to identify new microorganisms especially from unusual environments (e.g. crude oil) with high bioflocculant-producing ability and improve upon the flocculation efficiency of the known bioflocculants. It is known

that crude oil or oil-contaminated sites are the main sources of biosurfactant-producing microorganisms (Yin *et al.*, 2009). However, no previous report exists on the isolation and characterization of biofloculant-producing microorganisms from crude petroleum oil. In the present study, a total of 36 colonies were isolated from crude petroleum oil sample, and 10 pure culture strains with slimy or mucoid appearance were screened on the basis of kaolin suspension flocculation activity over 85 % (data not shown). Among them, three strains CPO8, CPO13 and CPO14 having kaolin at flocculation activity exceeding 90 % were selected for further studies.

According to their 16 S rDNA sequence analysis strains CPO8 and CPO13 were belonging to the species of *Bacillus subtilis*. While, strain CPO14 was identified as *Pseudomonas* sp., respectively. Their 16 S rDNA sequences have been deposited in the GenBank database with accession numbers HQ015371, HQ015373 and HQ015374. Phylogenetic tree based on 16 S rDNA

sequences was constructed using the neighbor-joining method (Fig.1). Out of these results, two among three selected biofloculant-producing bacteria are belonging to the genus *Bacillus*. This result was in the same direction with many previous studies which reported that several bacterial strains closely related to the genus *Bacillus* able to produce biofloculants (Shih *et al.*, 2001; Deng *et al.*, 2003; Abd-El-Haleem *et al.*, 2008; Xiong *et al.*, 2010).

The effect of the time course on both growth and biofloculants production of the three strains are given in Fig. 2a and 2b. The OD curve that may be represent the cell growth shows a sharp increment in the first 72 h then tends to be constant over the remaining studied culturing period. However, the flocculation activity during this period has been increased and reached its maximum value of 96.03 % (COP8), 92.17 % (CPO13) and 97.59 % (CPO14), respectively at 24 h, then decreased slightly until 132 h. Concerning both the biofloculant production and the cell growth profiles,




Fig. 1: Phylogenetic tree showing the relationships among the selected isolates (in boldface) and other closely related sequences collected from the Gene Bank. The dendrogram was generated by the neighbor-joining method. Bootstrap values per 100 bootstrap analysis presented for values greater than 52 %. The microorganism *Bacillus subtilis* FJ 392726 was used as an out-group




Fig. 2: Time course of growth (a) and bioflocculation activity (b) of strains CPO8, CPO13 and CPO14 in fermentation medium

it is evident that the bioflocculant production was almost in parallel with cell growth till 132 h studied (Liu *et al.*, 2010). Where the produced bioflocculant reached its maximum flocculating efficiency in the early stationary phase (at 24 h), while the cell production reached its maximum in the stationary phase (at 72 h), indicating that the bioflocculant was produced by biosynthesis during its growth, not by cell autolysis (Lu *et al.*, 2005). However, as the incubation time increased, the flocculating rate started to decrease in spite of the OD still recorded high values that may be related to the turbidity of the produced bioflocculant. This result indicated that all tested strains produce extracellular bioflocculants. Accordingly, in order to produce bioflocculant with high flocculating efficiency, 24 h culturing time has been chosen. Previously, the maximum flocculant production of *Alcaligenes latus* was achieved at the middle and late stage of the logarithmic growth phase (2–3 days), and flocculation activity began to decrease during the late stationary phase due to the activity of deflocculation enzymes (Kurane and Nohata, 1991). The growth curve of *P. mirabilis* TJ1 showed that the production of its bioflocculant was almost parallel with cell growth. The bioflocculant of *P. mirabilis* TJ1 reached its maximum flocculation rate (93.13 %) at stationary phase (Xia *et al.*, 2008). In the same way, production of the bioflocculant by *Enterobacter aerogenes* was in parallel with cell growth and reached its maximum flocculation rate in early stationary phase (Lu *et al.*, 2005). The flocculation rate of the bioflocculant from *Serratia ficaria* and *Bacillus* sp. F19 reached its

maximum (94.3 % and 97 %, respectively) in early stationary phase (Gong *et al.*, 2008).

Characterization of bioflocculants

The phenol-sulfuric acid method showed that the partially purified biopolymers contained 74 %, 40 % and 47 % of total sugars, while protein contents detected by the Bradford method were 13 %, 17 % and 14 % for the strains CPO8, CPO13, and CPO14, respectively, indicating the composition of all bioflocculants are glycoproteins. The Fourier-transform Infra-red (FTIR) spectra (Fig. 3 a, b and c) of the three bioflocculants were analyzed according to Xiong *et al.* (2010). It clearly showed the presence of carboxyl, hydroxyl, amide and amino groups in its molecules, which are the preferred groups for the flocculation, similar to those observed in polyelectrolytes (Ganesh-Kumar *et al.*, 2004). The spectra showed a broad stretching intense peak displayed around 3440/cm that was characteristic of hydroxyl and amino groups (Liu *et al.*, 2009). The weak stretching band at ~ 2955/cm known to be typical of carbohydrates, indicated COH asymmetrical stretching vibration. An asymmetrical stretching peak observed at ~1,640/cm was characteristic of CAO stretching vibration in ONHCOCH₃ (Xiong *et al.*, 2010). The weak band near 1405/cm indicates the bending vibration of CH₃ and the scissor vibration of CH₂. The weak peak at 1,261/cm in the bioflocculants produced by the *Bacillus* strains CPO8 and CPO13 may be assigned to be CAO symmetrical and asymmetrical stretching of a carboxylate group in the bioflocculant. However, this

peak was absent in the biofloculant produced by strain CPO14. In addition, the strong absorption band at the range of 1049 - 1097/cm indicated asymmetrical stretching vibration of a C-O-C ester linkage. These infrared spectra showed characteristic peaks for carbohydrates and amides. Therefore, it can be inferred that the biofloculant is a types of glycoproteins.

To confirm FTIR results, Proton nuclear magnetic resonance (¹H NMR) spectra analysis of the three biofloculants were performed. As shown in Fig. 4 a, b and c, the first notable result of the ¹H NMR spectra was that the spectra were virtually identical. In addition, the presence of carbohydrates is recognized by the intense signals in the characteristic bulk region at 3.4-

4.3 ppm. In the region 4.8-5.2 ppm severe overlap of the carbohydrate anomeric signals with those stemming from C-protons of the protein backbone is observed. Previously, de-Beer *et al.* (1994) reported that most cross peaks stemming from carbohydrates attached to an intact glycoprotein appear in a region (3.4-5.2 ppm) in the spectrum essentially devoid of cross peaks of the protein backbone. However, the strong water (HOD) signal at 4.8 ppm obscures the beta-anomeric proton signal at about 4.75 ppm. The signals for the methyl protons of the 6-deoxy sugars are seen at about at 1.44–1.48 ppm (Kumar *et al.*, 2004). The results obtained from the NMR analysis confirmed the FTIR results and indicated that all of selected bacterial strains produced




Fig. 3: FT-IR spectra of the partially purified biopolymers from strains CPO8 (a), CPO13 (b) and CPO14 (c)




Fig. 4: ¹H NMR spectra of the partially purified polymers from strains CPO8 (a), CPO13 (b) and CPO14 (c)

glycoproteins. Out of these results, the protein and the carbohydrate contents of the purified biopolymers may explain their stability toward all studied organic solvents (acetone, methanol, ethanol, chloroform, dimethylsulphoxide). However, it has a tendency to be soluble in water as well as both acidic and alkaline media.

Effect of biofloculants concentration pH and temperature on bioflocculation activity

Fig. 5 shows the relationship between the flocculants dosages and flocculation activities. When the culture broth in kaolin suspension (5.0 g/L) was tested in the dosage range of 0.5- 39 mg/L, it was apparent that the flocculation activity increased proportionally to the flocculant dosage of 0.5 to 7.0 mg/L and was highest at 3.5 mg/L of strains CPO8 and CPO13 and at 5.0 mg/L of strain CPO14, respectively. These results could be clarified as follows: a) The incomplete dispersion of excess polysaccharide, only the kaolin particles around the polysaccharides participated in the flocculation reaction, therefore, other kaolin particles did not participate in the reaction (Yokoi *et al.*, 1997) and b) The excess polysaccharide was oversaturated on many binding sites of the surface of kaolin particles, thus the attractive force of the other particles was reduced and the flocculation activity decreased (Kwon *et al.*, 1996). Thus, either the deficiency or excess amount of polysaccharide and kaolin clay decreased or even prevented the flocculation activity (Lee *et al.*, 1995).

The partially purified biofloculants had the optimum pH for flocculation activity at pH 7 and were not

affected by the pH range of 3–9 (Fig. 6a). At higher pH of 9–11, the flocculation activity decreased gradually with all strains except of strain CPO13 which still has high flocculation activity (96.4 %) up to pH 11. The pH range for flocculation reaction of these biopolymers were wider than those of the polyglutamate from *Bacillus subtilis* PY-90 (pH range of 3–5) (Yokoi *et al.*, 1995) and the cationic polysaccharide from *Paecilomyces sp.* I-1 (pH range of 4–8) (Takagi and Kadowaki, 1985). It is demonstrated that the biopolymer solution is suitable to be applied in neutral, weakly acid and weakly alkaline circumstances. However, its flocculation efficiency slightly reduces at highly acidic (pH<5) and highly alkaline (pH>9) circumstances. This may be returned to the biopolymer shows different electric states at different pH, in turn affects the bridging efficiency of the biopolymer for clay powder (Yong *et al.*, 2009).

However, the flocculation activity of the biofloculants produced by strains CPO8 and CPO13 was not affected by increasing the reaction temperatures in the range of 5–70 °C (Fig. 6b). While, the flocculation activity of strain CPO14 was decreased gradually by increasing the reaction temperature above 50 °C. It is elucidated that flocculation efficiency increments as the clay suspension temperature increased from 5-70 °C, but decreased as the clay suspension temperature increased above 70 °C. This result indicates that the too high clay suspension temperature is unfavorable for the flocculation performance of the biopolymer. This can be explained by chemical kinetics. Both the rate of biopolymer




Fig. 5: The effect of dosage on the flocculation activity of the partially purified biopolymers from strains CPO8, CPO13 and CPO14




Fig. 6: The influence of initial pH (a) and temperature (b) of Kaolin clay suspensions on the flocculation activity of the partially purified biopolymers from strains CPO8, CPO13 and CPO14

diffusion and suspended particle collision frequency were improved at higher temperatures, this contributes to the increase in rate of reaction (Lu *et al.*, 2005; Gong *et al.*, 2008). The flocculating activity of strain CPO14 that decreased significantly when the temperature exceeded 50 °C. This can be explained by denaturalization of proteins in the biofloculant and an increase in hot movement of kaolin particles (Liu *et al.*, 2009).

CONCLUSION

Three biofloculant-producing strains were isolated from contaminated crude petroleum oil. Two of them CPO8 and CPO13 are belonging to the genus *Bacillus*,

while strain CPO14 identified as *Pseudomonas*. Chemical analysis showed that the main compositions of their partially purified biofloculants were carbohydrates containing some proteins. FTIR indicated the presence of carboxyl, hydroxyl, and methoxyl groups. ¹HNMR analysis confirmed that the extracted biofloculants were glycoproteins. The highest flocculation activities of strains CPO8, CPO13 and CPO14 were 96.03 %, 92.17 % and 97.59 %, respectively at 24 h of cultivation. The thermal and pH stabilities of the biopolymer solution indicated its thermal stability over the temperature range of (5-100 °C). Moreover, the biopolymer solution was suitable to be applied in neutral, weakly acid and

weakly alkaline circumstances. The advantageous properties of these bioflocculants, such as stable thermal and pH characteristics and dosing rate comparable to those of chemical flocculants to flocculate suspended solids, suggest its potential industrial utility.

ACKNOWLEDGEMENTS

This work was supported by a grant from the Egyptian Science and Technology Development Fund (STDF) (Grant No. 743). The study was conducted at Environmental Biotechnology Department, Genetic Engineering and biotechnology Institute, City for Scientific Research and Technology Applications, Alexandria, Egypt.

REFERENCES

- Abd-El-Haleem, D.; Al-Thani, R.; Al-Mokemy, T.; Al-Marii, S.; Hassan, F., (2008). Isolation and characterization of extracellular bioflocculants produced by bacteria isolated from Qatari Ecosystems. *Pol. J. Microbiol.*, 57 (3), 231–239 (9 pages).
- Al-Thani, R.; Abd-El-Haleem, D.; Al-Shammri, M., (2009). Isolation and characterization of polyaromatic hydrocarbons-degrading bacteria from different qatari soils. *Afr. J. Microbiol. Res.*, 3 (11), 761-766 (6 pages).
- Besra, L.; Sengupta, D. K.; Roy, S. K.; Ay, P., (2003). Influence of surfactants on flocculation and dewatering of kaolin suspensions by cationic polyacrylamide (PAM-C) flocculant. *Sep. Purif. Tech.*, 30 (3), 251-264 (14 pages).
- Bradford, M. M., (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of proteindye binding. *Analys. Biochem.*, 72, 248–254 (5 pages).
- Chaplin, M. F.; Kennedy, J. F., (1994). *Carbohydrate Analysis*, seconded. Oxford University Press, New York, pp. 138 - 141 (4 pages).
- Collins, E. A.; Bares, J.; Billmeyer, F. W., (1973). *Experiments in Polymer Science*. New York: Wiley, pp. 125-130 (6 pages).
- De-Beer, T.; van-Zuylen, C. W. E. M.; HaË, K.; Boelens, R.; Kaptein, R.; Kamerling, J. P.; Vliegthart, J. F. G., (1994). Rapid and simple approach for the NMR resonance assignment of the carbohydrate chains of an intact glycoprotein. *FEBS Lett.*, 348 (1), 1-6 (6 pages).
- Deng, S. B.; Bai, R. B.; Hu, X. M.; Luo, Q., (2003). Characteristics of a bioflocculant produced by *Bacillus mucilaginosus* and its use in starch wastewater treatment. *Appl. Microbiol. Biotech.*, 60 (5), 588–593 (6 pages).
- Dermlim, W.; Prasertsan, P.; Doelle, H., (1999). Screening and characterization of bioflocculant produced by isolated *Klebsiella* sp. *Appl. Microbiol. Biotech.*, 52 (5), 698-703 (6 pages).
- Divakaran, R.; Sivasankara, V. N., (2001). Flocculation of kaolinite suspensions in water by chitosan. *Water Res.*, 35 (16), 3904–3908 (5 pages).
- Faust, S. D.; Aly, O. M., (1983). Removal of particulate matter by coagulation. In *Chemistry of Water Treatment*, Butterworth Publishers, Boston, pp. 277–367 (91 pages).
- Francy, D. S.; Thomas, J. M.; Raymond, R. L.; Ward, C. H., (1991). Emulsification of hydrocarbons by subsurface bacteria. *J. Ind. Microbiol.*, 8 (4), 237–246 (10 pages).
- Ganesh Kumar, C.; Joo, H. S.; Choi, J. W.; Koo, Y. M.; Chang, C. S., (2004). Purification and characterization of an extracellular polysaccharide from haloalkalophilic *Bacillus* sp. I-450. *Enzyme Microb. Tech.*, 34 (7), 673–681 (9 pages).
- Gong, W. X.; Wang, S. G.; Sun, X. F.; Liu, X. W.; Yue, Q. Y.; Gao, B. Y., (2008). Bioflocculant production by culture of *Serratia ficaria* and its application in wastewater treatment. *Bioresour. Tech.*, 99 (11), 4668–4674 (7 pages).
- Kim, L. S.; Hong, S. J.; Son, M. K.; Lee, Y. H., (2006). Polymeric and compositional properties of novel extracellular microbial polyglucosamine biopolymer from new strain of *Citrobacter* sp. BL-4. *Biotech. Lett.*, 28 (4), 241–245. (5 pages)
- Kumar, C. G.; Joo, H. S.; Choi, J. W., (2004). Purification and characterization of an extracellular polysaccharide from haloalkalophilic *Bacillus* sp. I-450[J]. *Enzyme Microb. Tech.*, 34 (7), 673–681 (9 pages).
- Kurane, R.; Nohata, Y., (1991). Microbial flocculation of waste liquids and oil emulsions by a bioflocculant from *Alcaligenes latus*. *Agr. Biol. Chem.*, 55 (4), 1127–1129. (3 pages)
- Kurane, R.; Takeda, K.; Suzuki, T., (1986). Screening for characteristics of microbial flocculants. *Agr. Biol. Chem.*, 50, 2301-2307 (7 pages).
- Kwon, G. S.; Moon, S. H.; Hong, S. D.; Lee, H. M.; Kim, H. S.; OH, H. M., (1996). A novel flocculant biopolymer produced by *Pestalotiopsis* sp. KCTC 8637P. *Biotech. Lett.*, 18 (12), 1459–1464 (6 pages).
- Lee, S. H.; Lee, S. O.; Jang, K. L.; Lee, T. H., (1995). Microbial flocculant from *Arcuadendron* sp. TS-49. *Biotech. Lett.*, 17 (1), 95–100 (6 pages).
- Liu, W.; Wang, K.; Li, B.; Yuan, H.; Yang, J., (2010). Production and characterization of an intracellular bioflocculant by *Chryseobacterium daeguense* W6 cultured in low nutrition medium. *Biores. Tech.*, 101 (3), 1044–1048 (5 pages).
- Liu, W.; Yuan, H.; Yang, J.; Li, B., (2009). Characterization of bioflocculants from biologically aerated filter backwashed sludge and its application in dyeing wastewater treatment. *Biores. Tech.*, 100 (9), 2629–2632 (4 pages).
- Lu, W.; Zhang, T.; Zhang, D.; Li, C.; Wen, J.; Du, L., (2005). A bioflocculant produced by *Enterobacter aerogenes* and its use in defecating the trona suspension. *Biochem. Eng. J.*, 27 (1), 1–7 (7 pages).
- Murray, E.; Doetsch R. N.; Robinow, C. F., (1994). Determinative and cytological light microscopy. In: Gerhardt P, Murray RGE, Wood WA, Krieg NR (eds) *Methods for general and molecular bacteriology*. American Society for Microbiology, Washington, DC, pp 21-41(21 pages).
- Rossini, M.; Garcia, J.; Galluzzo, M., (1999). Optimization of the coagulation-flocculation treatment: influence of rapid mix parameters. *Water Res.*, 33 (8), 1817-1826 (10 pages).
- Salehizadeh, H.; Shojaosadati, S. A., (2001). Extracellular biopolymeric flocculants: Recent trends and biotechnological importance. *Biotech. Adv.*, 19 (5), 371–385 (15 pages).
- Shih, I. L.; Van, Y. T.; Yeh, L. C., (2001). Production of a biopolymer flocculant from *Bacillus licheniformis* and its

- flocculation properties. *Bioresour. Tech.*, 78 (3), 267-272 (6 pages).
- Stauber, J. L.; Florence, T. M.; Davies, C. M.; Adams, M. S.; Buchanan, S. J., (1999). Bioavailability of Al in alum treated drinking water. *J. Am. Water Works Assoc.*, 91 (11), 84-93 (10 pages).
- Takagi, H.; Kadowaki, K., (1985). Flocculant production by *Paecilomyces* sp.: taxonomic studies and culture condition for production. *Agr. Biol. Chem.*, 49, 3151-3157 (7 pages).
- Xia, S. Q.; Zhang, Z. Q.; Wang, X. J.; Yang, A.; Chen, L.; Zhao, J. F.; Leonard, D.; Jaffrezic-Renault, N., (2008). Production and characterization of a bioflocculant by *Proteus mirabilis* TJ-1. *Bioresour. Tech.*, 99 (14), 6520-6527 (8 pages).
- Xiong, Y.; Wang, Y.; Yu, Y.; Li, Q.; Wang, H.; Chen, R.; He, N., (2010). Production and characterization of a novel bioflocculant from *Bacillus licheniformis*. *Appl. Environ. Microbiol.*, 76 (9), 2778-2782 (5 pages).
- Yin, H.; Qiang, J.; Jia, Y.; Ye, J.; Peng, H.; Qin, H.; Zhang, N.; He, B., (2009). Characteristics of biosurfactant produced by *Pseudomonas aeruginosa* S6 isolated from oil-containing wastewater. *Process Biochem.*, 44 (3), 302-308 (7 pages).
- Yokoi, H.; Natsuda, O.; Hirose, J.; Hayashi, S.; Takasaka, Y., (1995). Characteristics of a biopolymer flocculant produced by *Bacillus* sp. PY-90. *J. Ferment. Bioeng.*, 79 (4), 378-380 (3 pages).
- Yokoi, H.; Yoshida, T.; Mori, S.; Hirose, J.; Hayashi, S.; Takasaki, Y., (1997). Biopolymer flocculant produced by an *Enterobacter* sp. *Biotech. Lett.*, 19 (6), 569-573 (5 pages).
- Yong, P.; Bo, S.; Yu, Z., (2009). Research on flocculation property of bioflocculant PG.a21. *Ca. Mod. Appl. Sci.*, 3 (6), 106-112 (7 pages).
- Zhang, Z. Q.; Lin, B.; Xia, S. Q.; Wang, X. J.; Yang, A. M., (2007). Production and application of a novel bioflocculant by multiple-microorganism consortia using brewery wastewater as carbon source. *J. Environ. Sci. China.*, 19 (6), 667-673 (7 pages).

AUTHOR (S) BIOSKETCHES

Zaki, S., Ph.D., Associate Professor, Department of Environmental Biotechnology, Genetic Engineering and Biotechnology Research Institute, City for Scientific Research and Technology Applications, Burgelarab, Alexandria, Egypt, Email: saharzaki@yahoo.com

Farag, S., Ph.D., Assistant Professor, Department of Environmental Biotechnology, Genetic Engineering and Biotechnology Research Institute, City for Scientific Research and Technology Applications, Burgelarab, Alexandria, Egypt, Email: sohafarag88@yahoo.com

Abu Elreesh, G., Ph.D., Assistant Professor, Environmental Biotechnology Department, Genetic Engineering and Biotechnology Research Institute, City for Scientific Research and Technology Applications, Burgelarab, Alexandria, Egypt. Email: g_abelreesh@yahoo.com

Elkady, M., Ph.D., Assistant Professor, Department of Fabrication Technology, Advanced Technology and New Materials Research Institute (ATNMRI), Mubarak City for Scientific Research and Technology Applications, Alexandria, Egypt. Email: maroelkady@yahoo.com

Nosier, M., Ph.D. candidate, Department of Environmental Biotechnology, Genetic Engineering and Biotechnology Research Institute, City for Scientific Research and Technology Applications, Burgelarab City, Alexandria, Egypt. Email: mnosier1987@yahoo.com

Abd El Haleem, D., Ph.D., Professor and Head of the Department of Environmental Biotechnology, Genetic Engineering and Biotechnology Research Institute, City for Scientific Research and Technology Applications, Burgelarab, Alexandria, Egypt. Email: abdelhaleem@yahoo.de

How to cite this article: (Harvard style)

Zaki, S.; Farag, S.; Abu Elreesh, G.; Elkady, M.; Nosier, M.; Abd El Haleem, D., (2011). Characterization of bioflocculants produced by bacteria isolated from crude petroleum oil. *Int. J. Environ. Sci. Tech.*, 8 (4), 831-840.