Prevalence of Overweight and Obesity in 7 to 18 Year-Old Children in Birjand/ Iran

Fatemeh Taheri¹, MD; Toba Kazemi^{*1}, MD

1. Birjand University of Medical Sciences, Birjand, IR Iran

Received: Sep 25, 2008; Final Revision: Nov 03, 2008; Accepted: Feb 02, 2009

Abstract

Objective: The aim of this study was to estimate the prevalence of overweight and obesity in 7 to 18-year-old children in Birjand (east Iran) in 2005-2006. Individuals selected using multistage stratified random sampling.

Methods: This cross-sectional study was performed on 6093 students (2995 boys and 3098 girls) aged 7-18 years in Birjand (2005-2006). Subjects were selected via step-wised random sampling in four districts of the city. Body weight and height were measured directly. Percentiles were calculated for body mass index (BMI) Overweight and obesity was defined based on the 85th and 95th percentiles of body mass index for age and sex, respectively, as proposed by CDC in 2000.

Findings: The overall prevalence rates of overweight and obesity were 4.8% and 1.8%, respectively. Overweight prevalence varied by age from 1.6% to 9.1% in girls and 0.5% to 7.8% in boys, with obesity rate from 0.8% to 2.5% in girls and 0.5% to 3.7% in boys.

Conclusion: According to this study, prevalence rate of overweight and obesity in Birjandi children is lower than that in many other parts of Iran and some neighboring countries.

Iranian Journal of Pediatrics, Volume 19 (Number 2), June 2009, Pages: 135-140

Key Words: Obesity; Overweight; Students; Body mass index; Prevalence

Introduction

The prevalence of obesity in children and adolescents is increasing worldwide^[1-2]. In industrialized countries, increased prevalence of obesity among children was also reported

in studies periodically conducted by the Health Center of these countries^[3-5]. Accumulating evidence shows that the epidemic extends to the developing countries as well, in addition to an ongoing problem of under-nutrition in the

* Correspondence author;

Address: Valiassr Hospital, Ghaffari St, Birjand, South Khorassan, IR Iran E-mail: med_847@yahoo.com-09155610860

latter^[6-8]. The prevalence of obesity among US children is approximately of 22-30% and has increased at an accelerating rate in the past several years^[9-10].

Obesity in children and adolescents is a serious issue not only because of the health consequences in childhood and adolescence ^[11, 12] but also because of the greater risk of obesity in adulthood^[13]. Several methods have been created in order to measure body fat and obesity, during childhood and adolescence. Body mass index (BMI) is the preferred method of expressing body fat percentile of groups. It is widely accepted that a BMI between the 85th and 95th percentiles is defined as overweight, and a BMI greater than the 95th percentile as obesity ^[14].

In Birjand, east of Iran, there was limited research on childhood obesity and the present study aims at determining the prevalence of overweight and obesity among children aged 7-18 years, comparing the prevalence of overweight and obesity between males and females and between age groups. Children start school in Iran as soon as they become 6years old. For 5 years, they go to primary or elementary school (7-11 years old). Then another 3 years are passed as secondary or guidance school (12-14 years old). High school is 4 years (15-18 years old).

Subjects and Methods

The study was carried out at the schools of Birjand. We studied 6093 students (2995 boys and 3098 girls) aged 7-18 years from 2005-2006. Participants were selected through the systematic sampling technique

The measurements of body height and weight were carried out by trained medical students in the morning. Body weight (in kilograms) was measured to the nearest 0.1 kg with an electronic scale (SECA; Germany). Body height was measured to the nearest 0.5 cm as the adolescents stood erect against a vertical wall-mounted scale with heels, buttocks, and occiput in the Frankfort plane with anthropometric square. The subjects were dressed with light underclothes and wore no shoes throughout the measurements. BMI (kg/m²) was calculated as the ratio of the body weight to the square of body height.

Data were processed and analyzed by the EPSS software version 11.5. Estimations of the prevalence of overweight and obesity were based on CDC in 2000 (in excess of the 85th and 95th percentiles, respectively). The chi-square test, with a significance level of 5% (P<0.05) was used to compare the prevalence rates, taking into consideration gender and age group.

Findings

The sample, representing children aged between 7-18 years, consisted of 6093 subjects (3098 girls, 2995 boys). The overall prevalence of overweight and obesity was 4.8 and 1.8 percent, respectively.

Comparison of overweight and obesity prevalence in primary, secondary and high school children are shown in tables 1-4. Table 5 shows overweight and obesity prevalence in relation to sex. Prevalence of

Table 1: The prevalence of overweight and
obesity in primary school (using the CDC BMI
reference criteria)

Age (years)	Overweight	Obesity
<7y n=271	9(3.3%)	4(1.5%)
8y - 8y & 11 mo n=364	5(1.4%)	2(0.5%)
9 y- 9 y & 11 mo n=363	6(1.6%)	5(1.4%)
10y - 10 y & 11 mo n=383	6(1.6%)	4(1.1%)
11y - 11y & 11 mo n=415	18(4.3%)	7(1.7%)
All n=1796	44(2.4%)	22(1.2%)

Table 2: The prevalence of overweight andobesity in secondary school (using the CDCBMI reference criteria)

Age (years)	Overweight	Obesity
12y -12y & 11 mo n=683	24(3.5%)	6(0.9%)
13y -13y 11 mo n=667	38(5.7%)	13(1.9%)
14y - 14y & 11 mo n=617	33(5.3%)	11(1.8%)
All n=1967	95(4.8%)	30(1.5%)

Table 4: Comparison of prevalence ofoverweight and obesity in our subjects by typeof school

Type of school	Overweight	Obesity	
Primary school n=1796	44(2.4%)	22(1.2%)	
Mid school n=1967	95(4.8%)	30(1.5%)	
High school n=2330	154(6.6%)	61(2.6%)	
Total=6093	293(4.8%)	113(1.8%)	
X ² =52.2; df=4; <i>P</i> <0.001			

overweight and obesity was 5.8%, 1.5%, respectively, among girls and 3.7%, and 2.1% among boys. The prevalence of overweight was highest at 15 years of age among boys and girls (7.8%, 9.1%, respectively). The prevalence of obesity was highest at 18 years of age among boys (3.7%) and at 15 years among girls (2.5%).

Discussion

The purpose of this study was to provide data on the prevalence of overweight and obesity

Table 3: The prevalence of overweight andobesity in high-school (using the CDC BMIreference criteria)

Age (years)	Overweight	Obesity		
15y -15y & 11mo n=696	59(8.5%)	21(3%)		
16y -16y & 11mo n=641	41(6.4%)	14(2.2%)		
17y -17y & 11mo n=581	35(6%)	14(2.4%)		
18y -18y & 11mo n=412	19(4.6%)	12(2.9%)		
All n=2330	154(6.6%)	61(2.6%)		

among East Iranian children. The results of the present study demonstrated that the overall prevalence of overweight and obesity were 4.8% and 1.8%, respectively.

The prevalence of overweight and obesity among the East Iranian children (7-18 y) was lower than those reported in literature. Birjand, center of South-Khorasan province, is located in the east of Iran and one of the most deprived areas in Iran (Fig 1).

Moayeri reported that the prevalence of overweight and obesity among Tehrani students was 21.1% and 7.8%, respectively^[7]. In Tabriz, west Iran, the prevalence of



Fig 1: Map of Iran showing geographic regions and the location of Birjand

1 50	Girls			Boys		
Age	Total	Overweight	Obesity	Total	Overweight	Obesity
7y (n=271)	130	7(5/4%)	1(0/8%)	141	3(2/1%)	2(1/4%)
8y (n=364)	173	4(2/3%)	1(0/6%)	191	1(0/5%)	1(0/5%)
9y (n=363)	187	3(1/6%)	3(1/6%)	176	2(1/1%)	3(1/7%)
10y (n=383)	172	5(2/9%)	1(0/6%)	211	3(1/4%)	1(0/5%)
11y (n=415)	209	13(6/2%)	5(2/4%)	206	2(1%)	5(2/4%)
12y (n=683)	363	16(4/4%)	3(0/8%)	320	3(0/9%)	8(2/5%)
13y (n=667)	332	24(7/2%)	3(0/9%)	335	10(3%)	14(4/2%)
14y (n=617)	368	23(6/2%)	4(1/1%)	249	7(2/8%)	10(4/1%)
15y (n=696)	361	33(9/1%)	9(2/5%)	335	12(3/6%)	26(7/8%)
16y (n=641)	334	23(6/9%)	6(1/8%)	307	8(2/6%)	18(5/8%)
17y (n=581)	273	20(7/3%)	6(2/2%)	308	8(2/6%)	15(4/9%)
18y (n=412)	196	10(5/1%)	4(0/8%)	216	8(3/7%)	9(4/2%)
total	3098	181(5/8%)	46(1/5%)	2995	67(2/1%)	112(3/7%)

Table 5: The prevalence of overweight and obesity in our subjects by gender (using the CDC BMIreference criteria)

overweight and obesity in high-school girls was 11.1% and 3.6%, respectively ^[15]. In Shiraz, south Iran, the prevalence of overweight and obesity among adolescents (13-18 year-olds) was 11.3% and 2.9%, respectively ^[16].

The reasons for the lower prevalence of overweight and obesity among our subjects are largely unknown. But its reason may be sought in genetic factors, environmental factors (such as the hot weather), type of diet, extent of physical activity, different definitions of overweight and obesity and low socioeconomical status in east Iran.

The prevalence of overweight and obesity among 1253 students in Sana'a City (Yemen) was 6.2%, and 1.8%, respectively. The prevalence of overweight and obesity in 8300 Childhood in central Iraq was 6% and 1.3%, respectively. The results of our study was similar to those of these two reports^[17-18]. But other studies in Asia (Thailand, Saudi Arabia, China), Europe (France, Greece, Spain), Africa (South Africa) and America (Mexico, New York State) reported higher prevalence of overweight and obesity among students^[8,10,19, 21-4].

Some countries showed significant gender differences in overweight and obesity prevalence of children. In particular, most of the reports from Asia and Europe (Qatar, Turkey Greece), showed higher rates among boys than among girls^[6,21,25]. On the other hand, Saudi Arabia, Tehran and Kuwait demonstrated the opposite trend^[7,8,26]. But it was not different by gender in French and Gibraltar^[20,27]. We observed higher rates of overweight among adolescent girls but the rates of obesity were higher in boys.

The limitation of our study is that a descriptive study. In our study, we could not prove any causal relationship between obesity and its risk factors. There is no background for us to compare present status

with the past one. We recommend periodic studies in this field especially in this deprived area of Iran.

Conclusion

This is the first study on overweight in Birjandi children. In our study, the overall prevalence rates of overweight and obesity were 4.8% and 1.8%, respectively. Prevalence rate of overweight and obesity in Birjandian children is lower than that in many other parts of Iran and some neighboring countries.

Acknowledgment

This study was supported by grant from Birjand University of Medical Sciences. We would like to gratefully thank all the students who participated in this study.

References

- 1. World Health Organization (WHO). Obesity: preventing and managing the global epidemic. Report of a WHO consultation on obesity, Geneva, 3-5 June, 1997. Geneva: WHO, 1998.
- Kelishadi R. Childhood overweight, obesity, and the metabolic syndrome in developing countries. Epidemiol Rev. 2007;29(1):62-76.
- 3. Moreno LA, Sarría A, Fleta J, et al. Trends in body mass index and overweight prevalence among children and adolescents in the region of Aragon (Spain) from 1985 to 1995. Int J Obes Relat Metab Disord. 2000;24(7):925-31.
- Kalies H, Lenz J, von Kries R. Prevalence of overweight and obesity and trends in body mass index in German pre-school children, 1982-1997. Int J Obes Relat Metab Disord. 2002;26(9):1211-7.

- Kautiainen S, Rimpelä A, Vikat A, et al. Secular trends in overweight and obesity among Finnish adolescents in 1977-1999. Int J Obes Relat Metab Disord. 2002; 26(4):544-52.
- 6. Oner N, Vatansever U, Sari A, et al. Prevalence of underweight, overweight and obesity in Turkish adolescents. Swiss Med Wkly. 2004; 134(35-36): 529-33.
- Moayeri H, Bidad K, Aghamohammadi A, et al. Overweight and obesity and their associated factors in adolescents in Tehran, Iran, 2004-2005. Eur J Pediatr. 2006;165(7):489-93.
- 8. El-Hazmi MA, Warsy AS. A Comparative study of prevalence of overweight and obesity in children in different provinces of Saudi Arabia. J Trop Pediatr. 2002; 48(3):172-7.
- 9. Figueroa-Colon R, Franklin FA, Lee JY, et al. Prevalence of obesity with increased blood pressure in elementary school-aged children. South Med J. 1997;90(8):806-13.
- 10. Wang Y, Monteiro C, Popkin BM. Trends of obesity and underweight in older children and adolescents in the United States, Brazil, China, and Russia: Am J Clin Nutr. 2002;75(6):971-7.
- 11. Plourde G. Impact of obesity on glucose and lipid profiles in adolescents at different age groups in relation to adulthood. BMC Fam Pract. 2002;3:18.
- 12. Dietz WH. Childhood weight affects adult morbidity and mortality. J Nutr. 1998; 128(2 Suppl):411S-4S.
- Serdula MK, Ivery D, Coates RJ, et al. Do obese children become obese adults? A review of the literature. Prev Med. 1993; 22(2):167-77.
- 14. Dietz WH, Bellizzi MC. Introduction: the use of body mass index to assess obesity in children. Am J Clin Nutr. 1999;70(1): 123S-5S.
- 15. Gargari BP, Behzad MH, Ghassabpour S, et al. Prevalence of overweight and obesity among high-school girls in Tabriz, Iran, in 2001: Food Nutr Bull. 2004;25(3):288-91.
- 16. Mostafavi H, Dabagh Manesh MH, Zare N. Prevalence of obesity and over- weight in adolescents and adult population in

Shiraz: Iran J Endcorinol Metab. 2005; 7(1):57-66. (Persian)

- 17. Raja'a YA, Bin Mohanna MA. Overweight and obesity among school children in Sana'a City, Yemen. Ann Nutr Metab. 2005; 49(5): 342-5.
- Lafta RK, Kadhim MJ. Childhood obesity in Iraq: prevalence and possible risk factors. Ann Saudi Med. 2005;25(5):389-93.
- 19. Langendijk G, Wellings S, van Wyk M, et al. The prevalence of childhood obesity in primary school children in urban Khon Kaen, northeast Thailand. Asia Pac J Clin Nutr. 2003;12(1):66-72.
- 20. Klein-Platat C, Wagner A, Haan MC, et al. Prevalence and sociodemographic determinants of overweight in young French adolescents: Diabetes Metab Res Rev. 2003;19(2):153-8.
- 21. Krassas GE, Tzotzas T, Tsametis C, et al. Prevalence and trends in overweight and obesity among children and adolescents in Thessaloniki, Greece. J Pediatr Endocrinol Metab. 2001;14(Suppl 5):1319-26.

- 22. Armstrong ME, Lambert MI, Sharwood KA, Lambert EV. Obesity and over-weight in South African primary school children -the Health of the Nation Study. S Afr Med J. 2006;96(5):439-44.
- 23. Del Río-Navarro BE, Velázquez-Monroy O, Sánchez-Castillo CP, et al. The high prevalence of overweight and obesity in Mexican children. Obes Res. 2004;12(2): 215-23.
- 24. Wolfe WS, Campbell CC, Frongillo EA, et al. Overweight schoolchildren in New York State: prevalence and characteristics. Am J Public Health. 1994;84(5):807-13.
- Bener A. Prevalence of obesity, overweight, and underweight in Qatari adolescents. Food Nutr Bull. 2006;27(1): 39-45.
- 26. Ezzat KA, Fawzia A. Nutritional status survey of preschool children in Kuwait. East Mediter Health J. 1996;3(2):386-95.
- 27. Kirk SF, McLeod M. The prevalence of overweight and obesity in children aged 4 to 12 years in Gibraltar. Public Health Nutr. 2003;6(4):329-31.