

Alternating Ibuprofen and Acetaminophen in the Treatment of Febrile Hospitalized Children Aged 9-24 Months

Nader Pashapour*¹, MD; Ali A Maccooei¹, MD; Sariyeh Golmohammadlou², MD

1. Department of Pediatrics, Urmia University of Medical Sciences, Urmia, IR Iran
2. Department of Mother and Child Health, Urmia University of Medical Sciences, Urmia, IR Iran

Received: Aug 12, 2008; Final Revision: Dec 13, 2008; Accepted: Jan 13, 2009

Abstract

Objective: Recently, studies reported the emergence and increasing popularity of the alternate use of acetaminophen and ibuprofen. The aim of the present study was to compare the clinical effectiveness of acetaminophen alone with an alternative regimen of acetaminophen and ibuprofen in hospitalized infants aged 9-24 months with fever of non bacterial origin.

Methods: The study was performed between March 2006 and December 2007, in pediatric center of two hospitals in Urmia. Patients were randomly separated into two groups. Eligible cases were febrile hospitalized patients aged between 9 and 24 months; whose rectal temperature was $\geq 38.5^{\circ}\text{C}$. Infants of case group received 10 mg/kg ibuprofen alternating with 15 mg/kg acetaminophen every four hours. Infants of control group received 15 mg/kg acetaminophen every four hours. Temperature of patients in the two groups was registered 2, 4, 5, 7, and 8 hours after drug administration. The study was completed when each group received 35 cases.

Findings: Seventy infants were divided into two groups. The case group consisted of 19 males and 16 females, in control group there were 18 males and 17 females. The results revealed significant difference between the two groups in lowering fever at 4, 5, 7 and 8 hours after treatment was initiated, but there was no significant difference at 2 hours after drug administration.

Conclusion: An alternating treatment of acetaminophen (15 mg/kg) and ibuprofen (10mg/kg) 4 hours later is more effective than acetaminophen in lowering fever in 9 to 24 month old infants.

Iranian Journal of Pediatrics, Volume 19 (Number 2), June 2009, Pages: 164-168

Key Words: Acetaminophen; Ibuprofen; Antipyretic; Fever; Infant

* Correspondence author;

Address: Department of Pediatrics, Faculty of Medicine, Urmia University of Medical Science, Motahari Ave, Urmia, IR Iran

E-mail: npashapor@yahoo.com

Archive of SID

Introduction

Fever is one of the most common complaint and signs of illness in office-based primary care pediatric practice, accounting for 19% to 30% of visits^[1]. Although most experts consider fever a beneficial physiologic response to the infectious process, often it leads to patient irritability and stress as well as parental anxiety^[2].

Acetaminophen is very widely used as an antipyretic owing to its availability and good safety^[3]. Ibuprofen, like acetaminophen is an effective antipyretic and is well tolerated. There is no evidence of a difference with acetaminophen in the short term adverse effects, has good gastrointestinal tract absorption feature, reaching peak plasma concentration in one hour; the maximum temperature reduction peaks within 30 minutes^[4].

In addition, studies of parental knowledge of fever exposed unfounded fears (Fever phobia) and misconception forced many parents and some physicians to inappropriate care^[5]. New studies reported the emergence and increasing popularity of the alternative use of acetaminophen and ibuprofen^[6,7].

The aim of the present study was to compare the clinical effectiveness of acetaminophen alone with an alternative regimen of acetaminophen and ibuprofen in hospitalized infants aged 9-24 months with fever of non bacterial origin.

Subjects and Methods

This study was performed between March 2006 and December 2007, in 2 pediatric center in Urmia, which are tertiary level and affiliated to Urmia University of Medical Sciences. The study was approved by Research Center Board and the Ethics Committee of the University.

Eligible cases were febrile hospitalized patients aged between 9 and 24 months,

whose rectal temperature was $\geq 38.5^{\circ}\text{C}$. Exclusion criteria included any of the following: intolerance or major complications of the administered drug such as abdominal pain, vomiting, hypothermia, acute or chronic renal and hepatic disease, asthma, metabolic disease, bleeding disorder, convulsion, children with concurrent or previous intake of antibiotics and antipyretics, clinical or laboratory evidence of bacterial infection, and cases that did not complete the study. We chose non bacterial infection to limit scope of fever. Stool analysis, stool culture, cell blood count (CBC), C-reactive protein (CRP), erythrocyte sedimentation rate (ESR), chest x-ray, electrolytes, blood sugar, blood urea nitrogen and creatinine were evaluated in all cases.

In this clinical trial study, patients were randomly separated into two groups. Written consent was obtained from parents. Infants of case group received 10 mg/kg ibuprofen alternating with 15 mg/kg acetaminophen every four hours. Infants of control group received 15 mg/kg acetaminophen every four hours. Temperature of patients of the two groups was taken at admission, 2, 4, 5, 7, and 8 hours after drug administration. According to sample size calculation formula, for this study twenty patients would be sufficient to detect an assumed difference of 30% between groups in the reduction of fever.

Chi-square and t-test were used to compare the probability of sex, age and temperature difference between case and control groups.

Findings

During the study period, of the 76 cases, 7 cases were excluded because the parents did not want to continue participating in the study. The case groups consisted of 19 males and 16 females and in control group there were 18 males and 17 females.

Mean age on admission was 17.2 ± 5.0 months (CI 95% 15.5-18.9) for case and

Table 1: Data of patients with fever at admission (in case and control groups)

Variables	Patients*	Mean (SD)	P value
Age	Case (N=35)	17.2 (5.0)	0.8
Mean (SD)	Control (N=35)	17.0 (5.1)	
Sex (F/M)	Case (N=35)	16/19	0.7
Frequency	Control (N=35)	17/18	
Weight	Case (N=35)	11.4 (1.36)	0.2
Mean (SD)	Control (N=35)	11.9 (3.03)	
Fever (admission)	Case (N=35)	39.2 (0.52)	0.03
Mean (SD)	Control (N=35)	39.3 (0.47)	

*Case group: treated with ibuprofen alternating with acetaminophen
Control Group: treated with acetaminophen

17±5.0 months (CI 95% 15.3-18.7) for control group. Mean temperature on admission was 39.2°C (CI 95% 39.1-39.4) and 39.3°C (CI 95% 39.1-39.4) for case and control group respectively.

Mean weight on admission was 11.4 (CI 95% 10.9- 11.9) for case and 11.9 kg. (CI 95% 10.0- 13.0) for control group. Both groups had comparable clinical characteristics at

admission including age, weight and temperature (table 1). As table 2 shows, there is significant difference between two groups in lowering fever at 4, 5, 7 and 8 hours after treatment was initiated, but there is no significant difference at 2 hours after drug was administered. Major complications of administered antifebrile drugs were not observed in patients.

Table 2: Comparison of groups after drug administration

Temperature (times)*	Groups ‡	Mean (SD)	P Value
Fever 2	Case N=35)	38.8 (0.59)	0.7
	Control (N=35)	38.8 (0.47)	
Fever 4	Case (N=35)	38.4 (0.34)	0.048
	Control (N=35)	38.5 (0.30)	
Fever 5	Case (N=35)	38.0 (0.47)	0.04
	Control (N=35)	38.2 (0.38)	
Fever 7	Case (N=35)	38.0 (0.48)	0.04
	Control (N=35)	38.2 (0.57)	
Fever 8	Case (N=35)	37.7 (0.46)	0.02
	Control (N=35)	38.0 (0.52)	

* Each hour after drug administration that temperature of patients was measured

‡ Case group: treated with ibuprofen alternating with acetaminophen

Control Group: treated with acetaminophen

Discussion

The aim of this study was evaluation of clinical effectiveness of acetaminophen alone, in comparison with an alternative regimen of acetaminophen and ibuprofen. As data of table 1 shows, there was no significant difference between the two groups regarding distribution of age, sex, weight and degree of fever at admission. So any significant difference in reduction of fever could be attributed to the method of treatment.

As table 2 shows, there was no significant difference between the two groups in reduction of fever after two hours of antipyretic administration. Analysis of results at 4 hours post-treatment revealed that reduction of fever in group two which received ibuprofen (10mg/kg) was significantly less than the group treated with acetaminophen (15mg/kg). Meta analysis of seventeen double blind, randomized controlled trials in children who received ibuprofen and acetaminophen to treat fever, revealed that ibuprofen (5-10mg/kg) was a more effective antipyretic than acetaminophen (10-15mg/kg) at 2, 4 and 6 hours post-treatment^[8]. Robertson disagrees with aforementioned results and believes there is not enough evidence to support that ibuprofen more effectively reduces fever^[9]. Results of a study by Autert et al to find effect of the kind of drug on parent's opinion revealed that in spite of equivalent efficacy and tolerability of ibuprofen at a dose of 10 mg/kg and paracetamol at a dose of 15 mg/kg, parents' opinion was in favor of ibuprofen. Parents' positive opinion about ibuprofen, helped allay their anxiety over the treatment of their child^[10].

In this study comparison between two groups at 5, 7, 8 hours after treatment, showed significant lower temperature in the case group (table 2). Parallel to our results, another study was conducted to compare the antipyretic benefit of acetaminophen or ibuprofen monotherapy with an alternating regimen of both drugs in young children aged

6 to 36 months. Researchers concluded that an alternating treatment regimen of acetaminophen (12/5 mg/kg per dose) and ibuprofen (5 mg/kg per dose) every 4 hours for 3 days, regardless of the initial loading medication, is more effective than monotherapy in lowering fever in infants and children^[11].

Another study entitled as "Alternating ibuprofen and acetaminophen in treatment of febrile children", was conducted on patients aged 6 months to 14 years with rectal temperature ≥ 38.8 . Researchers concluded that a single dose of ibuprofen and acetaminophen was superior antipyretic regimen than ibuprofen monotherapy^[12].

Oral route was selected for administration of antipyretic drugs in this research. Nablusi et al designed a study to compare the antipyretic effectiveness of two rectal acetaminophen doses (15 mg/kg) and (35 mg/kg), to standard oral dose of 15mg/kg. They concluded that standard oral, rectal and high dose rectal acetaminophen had similar antipyretic effectiveness^[13].

There are not enough articles on alternating acetaminophen and ibuprofen use in febrile children. The short duration of monitoring and of follow-up was the limitation of our study. In spite of results of this study; we do not recommend alternating regimen in clinical practice until further studies confirm the results.

Conclusion

An alternating treatment of acetaminophen (15 mg/kg) and ibuprofen (10 mg/kg) 4 hours later is more effective than acetaminophen monotherapy in lowering fever at 5, 6, 7 and 8 hours after administration in 9 to 24 month old infants. At four hours after drug intake ibuprofen reduces fever better than acetaminophen. Further studies are needed to confirm this conclusion.

Acknowledgment

Authors would like to thank members of University Council of Research and Ethics Committee of the University for their guides in all stages of the study.

References

1. Jonna Briggs Institute. Management of the child with fever. Best Practice: Evidence Based Practice Information Sheets for Health Professionals. 2001;5(5):1-6. Available at: http://www.joannabriggs.edu.au/pdf/BPISEng_5_5.pdf. Access date: Jan 2002.
2. Crocetti M, Moghbeli N, Serwint J: Fever phobia revisited: have parental misconceptions about fever changed in 20 years? *Pediatrics*. 2001;107(6):1241-6.
3. Lesko SM, Mitchell AA: The safety of acetaminophen and ibuprofen among children younger than two years old. *Pediatrics*. 1999;104(4):e39.
4. Schmitt BD. Fever phobia. Misconceptions of parents about fever. *Am J Dis Child*. 1980;134(2):176-81.
5. Purssell E. Treating fever in children: Paracetamol or ibuprofen? *Br J Commun Nurs*. 2002;76(4):316-20.
6. Diez Domingo J, Burgos Ramirez A, Garrido Garcia J, et al. Use of alternating antipyretics in the treatment of fever in Spain. *An Esp Pediatr*. 2001; 55(6):503-10.
7. Mayoral CE, Marino RV, Rosenfeld W, et al. Alternating antipyretics: Is this an alternative? *Pediatrics* 2000; 105(5): 1009-12.
8. Perrott DA, Piira T, Goodenough B, et al. Efficacy and safety of acetaminophen vs ibuprofen for treating children's pain or fever. *Arch Pediatr Adolesc Med*. 2004;158(6):521-6.
9. Robertson J. Review: no evidence exists that paracetamol and ibuprofen differ for short term pain relief or safety in children. *Evi Based Nurs*. 2005;8(1):10.
10. Autret-leca E, Gibb IA, Goulder MA. Ibuprofen versus paracetamol in pediatric fever: objective and subjective finding from a randomized, blinded study. *Curr Med Res Opin*. 2007;23(9): 2205-11.
11. Sarrell EM; Wielunsky E, Cohen HA. Antipyretic treatment in young children with fever: acetaminophen, ibuprofen, or both alternating in a randomized, double-blind study. *Arch Pediatr Adolesc Med*. 2006;160(2):197-202.
12. Nabulsi M, Tamim H, Mahfoud Z, et al. Alternating ibuprofen and acetaminophen in the treatment of febrile children: a pilot study. *BMC Med*. 2006;4:4.
13. Nablusi M, Tamim H, Sabra R, et al. Equal antipyretic effectiveness of oral and rectal acetaminophen: A randomized controlled trial. *BMC Pediatrics*. 2005;5:35.