

## Estimation of Macronutrient Content in Kindergartens Meals: Food Composition Tables or Chemical Analysis?

Lazarevic Konstansa<sup>1,2</sup>, MD; Stojanovic Dusica<sup>2,3</sup>, MD; Bogdanovic Dragan<sup>1,2</sup>, MD

<sup>1</sup>State University of Novi Pazar, Serbia; <sup>2</sup>Public Health Institute Nis; <sup>3</sup>School of Medicine, University of Nis, Serbia

Received: Nov 01, 2013; Accepted: Aug 09, 2014; First Online Available: Sep 27, 2014

### Abstract

**Objective:** The nutrition of children in kindergartens has a great significance for proper growth and development of children. In order to save time and money, the control of macronutrients content is performed by calculations using food composition tables instead of performing a chemical analysis.

**Methods:** We examined the macronutrients content of 240 whole day meals using food composition tables and performed chemical analysis of meals to determine adequacy and validity of food composition tables in calculation of macronutrient contents in kindergarten meals.

**Findings:** We established no correlation ( $P > 0.05$ ) between the value of proteins, fats and carbohydrates. Significant difference was established between the average content of proteins ( $t = 2.57$ ;  $P < 0.05$ ), and carbohydrates ( $t = 3.20$ ;  $P < 0.01$ ), but not with the content of fats in the meals ( $t = 1.26$ ;  $P > 0.05$ ) (food composition tables vs chemical analysis).

**Conclusion:** Until we establish new food composition tables, chemical analysis remains the only valid method for assessment of macronutrients content and energy value of a meal in kindergarten.

*Iranian Journal of Pediatrics, Volume 24 (Number 5), October 2014, Pages: 643-646*

**Key Words:** Macronutrient; Preschool Children; Diet; Kindergarten

### Introduction

Proper nutrition during the period of childhood is the foundation of good health in the adulthood and necessary for proper growth and development of children<sup>[1]</sup>.

As a consequence of the modern style of life, characterized by long working day of parents, most of children in developing countries spend 8-12 hours during the day in kindergartens<sup>[2,3]</sup>. For this reason the kindergarten diet has an important role in the daily intake of energy and nutrients. Proteins from diet are important for growth and

development during childhood, and dietary requirements (g/kg of body weight) are significantly increased in this period. Dietary fat and carbohydrates are energy sources, but dietary fats are also source of fat soluble vitamins and some fatty acids important for children<sup>[4]</sup>.

There are different methods for estimation of the composition of food: data from printed food composition tables, electronic databases, food industry data and data from scientific literature<sup>[5]</sup>. In order to save time and money, chemical analysis of meal (often cooked or prepared food) is increasingly rare and calculations by using food

\* Corresponding Author;

Address: Dr Milenka Hadzica 2/23 18000 Niš, Serbia

E-mail: koni33@hotmail.com

© 2014 by Pediatrics Center of Excellence, Children's Medical Center, Tehran University of Medical Sciences, All rights reserved.

composition tables are applied. In few of the studies, nutrient content in kindergarten meals was calculated by food composition tables - studies from Brazil<sup>[6]</sup>, China<sup>[7]</sup>, Finland<sup>[8]</sup> or by chemical analysis in Croatia<sup>[9]</sup> or by both methods as in Poland<sup>[10]</sup> and in our study.

The purpose of this article is to compare results of protein, carbohydrate and fat content in daily kindergartens meals obtained from food composition table and by chemical analysis, and to suggest how to ensure the correct calculation of macronutrients contents in kindergarten meals.

## Subjects and Methods

There are three meals in Niš kindergarten (breakfast, lunch and snack), planned by a nutritionist, a physician and a nurse.

**Use of food composition tables.** It was necessary to calculate the amount of consumed food, and then multiply the food intake by the mean nutrient content obtained from the food composition database. Daily amount of food used for preparation of 240 meals is calculated based on the warehouse supply list. Nutrient contents of meals were calculated by Serbian food composition tables<sup>[11]</sup>.

**Chemical analysis:** After the collection, the samples of meals in duplicate were transported to the laboratory at 4°C within one hour. They were analyzed for moisture, protein, fat and ash<sup>[12]</sup> in the laboratory of the Institute for Public Health, Niš (Serbia).

Descriptive statistics (mean, standard deviation) of macronutrient contents (g) was calculated using Microsoft Excel software. Student's t-test and linear correlation was applied to compare the values between two methods.

## Findings

Results for contents of macronutrients in kindergartens meals were similar for fats, but not proteins and carbohydrates when calculated by chemical analysis and food composition tables (Table 1). Significant correlation between proteins, fats and carbohydrates intake obtained from warehouse supply lists and food composition tables were not observed (Table 2).

## Discussion

Data on macronutrients content of meals in the collective nutrition, obtained from food composition tables are mostly informative.

Depending on the type of food, nutrient and table or software selected for comparison, there were significant statistical differences between results of laboratory analyses and results calculated though tables and software data<sup>[13-15]</sup>. There are several explanations for these results.

Firstly, there were differences in macronutrient estimations depending upon the choice of the food composition tables. Some food composition tables are not adequately documented (small number of analyses, inappropriate choice of analysis). Results from 18 laboratories in Europe showed statistically different values for macronutrients in the well-homogenized samples of foods<sup>[16]</sup>. Using food composition databases from different countries may increase errors in calculation of macronutrient intake, and there is tendency to standardize food composition databases all over the world<sup>[17]</sup>.

Secondly, there is a variation in food composition depending on environmental, genetic and processing influences such as feed, soil,

**Table 1:** Mean macronutrient content in analyzed samples of meals by chemical analysis and food composition tables

Macronutrient	Food composition tables Mean (SD) (g)	Chemical analysis Mean (SD) (g)	t-test	P. value
<b>Protein</b>	40.80 (4.35)	35.0 (6.47)	2.57	0.02
<b>Fats</b>	38.18 (6.32)	34.99 (6.07)	1.26	0.2
<b>Carbohydrates</b>	151.80 (13.45)	133.93 (13.96)	3.20	0.004

SD: Standard Deviation

**Table 2:** Correlations between macronutrient content in analyzed samples of meals by chemical analysis and food composition tables

Parameter	Coefficient of correlation (r)	P. value
Protein	-0.171	0.6
Fat	0.290	0.4
Carbohydrate	0.010	0.97

climate, genetic resources (varieties/cultivars, breeds), storage conditions, processing, fortification and market share<sup>[18]</sup>.

Thirdly, the majority of nutrient databases provide information mainly on the composition of raw foods. Cooking methods, times, temperatures and food preparation habits vary widely by countries, and that can influence the macronutrient content<sup>[19]</sup>.

Fourthly, we don't have data about food plate waste in kindergartens.

Contrary to the results of previous studies, Bedogni et al concluded that food composition tables may be used to assess energy, carbohydrate, lipid and protein intake in the military settings<sup>[20]</sup>. Many food composition tables, including tables used in our study did not consider yield and retention factors. Camilli et al. found that when yield factor and retention factors were applied, the results of the computed nutrients according to food composition tables showed a satisfactory degree for protein and lipid contents<sup>[21]</sup>.

## Conclusion

It is evident that food composition tables must include the nutrient composition of cooked and prepared foods, as well as yield factor and retention factor.

Taking into account the time that preschool children spend in kindergartens and the importance of proper nutrition for child development in that life period, the chemical analysis of foods (dishes) must be applied for situations such as preschool institutional nutrition.

## Authors' Contribution

L. Konstansa: concept/design, acquisition of data, data analysis/interpretation, drafting of the manuscript

S. Dusica: acquisition of data, data critical revision of the manuscript

B. Dragan: concept/design, acquisition of data, data analysis/interpretation

**Conflict of Interest:** None

## References

1. Lynch JW, Smith GD. A life-course approach to chronic disease epidemiology. *Annu Rev Public Health* 2005;26:1-35.
2. Federal Interagency Forum on Child and Family Statistics. America's Children: Key National Indicators of Well-Being, 2002. Federal Interagency Forum on Child and Family Statistics, Washington, DC: U.S. Government Printing Office. Available at: [http://www.childstats.gov/pdf/ac2002/ac\\_02.pdf](http://www.childstats.gov/pdf/ac2002/ac_02.pdf) Access date: Nov 1, 2013.
3. Directorate for Health and Social Affairs. Food and mealtimes in childcare institutions. Oslo, 2005. Available at: [http://www.shdir.no/vp/multimedia/archive/00005/IS-1324\\_5639a.pdf](http://www.shdir.no/vp/multimedia/archive/00005/IS-1324_5639a.pdf). Access date: Dec 10, 2005. (In Norwegian)
4. Sardesai Vishwanath M. Requirements for energy, carbohydrates, fat and proteins. In: Introduction to Clinical Nutrition. Second edition, revised and expanded, CRC Press 2003; 33-40.
5. Pennington, JA. Applications of food composition data: Data sources and considerations for use. *J Food Composition Analysis* 2008;21:S3-S12.
6. Gomes RC, da Costa TH, Schmitz Bde A. Dietary assessment of pre-school children from Federal District Brazil. *Arch Latinoam Nutr* 2010;60(2):168-74.
7. Yin S, Su Y, Liu Q, et al. Dietary status of preschool children from day-care kindergartens in six cities of China. *Wei Sheng Yan Jiu* 2002;31(5):375-8.
8. Lehtisalo J, Erkkola M, Tapanainen H, et al. Food consumption and nutrient intake in day care and at home in 3-year-old Finnish children. *Public Health Nutr* 2010;13(6A):957-96.
9. Jagic V, Bosnir J, Racz A, Jelusic S. Caloric and nutritive value of kindergarten meals in Zagreb after implementation of the new national nutritive recommendation and standards 2007. *Paediatr Croat* 2011;55:11-16. [in Croatian]

10. Frackiewicz J, Ring-Andrzejczuk K, Gronowska-Senger A. Energy and selected nutrients content in pre-school children diet of Warsaw district. *Rocz Panstw Zakl Hig* 2011;62(2):181-5.
11. Jokic N, Dimic M, Pavlica M, Mirilov M. Food composition tables: the content of basic nutrients and protective substances and energy value of food. Institute for Economics Households Serbia, Belgrade 1999 (in Serbian).
12. Association of Official Analytical Chemists. Official methods of analysis, 15th ed. Arlington: Association of Official Analytical Chemists, 1990.
13. Yoshita K, Miura K, Okayama A, et al. A validation study on food composition tables for the international cooperative INTERMAP study in Japan. *Environ Health Prev Med* 2005;10(3):150-6.
14. Ribeiro P, de Moraes TB, Colugnati FA, et al. Food composition tables: laboratory comparative analysis. *Rev Saude Publica* 2003;37(2):216-25.
15. Kim ES, Ko YS, Kim J, et al. Food composition table-based estimation of energy and major nutrient intake in comparison with chemical analysis: a validation study in Korea. *Tohoku J Exp Med* 2003; 200(1):7-15.
16. Hoilman PCH, Katan MB. Results of the Eurofoods trial on between-laboratory variation in the analysis of macronutrients in foods. *Fresenius Z Anal Chem* 1987;326:690-5.
17. Charrondière UR, Vignat J, Riboli E. Comparable nutrient intake across countries is only possible through standardization of existing food composition tables. *IARC Sci Publ* 2002;156:45-7.
18. FAO. International Network of Food Data Systems (INFOODS). Available from: <http://www.fao.org/infoods/infoods/food-composition-challenges/en/>
19. Bordin K, Kunitake MT, Aracava KK, et al. Changes in food caused by deep fat frying - a review. *Arch Latinoam Nutr* 2013;63(1):5-13.
20. Bedogni G, Bernini Carri E, Gatti G, et al. Comparison of food composition tables and direct chemical analysis for the assessment of macronutrient intake in a military community. *Int J Food Sci Nutr* 1999; 50(1):73-9.
21. Camilli E, Gabrielli P, Marconi S, et al. Recipe composition data: Calculation procedure can be considered a valid alternative to chemical analysis for all nutrients? Abstracts of "ChimAlSi\_2012" IX° Italian Congress of Food Chemistry. "Food, Functional Foods and Nutraceuticals" Ischia (NA), Italy, June 03-07. 2012; P: 61.