

Short Term Outcome of Neonates Born With Abnormal Umbilical Cord Arterial Blood Gases

Mousa Ahmadpour-Kacho^{1,*}; Yadollah Zahedpasha¹; Mohsen Hagshenas¹; Zahra Akbarian Rad¹; Bahram Sadat Nasser¹; Ali Bijani¹

¹Non-Communicable Pediatric Diseases Research Center, Department of Pediatrics, Babol University of Medical Sciences, Babol, IR Iran

*Corresponding author: Mousa Ahmadpour-Kacho, Non-Communicable Pediatric Diseases Research Center, Department of Pediatrics, Babol University of Medical Sciences, Babol, IR Iran. E-mail: mousa_ahmadpour@hotmail.com

Received: December 16, 2014; Revised: April 9, 2015; Accepted: April 21, 2015

Background: Umbilical arterial blood gas (UABG) analysis is more objective than other methods for predicting neonatal outcome. Acidemic neonates may be at risk for unfavorable outcome after birth, but all neonates with abnormal arterial blood gas (ABG) analysis do not always have poor outcome.

Objectives: This study was carried out to determine the short term outcome of the neonates born with an abnormal ABG.

Patients and Methods: In a cohort prospective study 120 high risk mother-neonate pairs were enrolled and UABG was taken immediately after birth. All neonates with an umbilical cord pH less than 7.2 were considered as case group and more than 7.2 as controls. Outcomes like need to resuscitation, admission to newborn services and/or NICU, seizure occurrence, hypoxic ischemic encephalopathy (HIE), delayed initiation of oral feeding and length of hospital stay were recorded and compared between the two groups. P value less than 0.05 was considered as being significant.

Results: Comparison of short term outcomes between normal and abnormal ABG groups were as the following: need for advanced resuscitation 4 vs. 0 ($P = 0.001$), NICU admission 16 vs. 4 ($P = 0.001$), convulsion 2 vs. 0 ($P = 0.496$), HIE 17 vs. 4 ($P = 0.002$), delay to start oral feeding 16 vs. 4 ($P = 0.001$), mean hospital stay 4 vs. 3 days ($P = 0.001$). None of the neonates died in study groups.

Conclusions: An umbilical cord PH less than 7.2 immediately after birth can be used as a prognostic factor for unfavorable short term outcome in newborns.

Keywords: Umbilical Arterial Blood Gas; pH; Propionic Acidemia; Assessment, Patient Outcome

1. Backgrounds

Various clinical and laboratory variables have been used to predict the development of poor neonatal outcome immediately after birth.

pH assessment of cord ABG analysis is an objective diagnostic method for high risk newborns that may be at risk for poor neurologic outcome like birth asphyxia (1). Apgar score is not an objective method to assess the outcome. Most cerebral palsy patients had normal Apgar score at birth (2).

It has since become widely accepted that umbilical cord blood gas analysis can provide important information about the past, present and possibly the future condition of the infant. Umbilical cord blood gas analysis is now recommended in all high risk deliveries by both the British and American Colleges of Obstetrics and Gynecology (3, 4), and in some centers it is practiced routinely following all deliveries.

Low cord pH in infants who are vigorous at birth and free of cardiopulmonary compromise does not indicate an increased risk of adverse outcome (5). A good blood

gas in a depressed newborn should alert the neonatologist to search more diligently for other causes of neonatal depression like sepsis, trauma or congenital abnormalities (6).

Considerable effort has been spent to determine risk factors and the role of intrapartum asphyxia for adverse neonatal outcome (7). Based on different studies, mean values of umbilical cord blood pH range from 7.25-7.28 (8). A baby with abnormal umbilical cord pH does not always present with poor outcome.

2. Objectives

So this study was carried out to determine the short term outcome after birth in the neonates born with an abnormal umbilical cord pH.

3. Patients and Methods

This study is a prospective observational cohort study carried out at Rohani hospital, affiliated to Babol Univer-

sity of Medical Sciences in Babol, Mazandaran, Iran. The study was approved by the university ethic committee and informed consents were taken from the parents. All term neonates born between November 2011 and June 2012 by Cesarean Section (C/S) to a high risk mother based on neonatal resuscitation program (NRP) (9) entered the study; an umbilical cord arterial blood sample was taken according to the AAP protocol. All samples were sent in pre-heparinized syringes and prepared under aseptic condition and occupational safety. The samples were analyzed by ABG analyzer (Gem Premier, USA) 30 minutes after birth. Neonates with umbilical cord pH less than 7.2 were considered as cases and neonates with umbilical cord pH more than 7.2 as control group.

Data including pH, PO₂, PCO₂, BE and bicarbonate along with demographic data like maternal risk factor and age, gestational age, weight and sex of neonates were recorded by a questioner.

Inclusion criteria were term delivery to a high risk mother based on neonatal resuscitation program (NRP) schedule. Extraction criteria were prenatal infection and any major congenital anomaly. Considering $d = 20$ with an 80% power and 95% confidence interval, sample size was calculated 120 (60 mother-neonate pairs in each arm).

3.1. Outcome Variables

3.1.1. Need to Resuscitation

On the basis of NRP, we began to resuscitate neonates with meconium stained amniotic fluid, lack of spontaneous or effective respiratory effort and lack of appropriate muscular tone.

3.1.2. Need to Being Admitted to Neonatal Ward or NICU

Neonates who required positive pressure ventilation were admitted to newborn services and/or NICU.

3.1.3. Convulsion

Diagnosis of convulsion made by clinical observation by attending physician or nurse's reports. All neonates with convulsion were admitted and treated in neonatal ward or NICU.

3.1.4. Encephalopathy

Any asphyxiated neonate with neurologic signs and symptoms defined and classified based on the Sarnat and Sarnat staging system (10).

3.1.5. Delayed Start of Oral Feeding and/or Feeding Intolerance

Because of feeding intolerance or respiratory disorder, oral feeding started after stabilization. All neonates admitted to the NICU and/or SCN were kept NPO and IV fluid was started.

3.1.6. Hospital Stay

This period is calculated from birth to discharge and declared as days.

3.1.7. Death

Any cases of death in both groups were recorded.

3.2. Statistical Analysis

Data are analyzed by SPSS, version 18; X² test, Mann-Whitney and Spearman test were used. P less than 0.05 was considered as being significant.

4. Results

The baseline demographic data of the two groups are shown in Table 1. These variables did not show any significant differences between the two groups. Comparison of ABG parameters in two neonatal groups with normal and abnormal umbilical cord pH are as in Table 2. According to this table, differences between pH, PCO₂, PO₂, BE and bicarbonate were significant.

Comparison of short term outcomes including need to resuscitation, convulsion, encephalopathy and its severity, neonatal ward and NICU admission, delay in commencement of feeding and length of hospital stay clearly were better in non-acidemic neonates than in acidemic newborns (Table 3). Need to resuscitation including initial steps in acidemic neonates was higher than the non-acidemic neonates (42 vs. 13 neonates; $P = 0.001$) but there was no significant difference for PPV requirement (advanced resuscitation) between acidemic and non-acidemic neonates (4 vs. 0 neonates; $P < 0.119$).

One neonate needed chest compression, had cord arterial pH 6.99 and BD 12.7. This case stayed in NICU for 11 days. We had two cases of convulsion. First case had horizontal deviation of eyes along with apnea. Second case had tonic lower limb extension like decerebrated posture along with bradycardia. This case had paroxysmal tachycardia with hypertension. He received midazolam to control the seizure. Although in acidemic neonates we have 3.3% convulsive disorders but in non-acidemic newborns there were no cases of seizure. So in comparison with non-acidemic neonates there was no significant difference in seizure between the two groups ($P = 0.496$).

4.1. Hypoxic-Ischemic Encephalopathy

There were 21 cases of encephalopathy in this study. Signs of HIE often had been presented during first hours or days after birth. Some patients presented with mild signs such as decreased alertness, irregular respiratory efforts, and hypotonicity along with normal pupillary reflex.

Convulsive disorders as signs of moderate encephalopathy had been presented in two cases of our patients who had perinatal acidemia and in 24 to 48 hours after birth

their neurologic symptoms became remarkable.

In acidemic group 17 (28.3%) cases had encephalopathy but in nonacidemic group only 4 (6.7%) cases were encephalopathic with a calculated risk ratio of 4.25. So there was significant relationship between encephalopathy and perinatal acidemia ($P = 0.002$).

Severity of encephalopathy: Fifteen cases of grade 1 HIE occurred in acidemic group versus only 4 cases in non-acidemic group. On the other hand, in acidemic group we had two cases of grade 2 HIE but in non-acidemic group there were no cases of severe HIE. Also we found a linear relationship between the intensity of acidemia and severity of encephalopathy and all results of parametric tests were like nonparametric results ($P = 0.002$).

4.2. Death

There was no case of death in the two groups of the study.

4.3. Time of Oral Feeding Initiation

In acidemic group approximately half of the patients had oral feeding intolerance in the first hour after birth, 14 cases after two hours and 5 cases after four hours could tolerate oral feeding. These patients received D/W 10% IV and were transferred to the ward. Commencement and Tolerance of Oral Feeding in Patients With Abnormal Umbilical Cord pH has been shown in Figure 1.

4.4. Delay in Oral Feeding Initiation

In acidemic group we commenced oral feeding for 6 cases at 24 hours of life and for 3 cases at 48 hours of life. One case had oral feeding intolerance until 5 days after birth. Finally 31 (51.7%) cases had oral feeding and 29 (48.3%) cases had oral feeding intolerance in the first hour of life.

In non-acidemic group 4 neonates had begun feeding at 2, and 2 cases after 4 hours of life, and 3 neonates gradually begun breast feeding after 48 hours and in one case at 96 hours of life. Finally in 50 (83.3%) cases of newborns in non-acidemic group oral feeding had begun in the first hour of life (Figures 2 and 3). Also Duration of NICU and Neonatal Ward Stay in Non-Acidemic Neonates has been shown in Figure 4.

4.5. Hospital Stay

Mean hospital stay in acidemic neonates was 3.78 ± 3.07 days. In acidemic neonates the length of hospital stay had reverse relationship with umbilical cord pH ($r = -0.363$, $P < 0.001$), NaHCO_3 ($r = -0.304$, $P < 0.001$) and BE ($r = -0.456$, $P < 0.001$) whereas there was no linear relationship with PO_2 . The result of nonparametric tests was the same as the parametric tests (Figures 3 and 4).

4.6. Risk Factors

Meconium stained amniotic fluid was the most common risk factor which has been seen in both groups. Premature rupture of membranes (PROM) was the second most common risk factor in our study.




Figure 1. Commencement and Tolerance of Oral Feeding in Patients With Abnormal Umbilical Cord pH




Figure 2. Commencement and Tolerance of Oral Feeding in Neonates With Normal Umbilical Cord pH




Figure 3. Duration of NICU and Neonatal Ward Stay in Acidemic Neonates




Figure 4. Duration of NICU and Neonatal Ward Stay in Non-Acidemic Neonates

Table 1. Baseline Demographic Data in Abnormal and Normal Umbilical Cord pH Groups

Variables	Abnormal pH Group ^a	Normal pH Group ^a	P value
Gender^b			0.144
Female	34 (56.6)	25 (41.7)	
Male	26 (43.3)	35 (58.3)	
Weight^c, gr	32640 ± 554	3204 ± 370	0.622
Gestational age^c, week	39 ± 1	38 ± 1	0.549
Maternal age^c, year	28.8 ± 2	25 ± 4.2	0.001

^a Data are presented for n = 60.^b Data are presented as No. (%).^c Data are presented as Mean ± SD.**Table 2.** Comparison of ABG Parameters in the Two Neonatal Groups With Normal and Abnormal Cord ABG^a

Parameters	Abnormal pH Group ^b	Normal pH Group ^b	P value
pH	7.135 ± 0.066	7.284 ± 0.037	> 0.001
PO², Torr	11 ± 6	14 ± 6	0.003
PCO², Torr	63.7 ± 13.7	52.1 ± 7.9	0.001
BE, mmol	-8.34 ± 3.57	-2.48 ± 2.34	0.001
Bicarbonate, mEq	21.36 ± 3.54	24.13 ± 3.25	0.001

^a Data are presented for n = 60.^b Data are presented as Mean ± SD.**Table 3.** Comparison of Short Term Outcome Between Abnormal and Normal Umbilical Cord pH Groups

Outcome Variables	Abnormal pH Group ^a	Normal pH Group ^a	P value
Need for resuscitation	42 (70)	13 (21.7)	< 0.001
Need for PPV	4 (6.7)	0 (0)	0.119
Convulsion	2 (3.3)	0 (0)	0.496
HIE	17 (28.3)	4 (6.7)	0.002
Need for NICU admission	16 (26.7)	3 (5)	0.001
Delays in oral feeding initiation until an hour	29 (48.3)	10 (17.7)	0.012
Hospital stay	3.78 ± 3.07 ^b	2.28 ± 0.98 ^b	< 0.001

^a Data are presented for n = 3 and as No. (%).^b Data are presented as Mean ± SD.

5. Discussion

Our study which has been carried out for comparison of short outcomes in normal and abnormal umbilical cord pH groups, showed that short outcomes including need to resuscitation, encephalopathy, convulsion, NICU and neonatal ward admission and length of hospital stay, delay in commencement of oral feeding in abnormal umbilical cord pH group are more common than in normal umbilical cord pH group (11).

Andres et al. assessed ninety three neonates for the correlation between the umbilical cord PCO², PO², pH, bicarbonate and BD with death, need to cardiopulmonary resuscitation, convulsion, RDS, PDA, NEC, IUGR, sepsis and HIE. Although in their study there was no correlation between sepsis, RDS, NEC, PDA and UABG parameters,

need to cardiopulmonary resuscitation, need to intubation, convulsion, HIE and IUGR were correlated with low UABG pH (11). In their study PO² had no correlation with outcomes. In our study, abnormal UABG pH had no significant correlation with convulsion.

Goldaber and colleagues studied the association between umbilical arterial acidosis and neurologic complications (encephalopathy and convulsion) among 3506 term, singleton neonates with cord arterial pH < 7.20 (12). Neonatal death was much more likely at pH < 7.00. The cutoff value at which seizures became more likely was pH < 7.05, and for unexplained seizures pH < 7.00. They suggested that a realistic value for defining pathological acidemia was pH < 7.00.

In Williams et al. study, neonatal seizure was predicted only by low umbilical artery pH. A pH of less than 7 was more (73.8%) sensitive than a base excess of -16 (52.5%) in predicting the development of neonatal seizures (13). In our study only two cases of the acidemic neonates developed seizure both of whom had pH 6.99, 7.08 and base excess of -12.7 and -9.2.

Victory et al. found that there is a progression of risk in term infants for NICU admission and need for assisted ventilation with worsening acidosis at birth (7).

Not all neonates with acidosis at birth require resuscitation or have adverse outcome. In the present study 60 percent did not require admission. King et al. concluded that infants with an umbilical artery pH \leq 7.0 and assessed to be clinically well can be treated similar to non-acidemic infants (14). The cut off value for pH in our study is somewhat higher than in King's study.

In a prospective study performed by Loh and his colleagues on umbilical cord ABG analysis in labor ward on 200 neonates with purpose to establish the normal range of umbilical cord blood gas values, only 5 cases had umbilical cord pH under 7.05. An important point in this study was correlation between the significant metabolic acidosis compared to respiratory acidosis and poor outcome like prolonged stay in NICU and death (6). The poorest outcome in our study belonged to a baby who had received PPV at the time of birth and had pH less than 6.99 and base excess of -12.7.

Malin and her colleagues in their study (2010) declared that umbilical cord pH less than 7.2 was an appropriate prognostic agent for presenting asphyxia and other short outcomes. In their systematic review they showed that low arterial cord pH was significantly associated with perinatal and long term neonatal outcomes like mortality, hypoxic ischemic encephalopathy, intraventricular hemorrhage or periventricular leucomalacia and cerebral palsy (15).

Williams and Singh (2004) studied short term outcomes in 47 acidemic neonates. Early convulsion as most clear short outcome had been seen in moderate to severe encephalopathy in their study. Interestingly upper PO₂ in their patients corresponded with poorer short outcomes, so they took PO₂ for a poor prognostic agent. Our study didn't establish this opinion. In their study there was no correlation with pH, BD, HIE and convulsive disorder (16). These findings differ from those of ours and majority of researchers.

Bekebam and his colleagues in 2002 with the purpose of evaluating incidence of prenatal acidemia carried out a prospective cohort study. They showed that perinatal acidemia was more common in male neonates ($P < 0.0001$) and male neonates were more prone to fetal distress and low Apgar score (17). This finding corresponds with our study, which showed mild predominance of perinatal acidemia incidence in male newborns.

One of limitations which may be attributed to our study may be related to study population, all of whom were

mother-baby pairs from cesarean section group. The umbilical cord pH value may differ between cesarean section and normal vaginal delivery groups as indicated by Loh et al. In their study, the mean and standard deviation values of umbilical artery pH were 7.21 and 0.08 for vaginal deliveries, and 7.22 and 0.07 for caesarean sections, respectively (6).

The results of present study showed that short term outcomes like need for advanced resuscitation, NICU admission, HIE, delay to start oral feeding, mean hospital stay were higher in acidemic in comparison to non-acidemic neonates. Therefore, an umbilical cord PH less than 7.2 immediately after birth can be used as a prognostic factor for prediction of unfavorable short term outcomes in newborns.

Acknowledgements

We would like to thank Dr Firozjahi for his laboratory support.

Funding/Support

This study was funded by Research deputy of Babol University of Medical Sciences (Proposal No: 1294).

References

- White CR, Doherty DA, Henderson JJ, Kohan R, Newnham JP, Pennell CE. Benefits of introducing universal umbilical cord blood gas and lactate analysis into an obstetric unit. *Aust N Z J Obstet Gynaecol.* 2010;**50**(4):318-28.
- Vukojevic M, Soldo I, Granic D. Risk factors associated with cerebral palsy in newborns. *Coll Antropol.* 2009;**33** Suppl 2:199-201.
- Royal College of Obstetricians and Gynaecologists. The use of electronic fetal monitoring: the use and interpretation of cardiotocography in intrapartum fetal surveillance. In: National Institute for Health and Care Excellence, editor. London: 2001.
- Acog Committee on Obstetric Practice. ACOG Committee Opinion No. 348, November 2006: Umbilical cord blood gas and acid-base analysis. *Obstet Gynecol.* 2006;**108**(5):1319-22.
- Armstrong L, Stenson BJ. Use of umbilical cord blood gas analysis in the assessment of the newborn. *Arch Dis Child Fetal Neonatal Ed.* 2007;**92**(6):F430-4.
- Loh SF, Woodworth A, Yeo GS. Umbilical cord blood gas analysis at delivery. *Singapore Med J.* 1998;**39**(4):151-5.
- Victory R, Penava D, Da Silva O, Natale R, Richardson B. Umbilical cord pH and base excess values in relation to adverse outcome events for infants delivering at term. *Am J Obstet Gynecol.* 2004;**191**(6):2021-8.
- Ip FF, di Pierro I, Brown R, Cunningham I, Handelsman DJ, Liu PY. Trough serum testosterone predicts the development of polycythemia in hypogonadal men treated for up to 21 years with subcutaneous testosterone pellets. *Eur J Endocrinol.* 2010;**162**(2):385-90.
- Kattwinkel J. *NRP Neonatal Resuscitation Textbook 6th Edition (English version)*. 6th Edition ed: American Academy of Pediatrics and American Heart Association Guidelines for Neonatal Resuscitation; 2011.
- Levene MI, DeVries LS. Hypoxic-ischemic encephalopathy. In: Martin RJ, Fanaroff AA, Walsh MC editors. *Fanaroff and Martin's Neonatal-Perinatal Medicine: Diseases of the Fetus and Infant.*: Elsevier Health Sciences; 2010.
- Andres RL, Saade G, Gilstrap LC, Wilkins I, Witlin A, Zlatnik F, et al. Association between umbilical blood gas parameters and neonatal morbidity and death in neonates with pathologic fetal acidemia. *Am J Obstet Gynecol.* 1999;**181**(4):867-71.
- Goldaber KG, Gilstrap L3, Leveno KJ, Dax JS, McIntire DD. Patho-

- logic fetal acidemia. *Obstet Gynecol.* 1991;**78**(6):1103-7.
13. Williams KP, Singh A. The correlation of seizures in newborn infants with significant acidosis at birth with umbilical artery cord gas values. *Obstet Gynecol.* 2002;**100**(3):557-60.
 14. King TA, Jackson GL, Josey AS, Vedro DA, Hawkins H, Burton KM, et al. The effect of profound umbilical artery acidemia in term neonates admitted to a newborn nursery. *J Pediatr.* 1998;**132**(4):624-9.
 15. Malin GL, Morris RK, Khan KS. Strength of association between umbilical cord pH and perinatal and long term outcomes: systematic review and meta-analysis. *BMJ.* 2010;**340**:c1471.
 16. Williams KP, Singh A. Umbilical artery blood gas parameters in neonates with early onset seizures who die. *BJOG.* 2004;**111**(10):1042-5.
 17. Bekedam DJ, Engelsbel S, Mol BW, Buitendijk SE, van der Pal-de Bruin KM. Male predominance in fetal distress during labor. *Am J Obstet Gynecol.* 2002;**187**(6):1605-7.

Archive of SID