Published online 2015 August 24.

Research Article

Bioelectric Impedance Analysis in the Diagnosis of Vesicoureteral Reflux

Meral Torun Bayram^{1,*}; Demet Alaygut¹; Mehmet Turkmen¹; Alper Soylu¹; Salih Kavukcu¹

¹Department of Pediatrics, Medical Faculty, Dokuz Eylul University, Izmir, Turkey

*Corresponding author: Meral Torun Bayram, Department of Pediatrics, Medical Faculty, Dokuz Eylul University, Izmir, Turkey. Tel: +90-2324123624, E-mail: meralt.bayram@yahoo.com

Received: August 1, 2014; Accepted: April 16, 2015

Background: Vesicoureteral reflux (VUR) is a common abnormality of the urinary tract in childhood.

Objectives: As urine enters the ureters and renal pelvis during voiding in vesicoureteral reflux (VUR), we hypothesized that change in body water composition before and after voiding may be less different in children with VUR.

Patients and Methods: Patients were grouped as those with VUR (Group 1) and without VUR (Group 2). Bioelectric impedance analysis was performed before and after voiding, and third space fluid (TSF) (L), percent of total body fluid (TBF%), extracellular fluid (ECF%), and intracellular fluid (ICF%) were recorded. After change of TSF, TBF, ECF, ICF (ΔTSF, ΔTBF%, ΔECF%, ΔICF%), urine volume(mL), and urine volume/ body weight (mL/kg) were calculated. Groups 1 and 2 were compared for these parameters. In addition, pre- and post-voiding body fluid values were compared in each group.

Results: TBF%, ECF%, ICF%, and TSF in both pre- and post-voiding states and Δ TBF%, Δ ECF%, Δ ICF%, and Δ TSF after voiding were not different between groups. However, while post-voiding TBF%, ECF% was significantly decreased in Group 1 (64.5 ± 8.1 vs 63.7 ± 7.2, P = 0.013 for TBF%), there was not post-voiding change in TSF in the same group. On the other hand, there was also a significant TSF decrease in Group 2. **Conclusions:** Bladder and ureter can be considered as the third space. Thus, we think that BIA has been useful in discriminating children with VUR as there was no decreased in patients with VUR, although there was decreased TSF in patients without VUR. However, further studies are needed to increase the accuracy of this hypothesis.

Keywords: Impedance; Vesicoureteral Reflux; Children

1. Background

Vesicoureteral reflux (VUR) is a common abnormality of the urinary tract in childhood. Diagnostic imaging of VUR is one of the most frequently used procedures in pediatric radiology departments all over the world. Voiding cystourethrography (VCUG) and radionuclide voiding cystography (RNC) are most commonly two methods used to diagnose VUR (1). However, there are three main problems associated with VCUG and RNC, including sedation, urethral catheterization, and ionizing radiation (2). Sedation issues come up regularly and some institutions use sedation in most of their bladder-imaged patients. Another problem is that urethral catheter is not well tolerated by children or their parents. Dysuria for a day or two almost always follows the catheterization. Stricture disease, UTI, and hematuria may result from catheter (2). Third problem is radiation exposure associated with VCUG and RNC. There is significant concern about radiation exposure in children in long term and cancer development. In children, the risk is not only associated with the amount of radiation exposure but how much time has to pass till it has to bring about effects. Because infants and children have a full life expectancy ahead of them, the overall risk is greater than in adults (2). Because of the unpleasant nature of the radiographic imaging and radiation exposure, significant efforts have been undertaken to utilize non-invasive techniques for diagnosis of VUR.

Bioelectric impedance analysis (BIA) is an accurate, non-invasive, and inexpensive technique for quantitative estimating of body water compartments in both healthy individuals and those affected by pathologic conditions (3, 4). The technique is based on the measurement of the resistance generated in the body against a low voltage electrical current. It is possible to estimate the TBF and to separate the ECF and ICF volumes by using various frequencies (5). Previously BIA has been used to assess body water changes in patients with renal failure, ascites, acute dehydration, in the postoperative period, and congestive heart failure (6-10). However, the role of BIA in diagnosis of VUR is not yet defined. Because VUR is the retrograde flow of urine from bladder to kidneys (11), body water in children with VUR after voiding may show small change.

2. Objectives

Therefore, we hypothesized that change in body water composition before and after voiding may be different in children with VUR compared to those without VUR using BIA. Thus, in this study, we have investigated whether BIA can be used as a method in diagnosis of VUR.

Copyright © 2015, Growth & Development Research Center. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non-Commercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

3. Patients and Methods

3.1. Patients and Control Cases

This study was conducted in our institution between March 2011 and September 2012. Children with continence who had VCUG (in the last three months) to diagnose VUR were included in the study. Any patients who had a history of heart disease, chronic kidney disease, anatomic obstruction (e.g. urethral valves or ectopic ureterocele), post-void residual urine volume (PVR) greater than 10 mL, urinary incontinence, or active urinary tract infection were excluded from study. The patients and control cases were evaluated with respect to present age, gender, VCUG, 99 mTc-DMSA renal scintigraphy and multifrequency bioelectrical impedance analysis (MF-BIA).

The patients were grouped as VUR (+) (Group 1) and VUR (-) (Group 2 = control group) with respect to the presence of VUR in VCUG. VUR has been graded from grade 1 to 5 according to the grading system of international reflux study committee (12). Therefore, Group 2 was grouped as low-grade and high-grade VUR. First- and second-grade reflux was considered as low grade and third to fifth grades as high-grade reflux. The study was approved by the local ethical committee of our institution. All patients and controls signed the informed consent for participation in the study.

3.2. Anthropometric Measurements

Body weight was determined on a SECA balance scale (Hamburg, Germany) to the nearest 0.1 kg, with subjects dressed in a light t-shirt and shorts. Body height of the patients was recorded on a stadiometer with a standing subject to the nearest 0.1 cm with a Harpenden fixed stadiometer (Holtain Ltd., Crymych, Dyfed, Britain). Body mass index was calculated as weight/height² (kg/m²).

3.3. Multifrequency Bioelectric Impedance Analysis

Body composition and body water compartment analysis were carried out with the subject lying in a supine position on a flat, nonconductive bed with MF-BIA. MF-BIA was done between 8 and 10 in the morning to avoid bias due to diurnal variation with a Bodystat Qudscan 4000 bioimpedance analyzer (Bodystat Limited, British Isles). All patients were evaluated two times after an overnight fasting, firstly with full bladder and secondly after voiding. Current-detecting electrodes were placed between the styloid processes of the right radius and ulna and between the medial and lateral malleoli of the right ankle. Current-introducing electrodes were then placed on the respective dorsal surfaces of the metacarpals and metatarsals, 5 cm distal to the proximal electrodes (standart tetrapolar placement). A minimal inter-electrode distance of 5 cm has been recommended to prevent interaction between electrodes (13, 14). Later, body weight and height of patient were put into the analyzer. By the MF-BIA, an alternating electrical current of constant frequency with low intensity was applied to the body.

Body mass index (BMI), fat mass index (FMI), fat-free mass index (FFMI), percent of total body fluid (TBF%), percent of extracellular fluid (ECF%), percent of intracellular fluid (ICF%), third space fluid (TSF) (L), impedance (a combination of resistance and reactance), resistance (R; i.e. opposition to the flow of an alternating current through intra- and extracelluar ionic solution), reactance (Xc; i.e. the capacitative component of tissue interfaces, cell membranes and organelles), and phase angle (PA; Xc/R) were calculated by using the manufacturer's software. These measurements were performed before and after voiding.

3.4. Calculations

Urine volume (mL) of all cases was calculated after voiding. Later, urine volume was divided by body weight (kg) and so volume/body weight (mL/kg) was determined. Therefore, the changes (liter or percent) in TSF (Δ TSF), TBF (Δ TBF%), ECF (Δ ECF%), ICF (Δ ICF%) were calculated by the formula as TSF, TBF, ECF, and ICF after voiding minus TSF, TBF, ECF, ICF before voiding, then divided by the before voiding TSF, TBF, ECF, and ICF in each patient. For example, one case had his before voiding ECF as 48%, and after voiding ECF as 45%, then the DECF% is 45 - 48/48 = 1.95%. All these parameters were compared between Group 1 and 2.

In addition, impedance (ohm) of a cylindrical conductor, in fact, is proportional to its specific impedivity and to its length. Therefore, resistance/height (R/H) and reactance/height (Xc/H) were also calculated.

3.5. Statistical Analysis

All results are expressed as mean \pm standard deviation unless otherwise specified. The data between groups were compared with Mann-Whitney U, Chi-Square test, paired sample t test, and Wilcoxon signed-ranks tests. Correlation between the grade of VUR and changes of body fluid was assessed using Pearson correlation. A P value < 0.05 was considered statistically significant. Statistical analyses were performed using the scientific package for social science 16.0.1 (SPSS Inc., Chicago, IL, USA).

4. Results

A total of 78 (25 males) patients were included in the study. There are 52 patients in Group 1 (with VUR) and 26 patients in Group 2 (without VUR). In Group 1, 25 (48%) patients had severe VUR (grade \geq 3) and 23 (44%) patients had bilateral VUR. Table 1 shows the demographic characteristics and body composition: age, sex, height, weight, BMI, FMI, and FFMI are similar in Groups 1 and 2.

Data on body fluid composition obtained by MF-BIA pre- and post-voiding are shown in Table 2. In Group 1, differences in pre- and post-voiding values of TBF%, ECF%, impedance, R, and R/H, showed statistically significant differences. In Group 2, differences in pre- and

post-voiding values of ECF%, TSF in L, PA, Xc, and Xc/H, showed statistically significant differences.

Pre- and post-voiding TSF in L, TBF%, ECF%, and ICF% values by MF-BIA and the ΔTSF%, ΔTBF%, ΔECF%, and ΔICF% were not different between Groups 1 and 2 (Table 3). Also, there was no difference of same parameters between patients with high-grade VUR and other patients (data not shown).

By analyzing the relationship of VUR grade with changes of body fluid after voiding, there was only a significant positive correlation between the VUR grade and R/H (Table 4).

Table 1. Demographic Characteristics and Body Composition in Group 1 and Group 2 ^a					
General Characteristics	Group 1, (With VUR), n = 52	Group 2, (Without VUR), n = 26	P Value		
Age, mo	74.1±35.6	76.6 ± 30.1	0.637		
Gender			0.074		
Male	13	12			
Female	39	14			
Renal scarring	37/12	4/12	0.001		
Weight, kg	22.3 ± 10.3	24.6 ± 8.4	0.117		
Height, cm	114.7 ± 19.5	119.4 ± 15.7	0.203		
BMI, kg/m ²	15.9 ± 2.9	16.8±2.2	0.071		
FMI, kg/m ²	4.2 ± 3.5	4.5 ± 2.5	0.316		
FFMI, kg/m ²	13.1±1.7	13.7±1.6	0.171		
a Abbraviations: PML body mass index. FEML fat free mass index. FML fat mass index. MLP. vesicouverant reflux					

^a Abbreviations: BMI, body mass index, FFMI, fat-free mass index; FMI, fat mass index; VUR, vesicoureteral reflux

Table 2. Pre- and Post-Voiding Distribution of Body Fluids and Body Weight in Group 1 and Group 2

	Group 1			Group 2		
	Pre-Voiding	Post-Voiding	P Value	Pre-voiding	Post-Voiding	P Value
Weight	22.3 ± 10.3	22.2 ± 10.2	0.067	24.6 ± 8.4	24.6 ± 8.3	0.414
TBF, %	64.5 ± 8.1	63.7 ± 7.2	0.013	63.8 ± 6.2	63.6 ± 6.2	0.125
ECF, %	42.6 ± 8.9	42.4 ± 8.8	0.001	39.6 ± 6.7	39.2 ± 6.4	0.015
ICF, %	40.5 ± 9.8	40.8 ± 10.2	0.415	40.8 ± 10.3	42.1 ± 10.7	0.479
TSF, L	-3.0 ± 2.0	-3.2 ± 2.1	0.077	-3.3 ± 2.2	-3.5 ± 2.2	0.013
Imp.50kHz	662.7 ± 90.2	670.4 ± 86.4	0.003	638.3 ± 65.4	641.6 ± 68.1	0.074
PA	4.7 ± 0.9	4.7 ± 0.7	0.620	4.9 ± 6.9	5.1 ± 0.8	0.010
R, Ω	660.4 ± 90.3	668.0 ± 86.5	0.004	635.9 ± 65.3	639.2 ± 68.2	0.080
Χc, Ω	45.3 ± 9.1	54.9 ± 9.3	0.053	54.3 ± 8.2	56.3 ± 8.1	0.001
R/H, Ω/ m	592.1 ± 127.8	599.9 ± 129.5	0.006	542.7 ± 94.5	545.8 ± 98.5	0.065
$\mathbf{Xc}/\mathbf{H}, \Omega/\mathbf{m}$	48.2 ± 9.6	$\textbf{48.7} \pm \textbf{9.4}$	0.126	45.9 ± 6.9	47.7 ± 7.8	0.002

^a Abbreviations: ECF, extracellular fluid; ICF, intracellular fluid; PA, phase angle; R, resistance at 50 kHz; R/H, resistance/height; TBF, total body fluid; TSF, third space fluid; Xc, reactance at 50 kHz; Xc/H, reactance/height.

Table 3. Pre-Voiding and Post-Voiding Distribution of Body Fluids and Urine Volume in Group 1 and Group 2					
	Group 1	Group 2	P Value		
Total Body Fluid (TBF), %					
Prevoiding TBF	64.5 ± 8.1	63.8 ± 6.2	0.845		
Postvoiding TBF	63.7 ± 7.2	63.6 ± 6.2	0.890		
ΔTBF	-0.011 ± 0.027	-0.004 ± 0.0124	0.450		
Extracellular Fluid (ECF), %					
Prevoiding TBF	42.6 ± 8.9	39.6 ± 6.7	0.155		
Postvoiding TBF	42.4 ± 8.8	39.2 ± 6.4	0.108		
ΔTBF	-0.004 ± 0.007	-0.009 ± 0.037	0.928		
Intracelluler Fluid (ICF), %					
Prevoiding TBF	40.5 ± 9.8	40.8 ± 10.3	0.853		
Postvoiding TBF	40.8 ± 10.2	42.1 ± 10.7	0.715		
ΔTBF	-0.009 ± 0.077	0.039 ± 0.134	0.746		
Third space Fluid (TSF), L					
Prevoiding TBF	-3.4 ± 2.0	-3.3 ± 2.2	0.393		
Postvoiding TBF	-3.2 ± 2.1	-3.5 ± 2.2	0.370		
ΔTBF	0.079 ± 0.296	$0,095 \pm 0.266$	0.765		
Urine Volume/Body Weight, mL	5.27 ± 4.33	5.59 ± 3.89	0.436		

Table 4. Relationship Between Grade of VUR With Changes of Body Fluid After Voiding

Measure	Grade of VUR		
	r	P Value	
ΔΤΒF, %	-0.097	0.402	
ΔΕCF, %	-0.114	0.323	
ΔICF, %	-0.093	0.419	
ΔTSF, L	-0.037	0.750	
∆Imp, 50 kHz	0.043	0.712	
∆Phase Angle	-0.218	0.057	
$\mathbf{R}/\mathbf{H}, \Omega/\mathbf{m}$	0.327	0.004	
$\mathbf{Xc}/\mathbf{H}, \Omega/\mathbf{m}$	0.134	0.247	

^a Abbreviations: Δ ECF, change in ECF after voiding; Δ ICF, change in ICF after voiding; Δ TBF, change in TBF after voiding; Δ TSF, change in TSF after voiding.

5. Discussion

Diagnostic methods used for the detection of VUR according to their characteristics can be categorized as direct/indirect, catheter-using/catheter-free, radiation-giving/radiation-free. Ideally, the diagnostic methods used for the diagnosis of VUR should be a safe, radiation-free, non-invasive, low-cost, high-sensitivity imaging method (2, 15). However, none of the current diagnostic methods include most of these criteria. Therefore, new diagnostic methods for the diagnosis of VUR must be investigated.

There are reference methods for measuring body fluid volumes such as isotope dilution techniques, total body nitrogen, and densitometry for TBF, bromide for ECF and radioactive potassium isotope for ICF (16-20). However, these methods are expensive and cumbersome, and expose children to radiation or invasive procedures (16-19, 21), and the use of these methods is limited in clinical practice (22). Approaches to overcome these restrictions are single-frequency BIA (SF-BIA), multi-frequency BIA (MF-BIA), and bioelectric impedance spectroscopy (BIS).

SF-BIA, MF-BIA, and BIS are used for estimating of TBF, ECF, ICF, and body composition on the basis of mathematical formulae using measurements of resistance, reactance, and impedance (23). These methods provide evaluation of body water compartments of healthy subjects and those affected by pathological situations (3, 4, 24). Among BIA techniques, MF-BIA seems to be a more accurate method for estimating the TBF compartment (24, 25). A meta-analysis showed that TBF in healthy individuals was significantly overestimated by SF-BIA or BIS in comparison with the reference values obtained using D₂O dilution. However, those studies that used MF-BIA only did not overestimate the TBF (24). The use of BIA as a bedside method has increased because the equipment is portable and safe, the procedure is simple and noninvasive, and the results are reproducible and rapidly obtained (25). BIA is also a painless method (26). The main advantage of this method, in patients with alterations in water metabolism, is that it works independently of the body weight. Because VUR is the retrograde flow of urine from bladder to kidneys (11), change after voiding in body water composition, especially TBF and TSF, might be different between children with and without VUR. In this study, we wanted to confirm this hypothesis and chose MF-BIA due to above-mentioned assumptions. Thus, we investigated whether a BIA can be used as a method in diagnosis of VUR.

Body fluid compartments may be affected by body composition. For example, TBF is strongly related to FFMI, similarly, body cell mass, which is an important nutritional parameter, is also closely connected to ICF (27, 28). Therefore, the differences in body compartments prevent correct evaluation of body water compartments. In our study, body weight and height, BMI, FFM, and FFMI were similar between patients with and without VUR. In addition, the urine volume and urine volume/body weight did not differ between groups. Although there was no difference in body composition affecting body fluid between patients with and without VUR, pre-and post-voiding TSF (L), TBF%, ECF%, and ICF% were not different between these groups. In addition, the changes (% or L) in TSF, TBF, ECF, and ICF post-voiding were similar between same groups. However, when the groups were examined separately, post-voiding TBF% and ECF% was found to be lower than the pre-voiding TBF% and ECF% in Group 1. Nevertheless, TSF was not different

In addition to the above results, the difference in impedance, R, R/H, Xc, and Xc/H values after voiding were severally determined in both groups at the same time. However, these values are pitfalls of conventional BIA (29). In previous studies it was obs. On the other hand, post-voiding TSF value was determined lower than prevoiding TSF value in Group 2. Urine in bladder and ureter is probably TSF and VUR is the retrograde flow of urine from the bladder to the kidneys (11). Therefore, change of TSF after voiding was likely found to be less in children with VUR.erved that R and R/H were strongly correlated with TBF, whereas Xc and Xc/H were more strongly related to ECF (30, 31). In this study, we also similarly determined decrease in TBF% and increase in R, after voiding among patients with VUR and decrease in ECF% and increase in Xc after voiding among patients without VUR.

VUR is associated with two related consequences: urinary tract infection and renal scarring. The management of VUR is based on preventing these sequelae. Renal scar causes secondary hypertension and chronic renal failure. One of the risk factors for renal scar formation is higher grade VUR (32). Also, prophylaxis in VUR is recommended for high-grade VUR. For these reasons, we also examined the availability of the BIA method in the diagnosis of high-grade VUR. However, there were no differences in TBF and TSF between patients with and without high-grade VUR in pre- and post-voiding states. In addition, there was no relationship between changes in TBF and TSF after voiding. Moreover, when the relationship between body fluid changes after voiding with grade of VUR was examined, a significant correlation was not determined except for R/H.

The main disadvantages of BIA in diagnosis of VUR are limited visualization of the urethra, and inadequacy in diagnosis of patients with passive VUR. Despite these disadvantages, we think that BIA is a valid alternative to conventional VCUG or RC in a screening population of girls and in follow-up. However, for the present, we could not exactly determine that BIA is an alternative to conventional VCUG or RC in VUR diagnosis. On the other hand, we found that while post-voiding TSF value was lower than pre-voiding TSF value in patients without VUR, there was no difference between these values in patients with VUR and this result suggests having need for further studies with more patients.

References

- Kis E, Nyitrai A, Varkonyi I, Mattyus I, Cseprekal O, Reusz G, et al. Voiding urosonography with second-generation contrast agent versus voiding cystourethrography. *Pediatr Nephrol.* 2010;25(11):2289-93.
- Snow BW, Taylor MB. Non-invasive vesicoureteral reflux imaging. J Pediatr Urol. 2010;6(6):543–9.
- Gudivaka R, Schoeller DA, Kushner RF, Bolt MJ. Single- and multifrequency models for bioelectrical impedance analysis of body water compartments. J Appl Physiol (1985). 1999;87(3):1087-96.
- Woodrow G, Oldroyd B, Turney JH, Davies PS, Day JM, Smith MA. Measurement of total body water by bioelectrical impedance in chronic renal failure. *Eur J Clin Nutr.* 1996;**50**(10):676–81.
- Dewit O, Ward L, Middleton SJ, Watson C, Friend PJ, Elia M. Multiple frequency bioimpedance: a bed-side technique for assessment of fluid shift patterns in a patient with severe dehydration. *Clin Nutr.* 1997;16(4):189–92.
- Mandolfo S, Farina M, Imbasciati E. Bioelectrical impedance and hemodialysis. Int J Artif Organs. 1995;18(11):700-4.
- Arkouche W, Fouque D, Pachiaudi C, Normand S, Laville M, Delawari E, et al. Total body water and body composition in chronic peritoneal dialysis patients. J Am Soc Nephrol. 1997;8(12):1906-14.
- Dal Cin S, Braga M, Molinari M, Cristallo M, Di Carlo V. Role of bioelectrical impedance analysis in acutely dehydrated subjects. *Clin Nutr.* 1992;11(3):128-33.
- Chiolero RL, Gay LJ, Cotting J, Gurtner C, Schutz Y. Assessment of changes in body water by bioimpedance in acutely ill surgical patients. *Intensive Care Med.* 1992;18(6):322–6.
- 10. Soderberg M, Hahn RG, Cederholm T. Bioelectric impedance analysis of acute body water changes in congestive heart failure. *Scand J Clin Lab Invest.* 2001;**61**(2):89–94.
- Mattoo TK, Mathews R. Vesicoureteral Reflux and Renal Scarring. In: Avner ED, Harmon WE, Niaudet P, Yoshikawa N, editors. *Pediatric Nephrology*. 6th ed. Springer-Verlag Berlin Heidelberg; 2009. pp. 1311–36.
- Medical versus surgical treatment of primary vesicoureteral reflux: a prospective international reflux study in children. *J Urol.* 1981;**125**(3):277–83.
- 13. De Palo T, Messina G, Edefonti A, Perfumo F, Pisanello L, Peruzzi L, et al. Normal values of the bioelectrical impedance vector in child-

hood and puberty. Nutrition. 2000;16(6):417-24.

- Piccoli A, Fanos V, Peruzzi L, Schena S, Pizzini C, Borgione S, et al. Reference values of the bioelectrical impedance vector in neonates in the first week after birth. *Nutrition*. 2002;18(5):383–7.
- 15. O'Hara SM. Vesicoureteral reflux: latest option for evaluation in children. *Radiology*. 2001;**221**(2):283–4.
- Gutin B, Litaker M, Islam S, Manos T, Smith C, Treiber F. Body-composition measurement in 9-11-y-old children by dual-energy Xray absorptiometry, skinfold-thickness measurements, and bioimpedance analysis. Am J Clin Nutr. 1996;63(3):287-92.
- Boot AM, Bouquet J, de Ridder MA, Krenning EP, de Muinck Keizer-Schrama SM. Determinants of body composition measured by dual-energy X-ray absorptiometry in Dutch children and adolescents. *Am J Clin Nutr.* 1997;66(2):232–8.
- Wells JC, Fewtrell MS, Davies PS, Williams JE, Coward WA, Cole TJ. Prediction of total body water in infants and children. Arch Dis Child. 2005;90(9):965–71.
- Foster BJ, Leonard MB. Measuring nutritional status in children with chronic kidney disease. *AmJ Clin Nutr.* 2004;80(4):801-14.
- Jaffrin MY, Morel H. Body fluid volumes measurements by impedance: A review of bioimpedance spectroscopy (BIS) and bioimpedance analysis (BIA) methods. *Med Eng Phys.* 2008;30(10):1257-69.
- Ellis KJ, Wong WW. Human hydrometry: comparison of multifrequency bioelectrical impedance with 2H2O and bromine dilution. J Appl Physiol (1985). 1998;85(3):1056–62.
- 22. Bozzetto S, Piccoli A, Montini G. Bioelectrical impedance vector analysis to evaluate relative hydration status. *Pediatr Nephrol.* 2010;**25**(2):329–34.
- De Lorenzo A, Andreoli A, Matthie J, Withers P. Predicting body cell mass with bioimpedance by using theoretical methods: a technological review. J Appl Physiol (1985). 1997;82(5):1542-58.
- 24. Martinoli R, Mohamed EI, Maiolo C, Cianci R, Denoth F, Salvadori S, et al. Total body water estimation using bioelectrical impedance: a meta-analysis of the data available in the literature. *Acta Diabetol.* 2003;**40 Suppl 1**:S203-6.
- 25. Kyle UG, Bosaeus I, De Lorenzo AD, Deurenberg P, Elia M, Gomez JM, et al. Bioelectrical impedance analysis--part I: review of principles and methods. *Clin Nutr.* 2004;**23**(5):1226-43.
- Tyrrell VJ, Richards G, Hofman P, Gillies GF, Robinson E, Cutfield WS. Foot-to-foot bioelectrical impedance analysis: a valuable tool for the measurement of body composition in children. *Int J Obes Relat Metab Disord*. 2001;25(2):273–8.
- 27. Genton L, Hans D, Kyle UG, Pichard C. Dual-energy X-ray absorptiometry and body composition: differences between devices and comparison with reference methods. *Nutrition*. 2002;**18**(1):66–70.
- Earthman CP, Matthie JR, Reid PM, Harper IT, Ravussin E, Howell WH. A comparison of bioimpedance methods for detection of body cell mass change in HIV infection. J Appl Physiol (1985). 2000;88(3):944–56.
- 29. Espinosa Cuevas MA, Navarrete Rodriguez G, Villeda Martinez ME, Atilano Carsi X, Miranda Alatriste P, Tostado Gutierrez T, et al. Body fluid volume and nutritional status in hemodialysis: vector bioelectric impedance analysis. *Clin Nephrol.* 2010;**73**(4):300-8.
- Lukaski HC, Bolonchuk WW. Estimation of body fluid volumes using tetrapolar bioelectrical impedance measurements. *Aviat* Space Environ Med. 1988;59(12):1163–9.
- Segal KR, Burastero S, Chun A, Coronel P, Pierson RJ, Wang J. Estimation of extracellular and total body water by multiple-frequency bioelectrical-impedance measurement. *Am J Clin Nutr.* 1991;54(1):26–9.
- Greenbaum LA, Mesrobian HG. Vesicoureteral reflux. Pediatr Clin North Am. 2006;53(3):413–27.

www.SID ir